

ISOMETRIC IMMERSIONS INTO 3-DIMENSIONAL HOMOGENEOUS MANIFOLDS

BENOÎT DANIEL

ABSTRACT. We give a necessary and sufficient condition for a 2-dimensional Riemannian manifold to be locally isometrically immersed into a 3-dimensional homogeneous manifold with a 4-dimensional isometry group. The condition is expressed in terms of the metric, the second fundamental form, and data arising from an ambient Killing field. This class of 3-manifolds includes in particular the Berger spheres, the Heisenberg space Nil_3 , the universal cover of the Lie group $\text{PSL}_2(\mathbb{R})$ and the product spaces $\mathbb{S}^2 \times \mathbb{R}$ and $\mathbb{H}^2 \times \mathbb{R}$. We give some applications to constant mean curvature (CMC) surfaces in these manifolds; in particular we prove the existence of a generalized Lawson correspondence, i.e., a local isometric correspondence between CMC surfaces in homogeneous 3-manifolds.

1. INTRODUCTION

A classical problem in geometry is to determine whether a Riemannian manifold \mathcal{V} can be isometrically immersed in another Riemannian manifold $\bar{\mathcal{V}}$. We will restrict ourselves to the case of codimension 1 immersions, i.e., \mathcal{V} has dimension n and $\bar{\mathcal{V}}$ has dimension $n + 1$.

It is well known that the Gauss and Codazzi equations are necessary conditions relating the Riemann curvature tensor $\bar{\mathbf{R}}$ of $\bar{\mathcal{V}}$, the Riemann curvature tensor \mathbf{R} of \mathcal{V} and the shape operator \mathbf{S} of \mathcal{V} . Denoting by ∇ the Riemannian connection of \mathcal{V} , these equations are the following:

$$\langle \mathbf{R}(X, Y)Z, W \rangle - \langle \bar{\mathbf{R}}(X, Y)Z, W \rangle = \langle \mathbf{S}X, Z \rangle \langle \mathbf{S}Y, W \rangle - \langle \mathbf{S}Y, Z \rangle \langle \mathbf{S}X, W \rangle$$
$$\nabla_X \mathbf{S}Y - \nabla_Y \mathbf{S}X - \mathbf{S}[X, Y] = \bar{\mathbf{R}}(X, Y)N,$$

for all vector fields X, Y, Z and W on \mathcal{V} .

Moreover, in the case where $\bar{\mathcal{V}}$ is a space-form, i.e., the sphere \mathbb{S}^{n+1} , the Euclidean space \mathbb{R}^{n+1} or the hyperbolic space \mathbb{H}^{n+1} , the Gauss and Codazzi equations are also a sufficient condition for \mathcal{V} to be locally isometrically immersed in $\bar{\mathcal{V}}$ with \mathbf{S} as shape operator. In this case the Gauss and Codazzi equations involve only the metric and the shape operator of \mathcal{V} .

The author studied this problem when $\bar{\mathcal{V}}$ is a product manifold $\mathbb{S}^n \times \mathbb{R}$ or $\mathbb{H}^n \times \mathbb{R}$ ([Dan04]). Then the Gauss and Codazzi equations involve the metric

1991 *Mathematics Subject Classification*. Primary: 53C42. Secondary: 53A35, 53B25.

Key words and phrases. Isometric immersions, constant mean curvature surfaces, homogeneous manifolds, Gauss and Codazzi equations.

of \mathcal{V} , its shape operator S , the projection T of the vertical vector field (i.e., the unit vector field corresponding to the factor \mathbb{R}) on the tangent space of \mathcal{V} and the normal component ν of the vertical vector field (i.e., its inner product with the unit normal of \mathcal{V}). The author proved that the Gauss and Codazzi equations, together with two other compatibility equations coming from the fact that the vertical vector field is parallel, are a necessary and sufficient condition for \mathcal{V} to be locally isometrically immersed in $\tilde{\mathcal{V}}$ with S as shape operator, T as tangent projection of the vertical vector field and ν as normal component of the vertical vector field.

It is natural to try to generalize this result to other homogeneous manifolds. We will investigate the case of surfaces in manifolds of dimension 3, i.e., $n = 2$. Indeed, the classification of simply connected 3-dimensional homogeneous manifolds is well known. Such a manifold has an isometry group of dimension 3, 4 or 6. When the dimension of the isometry group is 6, then we have a space form. When the dimension of the isometry group is 3, the manifold has the geometry of the Lie group Sol_3 .

In this paper we will consider the homogeneous manifolds whose isometry groups have dimension 4: such a manifold is a Riemannian fibration over a 2-dimensional space form, the fibers are geodesics and there exists a one-parameter family of translations along the fibers, generated by a unit Killing field ξ which will be called the vertical vector field. These manifolds are classified, up to isometry, by the curvature κ of the base surface of the fibration and the bundle curvature τ , where κ and τ can be any real numbers satisfying $\kappa \neq 4\tau^2$. The bundle curvature is the number τ such $\tilde{\nabla}_X \xi = \tau X \times \xi$ for any vector field X on $\tilde{\mathcal{V}}$, where $\tilde{\nabla}$ denotes the Riemannian connection of $\tilde{\mathcal{V}}$.

When the bundle curvature τ vanishes (and then $\kappa \neq 0$), we get a product manifold $\mathbb{M}^2(\kappa) \times \mathbb{R}$ where $\mathbb{M}^2(\kappa)$ is the simply connected 2-manifold of constant curvature κ . Their isometry group has 4 connected components. The vertical vector ξ is simply the vector corresponding to the factor \mathbb{R} . This case was treated in [Dan04].

When $\tau \neq 0$, the isometry group has 2 connected components: an isometry either preserves the orientations of both the fibers and the base of the fibration, or reverses both orientations. These manifolds are of three types: they have the isometry group of the Berger spheres for $\kappa > 0$, of the Heisenberg space Nil_3 for $\kappa = 0$, and of $\widetilde{\text{PSL}_2(\mathbb{R})}$ for $\kappa < 0$. In this paper we will deal with these three types of manifold. Like for $\mathbb{M}^2(\kappa) \times \mathbb{R}$, the Gauss and Codazzi equations involve the metric of \mathcal{V} , its shape operator S , the tangential projection T of ξ and the normal component ν of ξ . Denoting by K the curvature of ds^2 , these equations become

$$K = \det S + \tau^2 + (\kappa - 4\tau^2)\nu^2,$$

$$\nabla_X S Y - \nabla_Y S X - S[X, Y] = (\kappa - 4\tau^2)\nu(\langle Y, T \rangle X - \langle X, T \rangle Y)$$

The first theorem is the following one.

Theorem (theorem 4.3). *Let \mathcal{V} be a simply connected oriented Riemannian manifold of dimension 2, ds^2 its metric (which we also denote by $\langle \cdot, \cdot \rangle$), ∇ its Riemannian connection and J the rotation of angle $\frac{\pi}{2}$ on $T\mathcal{V}$. Let S be a field of symmetric operators $S_y : T_y\mathcal{V} \rightarrow T_y\mathcal{V}$, T a vector field on \mathcal{V} and ν a smooth function on \mathcal{V} such that $\|T\|^2 + \nu^2 = 1$.*

Let \mathbb{E} be a 3-dimensional homogeneous manifold with a 4-dimensional isometry group and ξ its vertical vector field. Let κ be its base curvature and τ its bundle curvature. Then there exists an isometric immersion $f : \mathcal{V} \rightarrow \mathbb{E}$ such that the shape operator with respect to the normal N associated to f is

$$df \circ S \circ df^{-1}$$

and such that

$$\xi = df(T) + \nu N$$

if and only if (ds^2, S, T, ν) satisfies the Gauss and Codazzi equations for \mathbb{E} and, for all vector fields X on \mathcal{V} , the following equations:

$$\nabla_X T = \nu(SX - \tau JX), \quad d\nu(X) + \langle SX - \tau JX, T \rangle = 0.$$

In this case, the immersion is unique up to a global isometry of \mathbb{E} preserving the orientations of both the fibers and the base of the fibration.

The two additional conditions come from the fact that $\bar{\nabla}_X \xi = \tau X \times \xi$ for all vector fields X .

We notice that this theorem seems specific to dimension 2, since the operator of rotation J is involved.

The method to prove this theorem is similar to that of [Dan04] and was inspired by that of Tenenblat ([Ten71]): it is based on differential forms, moving frames and integrable distributions. However, things are technically much more complicated here: in [Dan04] the proof was simplified by the fact that $\mathbb{S}^n \times \mathbb{R}$ and $\mathbb{H}^n \times \mathbb{R}$ can be included in \mathbb{R}^{n+2} and in the Lorentz space \mathbb{L}^{n+2} respectively. We will first present the models used for the 3-dimensional homogeneous manifolds, and then we will prove the theorem.

Finally we will give two applications of the main theorem to constant mean curvature (CMC) surfaces in 3-dimensional homogeneous manifolds with 4-dimensional isometry group.

The first application is the existence of an isometric correspondence between certain CMC surfaces in homogeneous 3-manifolds with the same anisotropy coefficient $\kappa - 4\tau^2$. This correspondence generalizes the classical Lawson correspondence between certain CMC surfaces in space-forms. This is the following theorem.

Theorem (see theorem 5.2). *Let \mathbb{E}_1 and \mathbb{E}_2 be two 3-dimensional homogeneous manifolds with 4-dimensional isometry groups, of base curvatures κ_1 and κ_2 and bundle curvatures τ_1 and τ_2 respectively, and such that*

$$\kappa_1 - 4\tau_1^2 = \kappa_2 - 4\tau_2^2.$$

Let H_1 and H_2 be two real numbers such that

$$\tau_1^2 + H_1^2 = \tau_2^2 + H_2^2.$$

Then there exists an isometric correspondence between simply connected CMC H_1 surfaces in \mathbb{E}_1 and simply connected CMC H_2 surfaces in \mathbb{E}_2 .

This correspondence is called the correspondence of the sister surfaces.

The second application is the existence of “twin immersions” of non-minimal CMC immersions in homogeneous 3-manifolds with non-vanishing bundle curvature. This twin immersion might be useful to prove an Alexandrov-type theorem in these manifolds.

Conventions and notations. In this paper we will use the following index conventions: Latin letters i, j , etc, denote integers between 1 and n (or the integers 1 and 2), Greek letters α, β , etc, denote integers between 1 and $n + 1$ (or between 1 and 3).

The set of vector fields on a Riemannian manifold \mathcal{V} will be denoted by $\mathfrak{X}(\mathcal{V})$.

The Riemann curvature tensor R of a Riemannian manifold \mathcal{V} of Riemannian connection ∇ is defined using the following convention:

$$R(X, Y)Z = \nabla_Y \nabla_X Z - \nabla_X \nabla_Y Z + \nabla_{[X, Y]} Z.$$

The shape operator of a hypersurface \mathcal{V} of a Riemannian manifold $\bar{\mathcal{V}}$ associated to its unit normal N is

$$SX = -\bar{\nabla}_X N$$

where $\bar{\nabla}$ is the Riemannian connection of $\bar{\mathcal{V}}$.

2. 3-DIMENSIONAL HOMOGENEOUS MANIFOLDS WITH 4-DIMENSIONAL ISOMETRY GROUP

In this section we will give the general setting for simply connected homogeneous 3-manifolds with 4-dimensional isometry group and we will describe the models used. We will consider only those having non-vanishing bundle curvature (since the product manifolds $\mathbb{M}^2(\kappa) \times \mathbb{R}$ were treated in [Dan04]). The reader can refer to [Sco83] for the geometry of 3-dimensional homogeneous manifolds.

2.1. Canonical frame. Let \mathbb{E} be a simply connected 3-dimensional homogeneous manifold with a 4-dimensional isometry group. Such a manifold is a Riemannian fibration over a simply connected 2-manifold of constant curvature κ . The fibers are geodesics. We will denote by ξ a unit vector field on \mathbb{E} tangent to the fibers; it will be called the vertical vector field. It is a Killing field (corresponding to translations along the fibers).

We will denote by $\bar{\nabla}$ and \bar{R} the Riemannian connection and the Riemannian curvature tensor of \mathbb{E} respectively.

We assume that \mathbb{E} is not a product manifold $\mathbb{M}^2(\kappa) \times \mathbb{R}$.

The manifold \mathbb{E} locally has a direct orthonormal frame (E_1, E_2, E_3) with

$$E_3 = \xi$$

whose non-vanishing Christoffel symbols $\bar{\Gamma}_{\beta\gamma}^\alpha = \langle \nabla_{E_\alpha} E_\beta, E_\gamma \rangle$ are the following:

$$\begin{aligned}\bar{\Gamma}_{12}^3 &= \bar{\Gamma}_{23}^1 = -\bar{\Gamma}_{21}^3 = -\bar{\Gamma}_{13}^2 = \tau, \\ \bar{\Gamma}_{32}^1 &= -\bar{\Gamma}_{31}^2 = \tau - \sigma,\end{aligned}$$

for some real numbers σ and $\tau \neq 0$ (this will be explicated in the sequel). Then we have

$$[E_1, E_2] = 2\tau E_3, \quad [E_2, E_3] = \sigma E_1, \quad [E_3, E_1] = \sigma E_2.$$

We will call (E_1, E_2, E_3) the canonical frame of \mathbb{E} . For all vector field X we have

$$\bar{\nabla}_X E_3 = \tau X \times E_3$$

where \times denotes the vector product in \mathbb{E} , i.e., for all vector fields X, Y, Z , $\langle X \times Y, Z \rangle = \det_{(E_1, E_2, E_3)}(X, Y, Z)$.

Setting

$$\langle \bar{\mathbf{R}}(X \wedge Y), Z \wedge W \rangle = \langle \bar{\mathbf{R}}(X, Y)Z, W \rangle,$$

the matrix of $\bar{\mathbf{R}}$ in the basis $(E_2 \wedge E_3, E_3 \wedge E_1, E_1 \wedge E_2)$ is

$$\bar{\mathbf{R}} = \text{diag}(a, a, b)$$

with

$$a = \tau^2, \quad b = -3\tau^2 + 2\sigma\tau.$$

We now compute the curvature κ of the base of the fibration. If $\bar{M} \rightarrow M$ is a Riemannian submersion, then the sectional curvature of a 2-plane Π in M generated by an orthonormal pair (X, Y) is

$$K(\Pi) = \bar{K}(\bar{\Pi}) + \frac{3}{4} \|[\bar{X}, \bar{Y}]^\vee\|^2$$

where \bar{X} and \bar{Y} are horizontal lifts of X and Y in \bar{M} , $\bar{K}(\bar{\Pi})$ is the sectional curvature of a 2-plane $\bar{\Pi}$ in \bar{M} generated by (\bar{X}, \bar{Y}) , and where Z^\vee denotes the vertical part of a vector field Z in \bar{M} (see [Car92], chapter 8). In our case we get

$$\kappa = \langle \bar{\mathbf{R}}(E_1, E_2)E_1, E_2 \rangle + \frac{3}{4} \|[E_1, E_2]^\vee\|^2 = b + \frac{3}{4} \|2\tau E_3^\vee\|^2 = b + 3\tau^2.$$

Thus we have $b = \kappa - 3\tau^2$, and so

$$\sigma = \frac{\kappa}{2\tau}.$$

Proposition 2.1. *For all vector fields X, Y, Z, W on \mathbb{E} we have*

$$\langle \bar{\mathbf{R}}(X, Y)Z, W \rangle = (\kappa - 3\tau^2) \langle \mathbf{R}_0(X, Y)Z, W \rangle + (\kappa - 4\tau^2) \langle \mathbf{R}_1(\xi; X, Y)Z, W \rangle$$

with

$$\mathbf{R}_0(X, Y)Z = \langle X, Z \rangle Y - \langle Y, Z \rangle X,$$

$$\begin{aligned} R_1(V; X, Y)Z &= \langle Y, V \rangle \langle Z, V \rangle X + \langle Y, Z \rangle \langle X, V \rangle V \\ &\quad - \langle X, Z \rangle \langle Y, V \rangle V - \langle X, V \rangle \langle Z, V \rangle Y. \end{aligned}$$

Proof. We set $X = \tilde{X} + x\xi$ with \tilde{X} horizontal and $x = \langle X, \xi \rangle$, etc. Using the multilinearity of the Riemann curvature tensor, we get a sum of 16 terms; the terms where ξ appears three or four times, or twice at positions 1, 2 or 3, 4, vanish by antisymmetry. The terms where ξ appears once vanish because the matrix of \bar{R} in the basis $(E_2 \wedge E_3, E_3 \wedge E_1, E_1 \wedge E_2)$ is diagonal. Hence we have

$$\begin{aligned} \langle \bar{R}(X, Y)Z, W \rangle &= \langle \bar{R}(\tilde{X}, \tilde{Y})\tilde{Z}, \tilde{W} \rangle \\ &\quad + yw \langle \bar{R}(\tilde{X}, \xi)\tilde{Z}, \xi \rangle + yz \langle \bar{R}(\tilde{X}, \xi)\xi, \tilde{W} \rangle \\ &\quad + xw \langle \bar{R}(\xi, \tilde{Y})\tilde{Z}, \xi \rangle + xz \langle \bar{R}(\xi, \tilde{Y})\xi, \tilde{W} \rangle \\ &= (\kappa - 3\tau^2)(\langle \tilde{X}, \tilde{Z} \rangle \langle \tilde{Y}, \tilde{W} \rangle - \langle \tilde{X}, \tilde{W} \rangle \langle \tilde{Y}, \tilde{Z} \rangle) \\ &\quad + \tau^2(yw \langle \tilde{X}, \tilde{Z} \rangle - yz \langle \tilde{X}, \tilde{W} \rangle - xw \langle \tilde{Y}, \tilde{Z} \rangle + xz \langle \tilde{Y}, \tilde{W} \rangle) \\ &= (\kappa - 3\tau^2)(\langle X, Z \rangle \langle Y, W \rangle - \langle X, W \rangle \langle Y, Z \rangle) \\ &\quad - (\kappa - 4\tau^2)(\langle X, Z \rangle \langle Y, \xi \rangle \langle W, \xi \rangle + \langle Y, W \rangle \langle X, \xi \rangle \langle Z, \xi \rangle \\ &\quad - \langle X, W \rangle \langle Y, \xi \rangle \langle Z, \xi \rangle - \langle Y, Z \rangle \langle X, \xi \rangle \langle W, \xi \rangle). \end{aligned}$$

□

2.2. The manifolds with the isometry group of the Berger spheres.

They occur when $\tau \neq 0$ and $\kappa > 0$; they are fibrations over round 2-spheres. They are obtained by deforming the metric of a round sphere in a way preserving the Hopf fibration but modifying the length of the fibers. Their isometry group is included in that of the round sphere. The reader can refer to [Pet98].

The sphere \mathbb{S}^3 is the universal covering of $\text{SO}_3(\mathbb{R})$, which can be identified with the unitary tangent bundle to the 2-sphere US^2 . Indeed, the group $\text{SO}_3(\mathbb{R})$ acts transitively on US^2 , and the stabilizer of any point in US^2 is trivial. The unitary tangent bundle US^2 can be endowed with the metric induced by the standard metric on the tangent bundle TS^2 . We will give an expression of this metric.

Let $(x, y) \mapsto \varphi(x, y)$ be a conformal parametrization of a domain D in \mathbb{S}^2 and let λ be the conformal factor, i.e., the metric of D is given by $\lambda^2(dx^2 + dy^2)$. Then a parametrization of UD is the following:

$$(x, y, \theta) \mapsto \left(\varphi(x, y), \frac{1}{\lambda}(\cos \theta \partial_x + \sin \theta \partial_y) \right).$$

Let $p = \varphi(x, y) \in D$, $v \in \text{T}_p D$ and $V \in \text{T}_{(p,v)}(UD)$. Let $\alpha(t) = (p(t), v(t))$ be a curve such that $v(t) \in \text{T}_{p(t)}\mathbb{H}^2$, $p(0) = p$, $v(0) = v$ and $\alpha'(0) = V$. Then the norm of V is given by

$$\|V\|_{(p,v)}^2 = \|d\pi(V)\|_p^2 + \left\| \frac{Dv}{dt}(0) \right\|_p^2$$

where $\pi : UD \rightarrow D$ is the canonical projection.

We set $\alpha(t) = (x(t), y(t), \theta(t))$. Then we have

$$v(t) = \frac{1}{\lambda}(\cos \theta(t)\partial_x + \sin \theta(t)\partial_y),$$

and thus

$$\begin{aligned} \frac{Dv}{dt} &= -\frac{\dot{\lambda}}{\lambda^2}(\cos \theta \partial_x + \sin \theta \partial_y) + \frac{\dot{\theta}}{\lambda}(-\sin \theta \partial_x + \cos \theta \partial_y) \\ &\quad + \frac{1}{\lambda}(\cos \theta(\dot{x}\nabla_{\partial_x}\partial_x + \dot{y}\nabla_{\partial_y}\partial_x) + \sin \theta(\dot{x}\nabla_{\partial_x}\partial_y + \dot{y}\nabla_{\partial_y}\partial_y)), \end{aligned}$$

where the dot denotes the derivation with respect to t . Since $\dot{\lambda} = \dot{x}\lambda_x + \dot{y}\lambda_y$, $\nabla_{\partial_x}\partial_x = \frac{\lambda_x}{\lambda}\partial_x - \frac{\lambda_y}{\lambda}\partial_y$, $\nabla_{\partial_y}\partial_y = -\frac{\lambda_x}{\lambda}\partial_x + \frac{\lambda_y}{\lambda}\partial_y$ and $\nabla_{\partial_x}\partial_y = \nabla_{\partial_y}\partial_x = \frac{\lambda_y}{\lambda}\partial_x + \frac{\lambda_x}{\lambda}\partial_y$, we get

$$\frac{Dv}{dt} = \frac{1}{\lambda^2}(\lambda\dot{\theta} + \dot{y}\lambda_x - \dot{x}\lambda_y)(\cos \theta \partial_y - \sin \theta \partial_x).$$

Thus

$$\|V\|_{(p,v)}^2 = \lambda^2(\dot{x}^2 + \dot{y}^2) + \frac{1}{\lambda^2}(\lambda\dot{\theta} + \dot{y}\lambda_x - \dot{x}\lambda_y)^2.$$

Setting $z = \theta$ on the universal covering, we get the following expression for the metric of \widetilde{UD} :

$$ds^2 = \lambda^2(dx^2 + dy^2) + \left(-\frac{\lambda_y}{\lambda}dx + \frac{\lambda_x}{\lambda}dy + dz\right)^2.$$

We now choose $D = \mathbb{S}^2 \setminus \{\infty\}$ with the metric of constant curvature 4 (i.e., the metric of the round sphere of radius $\frac{1}{2}$) given by the stereographic projection, i.e.,

$$\lambda = \frac{1}{1 + x^2 + y^2}.$$

Then we get

$$ds^2 = \lambda^2(dx^2 + dy^2) + (2\lambda(ydx - xdy) + dz)^2.$$

More generally, \mathbb{R}^3 endowed with the metric

$$ds^2 = \lambda^2(dx^2 + dy^2) + (\tau\lambda(ydx - xdy) + dz)^2$$

with

$$\lambda = \frac{1}{1 + \frac{\kappa}{4}(x^2 + y^2)}$$

is the universal cover of a homogeneous manifold \mathbb{E} of bundle curvature τ and of base curvature $\kappa > 0$ minus the fiber corresponding to the point $\infty \in \mathbb{S}^2$. The fibers are given by $\{x = x_0, y = y_0\}$ in these coordinates. The canonical frame is (E_1, E_2, E_3) with

$$\begin{aligned} (1) \quad E_1 &= \lambda^{-1}(\cos(\sigma z)\partial_x + \sin(\sigma z)\partial_y) + \tau(x \sin(\sigma z) - y \cos(\sigma z))\partial_z, \\ E_2 &= \lambda^{-1}(-\sin(\sigma z)\partial_x + \cos(\sigma z)\partial_y) + \tau(x \cos(\sigma z) + y \sin(\sigma z))\partial_z, \\ E_3 &= \partial_z \end{aligned}$$

with

$$\sigma = \frac{\kappa}{2\tau},$$

which satisfy

$$[E_1, E_2] = 2\tau E_3, \quad [E_2, E_3] = \frac{\kappa}{2\tau} E_1, \quad [E_3, E_1] = \frac{\kappa}{2\tau} E_2.$$

This frame is defined on the open set \mathbb{E}' which is \mathbb{E} minus the fiber corresponding to the point $\infty \in \mathbb{S}^2$.

The Berger spheres in the strict sense are the manifolds such that $\kappa = 4$.

2.3. The manifolds with the isometry group of the Heisenberg space Nil_3 . They occur when $\tau \neq 0$ and $\kappa = 0$; they are fibrations over the Euclidean plane.

The Heisenberg space is the Lie group

$$\text{Nil}_3 = \left\{ \left(\begin{array}{ccc} 1 & a & c \\ 0 & 1 & b \\ 0 & 0 & 1 \end{array} \right); (a, b, c) \in \mathbb{R}^3 \right\}$$

endowed with a left invariant metric.

It is useful to use exponential coordinates. In this model, the Heisenberg space Nil_3 is \mathbb{R}^3 endowed with the following metric:

$$ds^2 = dx^2 + dy^2 + (\tau(ydx - xdy) + dz)^2.$$

The fibers are given by $\{x = x_0, y = y_0\}$ in these coordinates.

The canonical frame is (E_1, E_2, E_3) with

$$(2) \quad E_1 = \partial_x - \tau y \partial_z, \quad E_2 = \partial_y + \tau x \partial_z, \quad E_3 = \partial_z,$$

which satisfy

$$[E_1, E_2] = 2\tau E_3, \quad [E_2, E_3] = 0, \quad [E_3, E_1] = 0.$$

The reader can refer to [FMP99] (where $\tau = \frac{1}{2}$).

2.4. The manifolds with the isometry group of $\widetilde{\text{PSL}}_2(\mathbb{R})$. They occur when $\tau \neq 0$ and $\kappa < 0$; they are fibrations over hyperbolic planes.

The Lie group $\widetilde{\text{PSL}}_2(\mathbb{R})$ with its standard metric can be identified with the universal covering of the unitary tangent bundle to the hyperbolic plane UH^2 equipped with its canonical metric. Indeed, the group $\text{PSL}_2(\mathbb{R})$ acts transitively on UH^2 , and the stabilizer of any point in UH^2 is trivial. The unitary tangent bundle UH^2 can be endowed with the metric induced by the standard metric on the tangent bundle TH^2 . The reader can refer to [Sco83]. We will give an expression of this metric.

Let $(x, y) \mapsto \varphi(x, y)$ be a conformal parametrization of \mathbb{H}^2 and let λ be the conformal factor, i.e., the metric of \mathbb{H}^2 is given by $\lambda^2(dx^2 + dy^2)$. Then, proceeding as in section 2.2, we obtain that a metric on $\widetilde{\text{PSL}}_2(\mathbb{R})$ is

$$ds^2 = \lambda^2(dx^2 + dy^2) + \left(-\frac{\lambda_y}{\lambda} dx + \frac{\lambda_x}{\lambda} dy + dz \right)^2.$$

This metric defines a homogeneous manifold with $\kappa = -1$ and $\tau = -\frac{1}{2}$.

More generally, we can take the Poincaré disk model for the hyperbolic plane of constant curvature $\kappa < 0$. The manifold $\mathbb{D}^2\left(\frac{2}{\sqrt{-\kappa}}\right) \times \mathbb{R}$, where $\mathbb{D}^2(\rho) = \{(x, y) \in \mathbb{R}^2; x^2 + y^2 < \rho^2\}$, endowed with the metric

$$ds^2 = \lambda^2(dx^2 + dy^2) + (\tau\lambda(ydx - xdy) + dz)^2$$

with

$$\lambda = \frac{1}{1 + \frac{\kappa}{4}(x^2 + y^2)}$$

is a homogeneous manifold of bundle curvature τ and of base curvature $\kappa < 0$. The fibers are given by $\{x = x_0, y = y_0\}$ in these coordinates. The canonical frame is (E_1, E_2, E_3) with

$$(3) \quad \begin{aligned} E_1 &= \lambda^{-1}(\cos(\sigma z)\partial_x + \sin(\sigma z)\partial_y) + \tau(x\sin(\sigma z) - y\cos(\sigma z))\partial_z, \\ E_2 &= \lambda^{-1}(-\sin(\sigma z)\partial_x + \cos(\sigma z)\partial_y) + \tau(x\cos(\sigma z) + y\sin(\sigma z))\partial_z, \\ E_3 &= \partial_z \end{aligned}$$

with

$$\sigma = \frac{\kappa}{2\tau},$$

which satisfy

$$[E_1, E_2] = 2\tau E_3, \quad [E_2, E_3] = \frac{\kappa}{2\tau} E_1, \quad [E_3, E_1] = \frac{\kappa}{2\tau} E_2.$$

3. PRELIMINARIES

3.1. The compatibility equations for surfaces in 3-dimensional homogeneous manifolds. We consider a 3-dimensional homogeneous manifold \mathbb{E} with an isometry group of dimension 4, of bundle curvature τ and of base curvature κ . Let \bar{R} be the Riemann curvature tensor of \mathbb{E} . Let \mathcal{V} be an oriented surface in \mathbb{E} , ∇ the Riemannian connection of \mathcal{V} , J the rotation of angle $\frac{\pi}{2}$ on $T\mathcal{V}$, N the unit normal to \mathcal{V} and S the shape operator of \mathcal{V} .

Proposition 3.1. *For $X, Y, Z, W \in \mathfrak{X}(\mathcal{V})$ we have*

$$\langle \bar{R}(X, Y)Z, W \rangle = (\kappa - 3\tau^2)\langle R_0(X, Y)Z, W \rangle + (\kappa - 4\tau^2)\langle R_1(T; X, Y)Z, W \rangle,$$

$$\bar{R}(X, Y)N = (\kappa - 4\tau^2)\nu(\langle Y, T \rangle X - \langle X, T \rangle Y),$$

where

$$\nu = \langle N, \xi \rangle,$$

T is the projection of ξ on $T\mathcal{V}$, i.e.,

$$T = \xi - \nu N,$$

and R_0 and R_1 are as in proposition 2.1.

Proof. This is a consequence of proposition 2.1, using the fact that X, Y and Z are tangent to the surface and N is normal to the surface. \square

Corollary 3.2. *The Gauss and Codazzi equations in \mathbb{E} are*

$$K = \det S + \tau^2 + (\kappa - 4\tau^2)\nu^2,$$

$$\nabla_X SY - \nabla_Y SX - S[X, Y] = (\kappa - 4\tau^2)\nu(\langle Y, T \rangle X - \langle X, T \rangle Y),$$

where K is the Gauss curvature of \mathcal{V} .

Proposition 3.3. *For $X \in \mathfrak{X}(\mathcal{V})$ we have*

$$\nabla_X T = \nu(SX - \tau JX), \quad d\nu(X) + \langle SX - \tau JX, T \rangle = 0.$$

Proof. On the one hand we have

$$\begin{aligned} \bar{\nabla}_X \xi &= \bar{\nabla}_X(T + \nu N) \\ &= \bar{\nabla}_X T + d\nu(X)N + \nu \bar{\nabla}_X N \\ &= \nabla_X T + \langle SX, T \rangle N + d\nu(X)N - \nu SX. \end{aligned}$$

On the other hand we have

$$\begin{aligned} \bar{\nabla}_X \xi &= \tau X \times \xi \\ &= \tau X \times (T + \nu N) \\ &= \tau(\langle JX, T \rangle N - \nu JX). \end{aligned}$$

We conclude taking the tangential and normal parts in both expressions. \square

3.2. Moving frames. In this section we introduce some material about the technique of moving frames.

Let \mathcal{V} be a Riemannian manifold of dimension n , ∇ its Levi-Civita connection, and R the Riemannian curvature tensor. Let S be a field of symmetric operators $S_y : T_y \mathcal{V} \rightarrow T_y \mathcal{V}$. Let (e_1, \dots, e_n) be a local orthonormal frame on \mathcal{V} and $(\omega^1, \dots, \omega^n)$ the dual basis of (e_1, \dots, e_n) , i.e.,

$$\omega^i(e_k) = \delta_k^i.$$

We also set

$$\omega^{n+1} = 0.$$

We define the forms $\omega_j^i, \omega_j^{n+1}, \omega_{n+1}^i$ and ω_{n+1}^{n+1} on \mathcal{V} by

$$\omega_j^i(e_k) = \langle \nabla_{e_k} e_j, e_i \rangle, \quad \omega_j^{n+1}(e_k) = \langle S e_k, e_j \rangle,$$

$$\omega_{n+1}^j = -\omega_j^{n+1}, \quad \omega_{n+1}^{n+1} = 0.$$

Then we have

$$\nabla_{e_k} e_j = \sum_i \omega_j^i(e_k) e_i, \quad S e_k = \sum_j \omega_j^{n+1}(e_k) e_j.$$

Finally we set $R_{klj}^i = \langle R(e_k, e_l) e_j, e_i \rangle$.

Proposition 3.4. *We have the following formulas:*

$$(4) \quad d\omega^i + \sum_p \omega_p^i \wedge \omega^p = 0,$$

$$(5) \quad \sum_p \omega_p^{n+1} \wedge \omega^p = 0,$$

$$(6) \quad d\omega_j^i + \sum_p \omega_p^i \wedge \omega_j^p = -\frac{1}{2} \sum_k \sum_l R_{klij}^i \omega^k \wedge \omega^l,$$

$$(7) \quad d\omega_j^{n+1} + \sum_p \omega_p^{n+1} \wedge \omega_j^p = \frac{1}{2} \sum_k \sum_l \langle \nabla_{e_k} S e_l - \nabla_{e_l} S e_k - S[e_k, e_l], e_j \rangle \omega^k \wedge \omega^l.$$

For a proof of these classical formulas, the reader can refer to [Dan04], proposition 2.4.

3.3. Some facts about hypersurfaces. In this section we consider an orientable hypersurface \mathcal{V} of an $(n+1)$ -dimensional Riemannian manifold $\bar{\mathcal{V}}$. Let (e_1, \dots, e_n) be a local orthonormal frame on \mathcal{V} , e_{n+1} the normal to \mathcal{V} , and (E_1, \dots, E_{n+1}) a local orthonormal frame on $\bar{\mathcal{V}}$. We denote by ∇ and $\bar{\nabla}$ the Riemannian connections on \mathcal{V} and $\bar{\mathcal{V}}$ respectively, and by S the shape operator of \mathcal{V} (with respect to the normal e_{n+1}). We define the forms $\omega^\alpha, \omega_\beta^\alpha$ on \mathcal{V} as in section 3.2. Then we have

$$\bar{\nabla}_{e_k} e_\beta = \sum_\gamma \omega_\beta^\gamma(e_k) e_\gamma.$$

Let $A \in \text{SO}_{n+1}(\mathbb{R})$ be the matrix whose columns are the coordinates of the e_β in the frame (E_α) , namely $A_\beta^\alpha = \langle e_\beta, E_\alpha \rangle$. Let $\Omega = (\omega_\beta^\alpha) \in \mathcal{M}_{n+1}(\mathbb{R})$.

Lemma 3.5. *The matrix A satisfies the following equation:*

$$A^{-1}dA = \Omega + L(A)$$

with

$$L(A)_\beta^\alpha = \sum_k \left(\sum_{\gamma, \delta, \varepsilon} A_\alpha^\varepsilon A_k^\gamma A_\beta^\delta \bar{\Gamma}_{\gamma\varepsilon}^\delta \right) \omega^k,$$

where the $\bar{\Gamma}_{\gamma\varepsilon}^\delta$ are the Christoffel symbols of the frame (E_α) .

Proof. We have

$$e_\beta = \sum_\alpha A_\beta^\alpha E_\alpha.$$

Then, on the one hand we have

$$\begin{aligned} \bar{\nabla}_{e_k} e_\beta &= \sum_\delta dA_\beta^\delta(e_k) E_\delta + \sum_\delta A_\beta^\delta \bar{\nabla}_{e_k} E_\delta \\ &= \sum_\varepsilon dA_\beta^\varepsilon(e_k) E_\delta + \sum_\gamma \sum_\delta \sum_\varepsilon A_\beta^\delta A_k^\gamma \bar{\Gamma}_{\gamma\delta}^\varepsilon E_\varepsilon, \end{aligned}$$

and on the other hand we have

$$\bar{\nabla}_{e_k} e_\beta = \sum_{\gamma} \sum_{\varepsilon} \omega_{\beta}^{\gamma}(e_k) A_{\gamma}^{\varepsilon} E_{\varepsilon}.$$

Identifying the coefficients we get

$$\begin{aligned} dA_{\beta}^{\varepsilon}(e_k) &= - \sum_{\gamma} \sum_{\delta} A_{\beta}^{\delta} A_k^{\gamma} \bar{\Gamma}_{\gamma\delta}^{\varepsilon} + \sum_{\gamma} \omega_{\beta}^{\gamma}(e_k) A_{\gamma}^{\varepsilon} \\ &= \sum_{\gamma} \sum_{\delta} A_{\beta}^{\delta} A_k^{\gamma} \bar{\Gamma}_{\gamma\varepsilon}^{\delta} + \sum_{\gamma} \omega_{\beta}^{\gamma}(e_k) A_{\gamma}^{\varepsilon} \end{aligned}$$

since the frame (E_{α}) is orthonormal.

We conclude using the fact that A^{-1} is the transpose of A . \square

4. ISOMETRIC IMMERSIONS OF SURFACES INTO 3-DIMENSIONAL HOMOGENEOUS MANIFOLDS

We consider a simply connected oriented Riemannian manifold \mathcal{V} of dimension 2. Let ds^2 be the metric on \mathcal{V} (we will also denote it by $\langle \cdot, \cdot \rangle$), ∇ the Riemannian connection of \mathcal{V} , R its Riemann curvature tensor and J the rotation of angle $\frac{\pi}{2}$ on $T\mathcal{V}$. Let S be a field of symmetric operators $S_y : T_y\mathcal{V} \rightarrow T_y\mathcal{V}$, T a vector field on \mathcal{V} such that $\|T\| \leq 1$ and ν a smooth function on \mathcal{V} such that $\nu^2 \leq 1$.

The compatibility equations for surfaces in 3-dimensional homogeneous manifolds with 4-dimensional isometry group established in section 3.1 suggest to introduce the following definition.

Definition 4.1. Let \mathbb{E} be a 3-dimensional homogeneous manifold with a 4-dimensional isometry group. Let κ be its base curvature and τ its bundle curvature. We say that (ds^2, S, T, ν) satisfies the compatibility equations for \mathbb{E} if

$$\|T\|^2 + \nu^2 = 1$$

and, for all $X, Y, Z \in \mathfrak{X}(\mathcal{V})$,

$$(8) \quad K = \det S + \tau^2 + (\kappa - 4\tau^2)\nu^2,$$

$$(9) \quad \nabla_X SY - \nabla_Y SX - S[X, Y] = (\kappa - 4\tau^2)\nu(\langle Y, T \rangle X - \langle X, T \rangle Y),$$

$$(10) \quad \nabla_X T = \nu(SX - \tau JX),$$

$$(11) \quad d\nu(X) + \langle SX - \tau JX, T \rangle = 0.$$

Remark 4.2. We notice that (10) implies (11) except when $\nu = 0$ (by differentiating the identity $\langle T, T \rangle + \nu^2 = 1$ with respect to X).

Theorem 4.3. *Let \mathcal{V} be a simply connected oriented Riemannian manifold of dimension 2, ds^2 its metric and ∇ its Riemannian connection. Let S be a field of symmetric operators $S_y : T_y\mathcal{V} \rightarrow T_y\mathcal{V}$, T a vector field on \mathcal{V} and ν a smooth function on \mathcal{V} such that $\|T\|^2 + \nu^2 = 1$.*

Let \mathbb{E} be a 3-dimensional homogeneous manifold with a 4-dimensional isometry group and ξ its vertical vector field. Let κ be its base curvature and τ its bundle curvature. Then there exists an isometric immersion $f : \mathcal{V} \rightarrow \mathbb{E}$ such that the shape operator with respect to the normal N associated to f is

$$df \circ S \circ df^{-1}$$

and such that

$$\xi = df(T) + \nu N$$

if and only if (ds^2, S, T, ν) satisfies the compatibility equations for \mathbb{E} . In this case, the immersion is unique up to a global isometry of \mathbb{E} preserving the orientations of both the fibers and the base of the fibration.

The fact that the compatibility equations are necessary was proved in section 3.1. To prove that they are sufficient, we consider a local orthonormal frame (e_1, e_2) on \mathcal{V} and the forms $\omega^i, \omega^3, \omega_j^i, \omega_j^3, \omega_3^i$ and ω_3^3 as in section 3.2 (with $n = 2$).

From now on we assume that $\tau \neq 0$ since the case $\tau = 0$ was treated in [Dan04].

We denote by (E_1, E_2, E_3) the canonical frame of \mathbb{E} (see section 2.1); in particular we have $E_3 = \xi$. We denote by \mathbb{E}' the open set where the canonical frame is defined (in particular we have $\mathbb{E}' = \mathbb{E}$ when $\kappa = 0$ or $\kappa < 0$; see sections 2.2, 2.3 and 2.4).

We set

$$T^k = \langle T, e_k \rangle, \quad T^3 = \nu.$$

We define the one-form η on \mathcal{V} by

$$\eta(X) = \langle T, X \rangle.$$

In the frame (e_1, e_2) we have $\eta = \sum_k T^k \omega^k$. We define the following matrix of one-forms:

$$\Omega = (\omega_{\beta}^{\alpha}) \in \mathcal{M}_3(\mathbb{R}).$$

For $Z \in \text{SO}_3(\mathbb{R})$, we set

$$L(Z)_{\beta}^{\alpha} = \sum_k \left(\sum_{\gamma, \delta, \varepsilon} Z_{\alpha}^{\varepsilon} Z_k^{\gamma} Z_{\beta}^{\delta} \bar{\Gamma}_{\gamma \varepsilon}^{\delta} \right) \omega^k,$$

where the $\bar{\Gamma}_{\gamma \varepsilon}^{\delta}$ are the Christoffel symbols of the frame (E_{α}) (see section 3.3). This defines an antisymmetric matrix of 1-forms.

We also set $\sigma = \frac{\kappa}{2\tau}$.

From now on we assume that the hypotheses of theorem 4.3 are satisfied. We first prove some technical lemmas that are consequences of the compatibility equations.

Lemma 4.4. *We have*

$$d\eta = -2\tau\nu\omega^1 \wedge \omega^2.$$

Proof. By (10) we have $d\eta(X, Y) = \langle \nabla_X T, Y \rangle - \langle \nabla_Y T, X \rangle = 2\tau\nu \langle X, JY \rangle$. Thus $d\eta(e_1, e_2) = -2\tau\nu$. \square

Lemma 4.5. *We have*

$$\begin{aligned} dT^1 &= \sum_{\gamma} T^{\gamma} \omega_1^{\gamma} + \tau T^3 \omega^2, \\ dT^2 &= \sum_{\gamma} T^{\gamma} \omega_2^{\gamma} - \tau T^3 \omega^1, \\ dT^3 &= \sum_{\gamma} T^{\gamma} \omega_3^{\gamma} - \tau T^1 \omega^2 + \tau T^2 \omega^1. \end{aligned}$$

Proof. The first two identities are a consequence of condition (10) and the last one of condition (11). \square

Lemma 4.6. *We have*

$$\begin{aligned} d\Omega + \Omega \wedge \Omega &= \begin{pmatrix} 0 & \tau^2 & 0 \\ -\tau^2 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \omega^1 \wedge \omega^2 \\ &+ (\kappa - 4\tau^2) T^3 \begin{pmatrix} 0 & T^3 & -T^2 \\ -T^3 & 0 & T^1 \\ T^2 & -T^1 & 0 \end{pmatrix} \omega^1 \wedge \omega^2. \end{aligned}$$

Proof. We set $\Psi = d\Omega + \Omega \wedge \Omega$ and $R_{klj}^i = \langle R(e_k, e_l)e_j, e_i \rangle$. By proposition 3.4 we have

$$\Psi_j^i = -\frac{1}{2} \sum_k \sum_l R_{klj}^i \omega^k \wedge \omega^l + \omega_3^i \wedge \omega_j^3,$$

and by the Gauss equation (8) we have $R_{klj}^i = \bar{R}_{klj}^i + \omega_j^3 \wedge \omega_i^3 \langle e_k, e_l \rangle$ with $\bar{R}_{klj}^i = (\kappa - 3\tau^2)(\delta_j^k \delta_i^l - \delta_j^l \delta_i^k) + (\kappa - 4\tau^2)(T^l T^j \delta_i^k + T^k T^i \delta_j^l - T^l T^i \delta_j^k - T^k T^j \delta_i^l)$.

Thus we get

$$\Psi_j^i = (\kappa - 3\tau^2) \omega^i \wedge \omega^j + (\kappa - 4\tau^2)(T^i \omega^j - T^j \omega^i) \wedge \eta.$$

In the same way, by proposition 3.4 we have

$$\Psi_j^3 = \frac{1}{2} \sum_k \sum_l \langle \nabla_{e_k} S e_l - \nabla_{e_l} S e_k - S[e_k, e_l], e_j \rangle \omega^k \wedge \omega^l,$$

and by the Codazzi equation (9) we have

$$\langle \nabla_{e_k} S e_l - \nabla_{e_l} S e_k - S[e_k, e_l], e_j \rangle = (\kappa - 4\tau^2) T^3 (T^l \delta_j^k - T^k \delta_j^l).$$

Thus we get

$$\Psi_j^3 = (\kappa - 4\tau^2) T^3 \omega^j \wedge \eta.$$

Hence we have

$$\begin{aligned}\Psi &= (\kappa - 3\tau^2) \begin{pmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \omega^1 \wedge \omega^2 \\ &\quad + (\kappa - 4\tau^2) \begin{pmatrix} 0 & -T^2 & -T^3 \\ T^2 & 0 & 0 \\ T^3 & 0 & 0 \end{pmatrix} \omega^1 \wedge \eta \\ &\quad + (\kappa - 4\tau^2) \begin{pmatrix} 0 & T^1 & -0 \\ -T^1 & 0 & -T^3 \\ 0 & T^3 & 0 \end{pmatrix} \omega^2 \wedge \eta.\end{aligned}$$

We conclude using that $\omega^1 \wedge \eta = T^2\omega^1 \wedge \omega^2$, $\omega^2 \wedge \eta = -T^1\omega^1 \wedge \omega^2$ and $(T^1)^2 + (T^2)^2 + (T^3)^2 = 1$. \square

Lemma 4.7. *We have*

$$\begin{aligned}L(Z) &= (2\tau - \sigma) \begin{pmatrix} 0 & -T^3 & T^2 \\ T^3 & 0 & -T^1 \\ -T^2 & T^1 & 0 \end{pmatrix} \eta \\ &\quad + \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & \tau \\ 0 & -\tau & 0 \end{pmatrix} \omega^1 + \begin{pmatrix} 0 & 0 & -\tau \\ 0 & 0 & 0 \\ \tau & 0 & 0 \end{pmatrix} \omega^2.\end{aligned}$$

Proof. We compute that

$$\begin{aligned}L(Z)_\beta^\alpha &= \sum_k \left(\sum_\gamma \sum_\delta \sum_\varepsilon Z_\alpha^\varepsilon Z_k^\gamma Z_\beta^\delta \bar{\Gamma}_{\gamma\varepsilon}^\delta \right) \omega^k \\ &= \sum_k (\tau(Z_\alpha^2 Z_k^1 Z_\beta^3 + Z_\alpha^3 Z_k^2 Z_\beta^1 - Z_\alpha^1 Z_k^2 Z_\beta^3 - Z_\alpha^3 Z_k^1 Z_\beta^2) \\ &\quad + (\tau - \sigma)(Z_\alpha^2 Z_k^3 Z_\beta^1 - Z_\alpha^1 Z_k^3 Z_\beta^2)) \omega^k \\ &= \sum_k (\tau T^\beta (Z_k^1 Z_\alpha^2 - Z_\alpha^1 Z_k^2) + \tau T^\alpha (Z_\beta^1 Z_k^2 - Z_k^1 Z_\beta^2) \\ &\quad + (\tau - \sigma) T^k (Z_\beta^1 Z_\alpha^2 - Z_\alpha^1 Z_\beta^2)) \omega^k.\end{aligned}$$

Moreover the matrix Z lies in $\text{SO}_3(\mathbb{R})$, so it is equal to its comatrix. Using this fact we compute that

$$L(Z)_2^1 = -(2\tau - \sigma) T^3 (T^1 \omega^1 + T^2 \omega^2),$$

$$L(Z)_3^1 = (2\tau - \sigma) T^1 T^2 \omega^1 + (2\tau - \sigma) (T^2)^2 \omega^2 - \tau \omega^2,$$

$$L(Z)_3^2 = -(2\tau - \sigma) (T^1)^2 \omega^1 - (2\tau - \sigma) T^1 T^2 \omega^2 + \tau \omega^1,$$

which proves the lemma. \square

Lemma 4.8. *We have*

$$\begin{aligned} L \wedge L &= \tau(2\tau - \sigma)T^3 \begin{pmatrix} 0 & -T^3 & T^2 \\ T^3 & 0 & -T^1 \\ -T^2 & T^1 & 0 \end{pmatrix} \omega^1 \wedge \omega^2 \\ &\quad + \tau(\tau - \sigma) \begin{pmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \omega^1 \wedge \omega^2. \end{aligned}$$

Proof. We compute that

$$\begin{aligned} L \wedge L &= \tau(2\tau - \sigma) \begin{pmatrix} 0 & T^1 & 0 \\ -T^1 & 0 & -T^3 \\ 0 & T^3 & 0 \end{pmatrix} \eta \wedge \omega^2 \\ &\quad + \tau(2\tau - \sigma) \begin{pmatrix} 0 & -T^2 & -T^3 \\ T^2 & 0 & 0 \\ T^3 & 0 & 0 \end{pmatrix} \eta \wedge \omega^1 \\ &\quad + \tau^2 \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \omega^1 \wedge \omega^2. \end{aligned}$$

We conclude using that $(T^1)^2 + (T^2)^2 + (T^3)^2 = 1$. □

Lemma 4.9. *We have*

$$\begin{aligned} L \wedge \Omega + \Omega \wedge L &= (2\tau - \sigma)\eta \wedge \begin{pmatrix} 0 & -dT^3 & dT^2 \\ dT^3 & 0 & -dT^1 \\ -dT^2 & dT^1 & 0 \end{pmatrix} \\ &\quad + \tau(2\tau - \sigma)T^3 \begin{pmatrix} 0 & T^3 & -T^2 \\ -T^3 & 0 & T^1 \\ T^2 & -T^1 & 0 \end{pmatrix} \omega^1 \wedge \omega^2 \\ &\quad + \tau(2\tau - \sigma) \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \omega^1 \wedge \omega^2 \\ &\quad + \tau \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix} d\omega^1 + \tau \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{pmatrix} d\omega^2. \end{aligned}$$

Proof. We compute that

$$\begin{aligned} L \wedge \Omega + \Omega \wedge L &= (2\tau - \sigma)\eta \wedge M \\ &\quad + \tau\omega^2 \wedge \begin{pmatrix} 0 & -\omega_2^3 & 0 \\ -\omega_3^2 & 0 & \omega_1^2 \\ 0 & \omega_2^1 & 0 \end{pmatrix} \\ &\quad + \tau\omega^1 \wedge \begin{pmatrix} 0 & \omega_3^1 & -\omega_2^1 \\ \omega_1^3 & 0 & 0 \\ -\omega_1^2 & 0 & 0 \end{pmatrix} \end{aligned}$$

with

$$M = \begin{pmatrix} 0 & T^2\omega_2^3 - T^1\omega_3^1 & -T^3\omega_3^2 + T^1\omega_2^1 \\ -T^1\omega_1^3 + T^2\omega_3^2 & 0 & T^3\omega_3^1 - T^2\omega_1^2 \\ T^1\omega_1^2 - T^3\omega_2^3 & -T^2\omega_2^1 + T^3\omega_1^3 & 0 \end{pmatrix}.$$

We conclude using lemma 4.5, formulas (4) and (5), and the fact that $(T^1)^2 + (T^2)^2 + (T^3)^2 = 1$. \square

For $y \in \mathcal{V}$, let $\mathcal{Z}(y)$ be the set of matrices $Z \in \text{SO}_3(\mathbb{R})$ such that the coefficients of the last line of Z are the $T^\beta(y)$. It is diffeomorphic to the circle \mathbb{S}^1 .

We now prove the following proposition.

Proposition 4.10. *Assume that the compatibility equations for \mathbb{E} are satisfied. Let $y_0 \in \mathcal{V}$ and $A_0 \in \mathcal{Z}(y_0)$. Then there exist a neighbourhood U_1 of y_0 in \mathcal{V} and a unique map $A : U_1 \rightarrow \text{SO}_3(\mathbb{R})$ such that*

$$\begin{aligned} A^{-1}dA &= \Omega, \\ \forall y \in U_1, \quad A(y) &\in \mathcal{Z}(y), \\ A(y_0) &= A_0. \end{aligned}$$

Proof. Let U be a coordinate neighbourhood in \mathcal{V} . The set

$$\mathcal{F} = \{(y, Z) \in U \times \text{SO}_3(\mathbb{R}); Z \in \mathcal{Z}(y)\}$$

is a manifold of dimension 3, and

$$\mathbb{T}_{(y,Z)}\mathcal{F} = \{(u, \zeta) \in \mathbb{T}_y U \oplus \mathbb{T}_Z \text{SO}_3(\mathbb{R}); \zeta_\beta^3 = (dT^\beta)_y(u)\}.$$

Let Z denote the projection $U \times \text{SO}_3(\mathbb{R}) \rightarrow \text{SO}_3(\mathbb{R}) \subset \mathcal{M}_3(\mathbb{R})$. We consider on \mathcal{F} the following matrix of 1-forms:

$$\Theta = Z^{-1}dZ - \Omega - L(Z)$$

where $L(Z)$ is defined in lemma 3.5, namely for $(y, Z) \in \mathcal{F}$ we have

$$\Theta_{(y,Z)} : \mathbb{T}_{(y,Z)}\mathcal{F} \rightarrow \mathcal{M}_3(\mathbb{R}),$$

$$\Theta_{(y,Z)}(u, \zeta) = Z^{-1}\zeta - \Omega_y(u) - L(Z)(u).$$

We claim that, for each $(y, Z) \in \mathcal{F}$, the space

$$\mathcal{D}(y, Z) = \ker \Theta_{(y,Z)}$$

has dimension 2. We first notice that the matrix Θ belongs to $\mathfrak{so}_3(\mathbb{R})$ since Ω , $L(Z)$ and $Z^{-1}dZ$ do. Moreover we have

$$(Z\Theta)_\beta^3 = dZ_\beta^3 - \sum_\gamma Z_\gamma^3 \omega_\beta^\gamma - \sum_\gamma Z_\gamma^3 L(Z)_\beta^\gamma = dT^\beta - \sum_\gamma T^\gamma \omega_\beta^\gamma - \sum_\gamma T^\gamma L(Z)_\beta^\gamma.$$

Using lemmas 4.5 and 4.7 we compute that

$$(Z\Theta)_\beta^3 = 0.$$

Thus the values of $\Theta_{(y,Z)}$ lie in the space

$$\mathcal{H} = \{H \in \mathfrak{so}_3(\mathbb{R}); (ZH)_\beta^3 = 0\},$$

which has dimension 1 (indeed, the map $F : \mathrm{SO}_3(\mathbb{R}) \rightarrow \mathbb{S}^2, Z \mapsto (Z^3)_\beta$ is a submersion, and we have $H \in \mathcal{H}$ if and only if $ZH \in \ker(dF)_Z$. Moreover, the space $\mathrm{T}_{(y,Z)}\mathcal{F}$ contains the subspace $\{(0, ZH); H \in \mathcal{H}\}$, and the restriction of $\Theta_{(y,Z)}$ on this subspace is the map $(0, ZH) \mapsto H$. Thus $\Theta_{(y,Z)}$ is onto \mathcal{H} , and consequently the linear map $\Theta_{(y,Z)}$ has rank 1. This finishes proving the claim.

We now prove that the distribution \mathcal{D} is involutive. We first compute that

$$\begin{aligned} d\Theta &= -Z^{-1}dZ \wedge Z^{-1}dZ - d\Omega - dL \\ &= -(\Theta + \Omega + L) \wedge (\Theta + \Omega + L) - d\Omega - dL \\ &= -\Theta \wedge \Theta - \Theta \wedge \Omega - \Omega \wedge \Theta - \Omega \wedge L - L \wedge \Omega \\ &\quad - \Omega \wedge \Omega - d\Omega - L \wedge L - dL. \end{aligned}$$

Using lemmas 4.4, 4.6, 4.8, 4.9 and the relation $\sigma = \frac{\kappa}{2\tau}$, we obtain

$$d\Theta = -\Theta \wedge \Theta - \Theta \wedge \Omega - \Omega \wedge \Theta.$$

From this formula we deduce that if $\xi_1, \xi_2 \in \mathcal{D}$, then $d\Theta(\xi_1, \xi_2) = 0$, and so $\Theta([\xi_1, \xi_2]) = \xi_1 \cdot \Theta(\xi_2) - \xi_2 \cdot \Theta(\xi_1) - d\Theta(\xi_1, \xi_2) = 0$, i.e., $[\xi_1, \xi_2] \in \mathcal{D}$. Thus the distribution \mathcal{D} is involutive, and so, by the theorem of Frobenius, it is integrable.

Let \mathcal{A} be the integral manifold through (y_0, A_0) . If $\zeta \in \mathrm{T}_{A_0}\mathrm{SO}_3(\mathbb{R})$ is such that $(0, \zeta) \in \mathrm{T}_{(y_0, A_0)}\mathcal{A} = \mathcal{D}(y_0, A_0)$, then we have $0 = \Theta_{(y_0, A_0)}(0, \zeta) = A_0^{-1}\zeta$. This proves that

$$\mathrm{T}_{(y_0, A_0)}\mathcal{A} \cap (\{0\} \times \mathrm{T}_{A_0}\mathrm{SO}_3(\mathbb{R})) = \{0\}.$$

Thus the manifold \mathcal{A} is locally the graph of a function $A : U_1 \rightarrow \mathrm{SO}_3(\mathbb{R})$ where U_1 is a neighbourhood of y_0 in U . By construction, this map satisfies the properties of proposition 4.10 and is unique. \square

Proposition 4.11. *Let $x_0 \in \mathbb{E}$ (without loss of generality we can assume that $x_0 \in \mathbb{E}'$). There exist a neighbourhood U_2 of y_0 contained in U_1 and a unique function $f : U_2 \rightarrow \mathbb{E}'$ such that*

$$df = (B \circ f)A\omega,$$

$$f(y_0) = x_0,$$

where ω is the column $(\omega^1, \omega^2, 0)$ and, for $x \in \mathbb{E}'$, $B(x) \in \mathcal{M}_3(\mathbb{R})$ is the matrix of the coordinates of the frame $(E_\alpha(x))$ in the frame (∂_{x^α}) .

Proof. We consider on $U_1 \times \mathbb{E}'$ the following matrix of 1-forms:

$$\Lambda = B^{-1}dx - A\omega,$$

namely, for $q \in U_1$ and $x \in \mathbb{E}'$ we have

$$\Lambda_{(q,x)} : \mathrm{T}_q U_1 \oplus \mathrm{T}_x \mathbb{E}' \rightarrow \mathcal{M}_{3,1}(\mathbb{R}),$$

$$\Lambda_{(q,x)}(u, v) = B(x)^{-1}v - A(q)\omega_q(u).$$

We first notice that for all $(q, x) \in U_1 \times \mathbb{E}'$ the linear map $\Lambda_{(q,x)}$ is onto $\mathcal{M}_{3,1}(\mathbb{R})$. Consequently the space

$$\mathcal{E}(q, x) = \ker \Lambda_{(q,x)}$$

has dimension 2. We will prove that this distribution \mathcal{E} is integrable.

We have

$$d\Lambda = -B^{-1}dB B^{-1} \wedge dx - dA \wedge \omega - A \wedge d\omega.$$

By equations (4) and (5) we have $d\omega = -\Omega \wedge \omega$; and by proposition 4.10 we have $dA = A\Omega + AL(A)$. Thus we get

$$d\Lambda = -B^{-1}dB \wedge \Lambda - B^{-1}dB \wedge A\omega - AL(A) \wedge \omega.$$

Using lemma 4.7 we compute that

$$L(A) \wedge \omega = -(2\tau - \sigma)T^3 \begin{pmatrix} T^1 \\ T^2 \\ T^3 \end{pmatrix} \omega^1 \wedge \omega^2 - \begin{pmatrix} 0 \\ 0 \\ \sigma \end{pmatrix} \omega^1 \wedge \omega^2,$$

and thus, using the fact that $A_\beta^3 = T^\beta$ and $A = \text{com}A$, we get

$$AL(A) \wedge \omega = \begin{pmatrix} -\sigma A_3^1 \\ -\sigma A_3^2 \\ -2\tau T^3 \end{pmatrix} \omega^1 \wedge \omega^2.$$

We will use the notation (x, y, z) instead of (x^1, x^2, x^3) for the coordinates in \mathbb{E} and we will use the local models described in sections 2.2, 2.3 and 2.4. Using formulas (1), (2) and (3), we get that the matrix B is

$$B = \begin{pmatrix} \lambda^{-1} \cos(\sigma z) & -\lambda^{-1} \sin(\sigma z) & 0 \\ \lambda^{-1} \sin(\sigma z) & \lambda^{-1} \cos(\sigma z) & 0 \\ \tau(x \sin \sigma z - y \cos \sigma z) & \tau(x \cos \sigma z + y \sin \sigma z) & 1 \end{pmatrix},$$

with

$$\lambda = \frac{1}{1 + \frac{\kappa}{4}(x^2 + y^2)}.$$

We will write

$$A\omega = \begin{pmatrix} \alpha^1 \\ \alpha^2 \\ \eta \end{pmatrix}$$

with

$$\alpha^j = A_1^j \omega^1 + A_2^j \omega^2.$$

Then we have

$$\Lambda = B^{-1}dX - A\omega = \begin{pmatrix} \lambda(\cos(\sigma z)dx + \sin(\sigma z)dy) - \alpha^1 \\ \lambda(-\sin(\sigma z)dx + \cos(\sigma z)dy) - \alpha^2 \\ \tau\lambda(ydx - xdy) + dz - \eta \end{pmatrix}.$$

We also compute that

$$B^{-1}dB = \begin{pmatrix} \frac{\kappa}{2}\lambda(xdx + ydy) & -\sigma dz & 0 \\ \sigma dz & \frac{\kappa}{2}\lambda(xdx + ydy) & 0 \\ a & b & 0 \end{pmatrix}$$

with

$$a = \frac{\tau\kappa}{2}\lambda(y \cos(\sigma z) - x \sin(\sigma z))(xdx + ydy) + \tau(\sin(\sigma z)dx - \cos(\sigma z)dy),$$

$$b = -\frac{\tau\kappa}{2}\lambda(x \cos(\sigma z) + y \sin(\sigma z))(xdx + ydy) + \tau(\cos(\sigma z)dx + \sin(\sigma z)dy).$$

Thus we have

$$\begin{aligned} B^{-1}dB \wedge A\omega + AL(A) \wedge \omega &= \begin{pmatrix} \frac{\kappa}{2}\lambda(xdx + ydy) \wedge \alpha^1 - \sigma dz \wedge \alpha^2 \\ \sigma dz \wedge \alpha^1 + \frac{\kappa}{2}\lambda(xdx + ydy) \wedge \alpha^2 \\ a \wedge \alpha^1 + b \wedge \alpha^2 \end{pmatrix} \\ &+ \begin{pmatrix} -\sigma A_3^1 \\ -\sigma A_3^2 \\ -2\tau T^3 \end{pmatrix} \omega^1 \wedge \omega^2. \end{aligned}$$

Using the above expression for Λ we get

$$\lambda dx = \cos(\sigma z)\Lambda^1 - \sin(\sigma z)\Lambda^2 + \cos(\sigma z)\alpha^1 - \sin(\sigma z)\alpha^2,$$

$$\lambda dy = \sin(\sigma z)\Lambda^1 + \cos(\sigma z)\Lambda^2 + \sin(\sigma z)\alpha^1 + \cos(\sigma z)\alpha^2,$$

$$dz = \Lambda^3 + \eta - \tau\lambda(ydx - xdy).$$

The term in the first line of the matrix $B^{-1}dB \wedge A\omega + AL(A)$ is

$$\begin{aligned} \frac{\kappa}{2}(y \cos(\sigma z) - x \sin(\sigma z))\alpha^2 \wedge \alpha^1 + \sigma\tau(y \cos(\sigma z) - x \sin(\sigma z))\alpha^1 \wedge \alpha^2 \\ -\sigma\eta \wedge \alpha^2 - \sigma A_3^1\omega^1 \wedge \omega^2 + \chi^1 \end{aligned}$$

where χ^1 is a linear combination of the Λ^α (the coefficients being 1-forms). Since $\sigma = \frac{\kappa}{2\tau}$, the first two terms in this expression cancel. Moreover we have $\eta \wedge \alpha^2 = (A_1^3 A_2^2 - A_2^3 A_1^2)\omega^1 \wedge \omega^2 = -A_3^1\omega^1 \wedge \omega^2$, hence the term in the first line of the matrix $B^{-1}dB \wedge A\omega + AL(A)$ is χ^1 . In the same way, the term in the second line of the matrix $B^{-1}dB \wedge A\omega + AL(A)$ is a linear combination of the Λ^α which will be denoted by χ^2 . Finally we compute that the term in the third line of the matrix $B^{-1}dB \wedge A\omega + AL(A)$ is

$$\left(\frac{2\tau}{\lambda} - \frac{\tau\kappa}{2}(x^2 + y^2) \right) \alpha^1 \wedge \alpha^2 - 2\tau T^3 \omega^1 \wedge \omega^2 + \chi^3$$

where χ^1 is a linear combination of the Λ^α . Since $\lambda^{-1} = 1 + \frac{\kappa}{4}(x^2 + y^2)$ and $\alpha^1 \wedge \alpha^2 = (A_1^1 A_2^2 - A_2^1 A_1^2)\omega^1 \wedge \omega^2 = T^3 \omega^1 \wedge \omega^2$, this term is simply χ^3 . We conclude that

$$B^{-1}dB \wedge A\omega + AL(A) = \chi$$

where χ is a matrix of 2-forms which are linear combinations of the coefficients of Λ . Finally we have

$$d\Lambda = -B^{-1}dB \wedge \Lambda - \chi.$$

From this formula we deduce that if $\xi_1, \xi_2 \in \mathcal{E}$, then $d\Lambda(\xi_1, \xi_2) = 0$, and so $[\xi_1, \xi_2] \in \mathcal{E}$. Thus the distribution \mathcal{E} is involutive, and so, by the theorem of Frobenius, it is integrable.

Let \mathcal{A} be the integral manifold through (y_0, x_0) . If $v \in T_{x_0}\mathbb{E}$ is such that $(0, v) \in T_{(y_0, x_0)}\mathcal{A} = \mathcal{D}(y_0, x_0)$, then we have $0 = \Lambda_{(y_0, x_0)}(0, v) = B(x_0)^{-1}v$. This proves that

$$T_{(y_0, x_0)}\mathcal{A} \cap (\{0\} \times T_{x_0}\mathbb{E}) = \{0\}.$$

Thus the manifold \mathcal{A} is locally the graph of a function $A : U_2 \rightarrow \mathbb{E}'$ where U_2 is a neighbourhood of y_0 in U_1 . By construction, this map satisfies the properties of proposition 4.10 and is unique. \square

We now prove the theorem.

Proof of theorem 4.3. Let $y_0 \in \mathcal{V}$, $A_0 \in \mathcal{Z}(y_0)$ and $x_0 \in \mathbb{E}'$. We consider on \mathcal{V} a local orthonormal frame (e_1, e_2) in the neighbourhood of y_0 and we keep the same notations. Then by propositions 4.10 and 4.11 there exists a unique map $A : U_2 \rightarrow \text{SO}^3(\mathbb{R})$ such that

$$A^{-1}dA = \Omega + L(A),$$

$$\forall y \in U_1, \quad A(y) \in \mathcal{Z}(y),$$

$$A(y_0) = A_0,$$

and a unique map $f : U_2 \rightarrow \mathbb{E}'$ such that

$$df = (B \circ f)A\omega,$$

$$f(y_0) = x_0,$$

where U_2 is a neighbourhood of y_0 , which we can assume simply connected. We will check that f has the properties required in the theorem on U_2 .

We have $df^\alpha(e_k) = (B(f)A)_k^\alpha$, so in the frame (∂_{x^α}) the vector $df(e_k)$ is given by the column k of the matrix BA , which is invertible. Hence df has rank 2, and thus f is an immersion. Moreover, in the frame (E_α) the vector $df(e_k)$ is given by the column k of the matrix A , which is orthogonal, and thus we have $\langle df(e_p), df(e_q) \rangle = \delta_q^p$, which means that f is an isometry.

The columns of $A(y)$ form a direct orthonormal frame of \mathbb{E} . The first and second columns form a direct orthonormal frame of $T_{f(y)}f(\mathcal{V})$. Thus the third column gives, in the frame (E_α) , the unit normal $N(f(y))$ to $f(\mathcal{V})$ in \mathbb{E} at the point $f(y)$.

We set $X_j = df(e_j)$. Then we have

$$\begin{aligned} dA_j^\alpha(e_k) &= \langle \bar{\nabla}_{X_k} X_j, E_\alpha \rangle + \langle X_j, \bar{\nabla}_{X_k} E_\alpha \rangle \\ &= \langle \bar{\nabla}_{X_k} X_j, E_\alpha \rangle + \sum_\gamma \sum_\delta A_k^\gamma A_j^\delta \bar{\Gamma}_{\gamma\alpha}^\delta \\ &= \langle \bar{\nabla}_{X_k} X_j, E_\alpha \rangle + (AL(A))_j^\alpha(e_k), \end{aligned}$$

so

$$\begin{aligned}
\langle \bar{\nabla}_{X_k} X_j, N \rangle &= \sum_{\alpha} \langle \bar{\nabla}_{X_k} X_j, E_{\alpha} \rangle A_3^{\alpha} = \sum_{\alpha} A_3^{\alpha} (dA - AL(A))_j^{\alpha}(e_k) \\
&= \sum_{\alpha} A_3^{\alpha} (A\Omega)_j^{\alpha}(e_k) = \sum_{\alpha} \sum_{\gamma} A_{\gamma}^{\alpha} A_3^{\alpha} \omega_j^{\gamma}(e_k) \\
&= \omega_j^3(e_k) = \langle Se_k, e_j \rangle.
\end{aligned}$$

This means that the shape operator of $f(\mathcal{V})$ in \mathbb{E} is $df \circ S \circ df^{-1}$.

Finally, the coefficients of the vertical vector $\xi = E_3$ in the orthonormal frame (X_1, X_2, N) are given by the last line of A . Since $A(y) \in \mathcal{Z}(y)$ for all $y \in U_2$ we get

$$\xi = \sum_j T^j X_j + T^3 N = df(T) + \nu N.$$

We now prove that the local immersion is unique up to a global isometry of \mathbb{E} preserving ξ (and also, consequently, the orientation of the base of the fibration). Let $\tilde{f} : U_3 \rightarrow \mathbb{E}$ be another immersion satisfying the conclusion of the theorem, where U_3 is a simply connected neighbourhood of y_0 included in U_2 , let (\tilde{X}_{β}) be the associated frame (i.e., $\tilde{X}_j = d\tilde{f}(e_j)$ and \tilde{X}_3 is the normal of $\tilde{f}(\mathcal{V})$) and let \tilde{A} the matrix of the coordinates of the frame (\tilde{X}_{β}) in the frame (E_{α}) . Up to an isometry of \mathbb{E} (which is necessarily direct), we can assume that $f(y_0) = \tilde{f}(y_0)$ and that the frames $(X_{\beta}(y_0))$ and $(\tilde{X}_{\beta}(y_0))$ coincide, i.e., $A(y_0) = \tilde{A}(y_0)$. We notice that this isometry necessarily fixes ξ since the T^{α} are the same for x and \tilde{x} . The matrices A and \tilde{A} satisfy $A^{-1}dA = \Omega + L(A)$ and $\tilde{A}^{-1}d\tilde{A} = \Omega + L(\tilde{A})$ (see section 3.3), $A(y), \tilde{A}(y) \in \mathcal{Z}(y)$ and $A(y_0) = \tilde{A}(y_0)$, thus by the uniqueness of the solution of the equation in proposition 4.10 we get $A(y) = \tilde{A}(y)$. We conclude similarly that $f = \tilde{f}$ on U_3 .

The proof that this local immersion f can be extended to the whole \mathcal{V} (since \mathcal{V} is simply connected) is exactly the same as the proof of the corresponding statement in theorem 3.3 in [Dan04] (it is a standard argument). \square

Remark 4.12. If (ds^2, S, T, ν) satisfies the compatibility equations and correspond to an immersion $f : \Sigma \rightarrow \mathbb{E}$, then $(ds^2, S, -T, -\nu)$ also satisfies the compatibility equations and corresponds to the immersion $\sigma \circ f$ where σ is an isometry of \mathbb{E} reversing the orientations of both the fibers and the base of the fibration.

5. CONSTANT MEAN CURVATURE SURFACES IN 3-DIMENSIONAL HOMOGENEOUS MANIFOLDS

In this section we will give an application of theorem 4.3 to constant mean curvature surfaces (CMC) in 3-dimensional homogeneous manifolds with 4-dimensional isometry group. Abresch and Rosenberg proved that there exists a holomorphic quadratic differential for CMC surfaces in $S^2 \times \mathbb{R}$ and

$\mathbb{H}^2 \times \mathbb{R}$, generalizing the Hopf differential for CMC surfaces in 3-dimensional space forms ([AR03]). Since the Hopf differential is a very useful tool for CMC surfaces, this motivated many works on CMC surfaces in $\mathbb{S}^2 \times \mathbb{R}$ and $\mathbb{H}^2 \times \mathbb{R}$. Recently, Abresch announced the existence of a holomorphic quadratic differential for CMC surfaces in all 3-dimensional homogeneous manifolds with 4-dimensional isometry group ([Abr04]). This indicates that the theory of CMC surfaces in these manifolds may be particularly interesting.

We will consider constant mean curvature immersions of oriented surfaces. Consequently the mean curvature will be defined with a sign: it will be positive if the mean curvature vector induces the same orientation as the initial orientation, and it will be negative if the mean curvature vector induces the opposite orientation.

We will denote by I and J the identity and the rotation of angle $\frac{\pi}{2}$ on the tangent bundle of a surface.

5.1. A generalized Lawson correspondence. It is well known that there exists an isometric correspondence between certain simply connected CMC surfaces in space-forms: more precisely, every simply connected CMC H_1 surface in $\mathbb{M}^3(K_1)$ is isometric to a simply connected CMC H_2 surface in $\mathbb{M}^3(K_2)$ with $K_1 - K_2 = H_2^2 - H_1^2$, and the shape operators of these two surfaces differ by $(H_2 - H_1)I$. Two such surfaces are called cousin surfaces. This correspondence is often called the Lawson correspondence. In particular, any simply connected minimal surface in \mathbb{S}^3 is isometric to a CMC 1 surface in \mathbb{R}^3 , and any minimal surface in \mathbb{R}^3 is isometric to a CMC 1 surface in \mathbb{H}^3 .

The Lawson correspondence is a consequence of the Gauss and Codazzi equations in the space-forms.

In this section we will use the compatibility equations for homogeneous 3-manifolds with 4-dimensional isometry group and theorem 4.3 to prove the existence of an isometric correspondence between certain simply connected CMC surfaces in these 3-manifolds. Hence this will be a generalisation of the Lawson correspondence.

The technique will be to start with some data (ds^2, S, T, ν) on a surface satisfying the compatibility equations for some homogeneous 3-manifold and to modify them in order to get data satisfying the compatibility equations for another homogeneous 3-manifold. An important fact is that the space of symmetric traceless operators is globally invariant by rotation. The easiest change is to keep ds^2 and ν , and to rotate T and the traceless part of S by some fixed angles; the Codazzi equation then implies that we need to take the same angle for T and the traceless part of S .

Proposition 5.1. *Let \mathbb{E}_1 and \mathbb{E}_2 be two 3-dimensional homogeneous manifolds with 4-dimensional isometry groups, of base curvatures κ_1 and κ_2 and bundle curvatures τ_1 and τ_2 respectively. Assume that*

$$\kappa_1 - 4\tau_1^2 = \kappa_2 - 4\tau_2^2.$$

Let H_1 and H_2 be two real numbers such that

$$\tau_1^2 + H_1^2 = \tau_2^2 + H_2^2.$$

Let \mathcal{V} be a surface with a quadruple (ds^2, S_1, T_1, ν) satisfying the compatibility equations for \mathbb{E}_1 and such that

$$\operatorname{tr} S_1 = 2H_1.$$

Let

$$\theta \in \mathbb{R},$$

$$T_2 = e^{\theta J} T_1,$$

$$S_2 = e^{\theta J} (S_1 - H_1 I) + H_2 I.$$

In particular S_2 is symmetric and satisfies

$$\operatorname{tr} S_2 = 2H_2.$$

If the real number θ satisfies

$$(12) \quad \tau_2 + iH_2 = e^{i\theta} (\tau_1 + iH_1),$$

then the quadruple (ds^2, S_2, T_2, ν) satisfies the compatibility equations for \mathbb{E}_2 .

Conversely, if the function ν is not identically zero and if the quadruple (ds^2, S_2, T_2, ν) satisfies the compatibility equations for \mathbb{E}_2 , then (12) holds.

Proof. The fact that S_2 is symmetric comes from the fact that the space of symmetric traceless operators is invariant by a rotation.

We have

$$\det(S_k - H_k I) = \det S_k - H_k^2$$

for $k = 1, 2$, and so

$$\det S_1 = \det S_2 + H_1^2 - H_2^2.$$

Let K be the Gauss curvature of the metric ds^2 . By the Gauss equation (8) we have

$$\begin{aligned} K &= \det S_1 + \tau_1^2 + (\kappa_1 - 4\tau_1^2)\nu^2 \\ &= \det S_2 + H_1^2 - H_2^2 + \tau_1^2 + (\kappa_1 - 4\tau_1^2)\nu^2 \\ &= \det S_2 + \tau_2^2 + (\kappa_2 - 4\tau_2^2)\nu^2 \end{aligned}$$

since $\kappa_1 - 4\tau_1^2 = \kappa_2 - 4\tau_2^2$ and $\tau_1^2 + H_1^2 = \tau_2^2 + H_2^2$. Thus the quadruple (ds^2, S_2, T_2, ν) satisfies the Gauss equation for \mathbb{E}_2 .

Since J commutes with ∇_X for all vector fields X , we have

$$\nabla_X S_2 Y - \nabla_Y S_2 X - S_2[X, Y] = e^{\theta J} (\nabla_X S_1 Y - \nabla_Y S_1 X - S_1[X, Y]).$$

On the other hand, a computation done in the proof of proposition 4.1 in [Dan04] shows that

$$\langle Y, T_2 \rangle X - \langle X, T_2 \rangle Y = e^{\theta J} (\langle Y, T_1 \rangle X - \langle X, T_1 \rangle Y).$$

Hence the Codazzi equation for \mathbb{E}_2 is satisfied by (ds^2, S_2, T_2, ν) .

To prove that the quadruple (ds^2, S_2, T_2, ν) satisfies the compatibility equations (10) and (11) for \mathbb{E}_2 , it suffices to prove that

$$(13) \quad S_2 - \tau_2 J = e^{\theta J} (S_1 - \tau_1 J).$$

Using the expression of S_2 , equation (13) is equivalent to

$$(14) \quad H_2 I - \tau_2 J = e^{\theta J} (H_1 I - \tau_1 J).$$

We notice that this is a purely algebraic condition: the shape operators are not involved anymore. We consider a local direct orthonormal frame and we will identify the operators with their matrix in this frame. Then we have

$$J = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}.$$

Then equation (14) is equivalent to

$$\begin{cases} H_2 &= H_1 \cos \theta + \tau_1 \sin \theta, \\ \tau_2 &= \tau_1 \cos \theta - H_1 \sin \theta. \end{cases} \quad ,$$

i.e., it is equivalent to equation (12). This proves the first assertion of the theorem.

Conversely, if (ds^2, S_2, T_2, ν) satisfies the compatibility equations for \mathbb{E}_2 , then the compatibility equations (10) for (ds^2, S_1, T_1, ν) and (ds^2, S_2, T_2, ν) imply that (13) holds at every point where $\nu \neq 0$. If there exists a point where $\nu \neq 0$, this implies that (12) holds. \square

Theorem 5.2. *Let \mathbb{E}_1 and \mathbb{E}_2 be two 3-dimensional homogeneous manifolds with 4-dimensional isometry groups, of base curvatures κ_1 and κ_2 and bundle curvatures τ_1 and τ_2 respectively, and such that*

$$\kappa_1 - 4\tau_1^2 = \kappa_2 - 4\tau_2^2.$$

Let ξ_1 and ξ_2 be the vertical vector fields of \mathbb{E}_1 and \mathbb{E}_2 respectively.

Let Σ be a simply connected Riemann surface and let $x_1 : \Sigma \rightarrow \mathbb{E}_1$ be a conformal constant mean curvature H_1 immersion with $H_1^2 \geq \tau_2^2 - \tau_1^2$. Let N_1 be the induced normal (compatible with the orientation of Σ). Let S_1 be the symmetric operator on Σ induced by the shape operator of $x_1(\Sigma)$ associated to the normal N_1 . Let T_1 be the vector field on Σ such that $dx_1(T_1)$ is the projection of ξ_1 onto $T(x_1(\Sigma))$. Let $\nu = \langle N_1, \xi_1 \rangle$.

Let $H_2 \in \mathbb{R}$ such that

$$\tau_1^2 + H_1^2 = \tau_2^2 + H_2^2.$$

Let $\theta \in \mathbb{R}$ such that

$$\tau_2 + iH_2 = e^{i\theta} (\tau_1 + iH_1).$$

Then there exists a conformal immersion $x_2 : \Sigma \rightarrow \mathbb{E}_2$ such that:

- (1) *the metrics induced on Σ by x_1 and x_2 are the same,*
- (2) *the symmetric operator on Σ induced by the shape operator of $x_2(\Sigma)$ is $e^{\theta J} (S_1 - H_1 I) + H_2 I$,*
- (3) *$\xi_2 = dx_2(e^{\theta J} T_1) + \nu N_2$ where N_2 is the unit normal to x_2 .*

Moreover, this immersion x_2 is unique up to isometries of \mathbb{E}_2 preserving the orientations of both the fibers and the base of the fibration, and it has constant mean curvature H_2 .

The immersions x_1 and x_2 are called sister immersions. The number θ is called the phase of (x_1, x_2) .

This means that there exists an isometric correspondence between CMC H_1 simply connected surfaces in \mathbb{E}_1 and CMC H_2 simply connected surfaces in \mathbb{E}_2 .

Proof. Let ds^2 be the metric on Σ induced by x_1 . Then (ds^2, S_1, T_1, ν) satisfies the compatibility equations for \mathbb{E}_1 . Thus, by proposition 5.1, the quadruple $(ds^2, S_2, e^{\theta J}T_1, \nu)$ with $S_2 = e^{\theta J}(S_1 - H_1I) + H_2I$ also does. Thus by theorem 4.3 there exists an immersion x_2 satisfying properties 1, 2, and 3, and this immersion is unique up to isometries of \mathbb{E}_2 preserving the orientations of both the fibers and the base of the fibration. Moreover, we have $\text{tr} S_2 = 2H_2$, i.e., the immersion x_2 has mean curvature H_2 . \square




FIGURE 1. The correspondence of the sister surfaces

Figure 1 helps visualizing which classes of CMC surfaces are related by the sister surface correspondence. We start from a CMC surface in some homogeneous 3-manifold. Then we can go horizontally on the graph. We can go to the left until reaching a manifold with $\tau = 0$; in this case the absolute mean curvature $|H|$ increases. We can go to the right until reaching $H = 0$; in this case $|H|$ decreases.

A particularly interesting case is when \mathbb{E}_1 is the Heisenberg space Nil_3 with its standard metric ($\kappa_1 = 0$, $\tau_1 = \frac{1}{2}$) and $\mathbb{E}_2 = \mathbb{H}^2 \times \mathbb{R}$ ($\kappa_2 = -1$, $\tau_2 = 0$). Then CMC H_1 surfaces in Nil_3 correspond isometrically to CMC H_2 surfaces in $\mathbb{H}^2 \times \mathbb{R}$ with $H_2^2 = H_1^2 + \frac{1}{4}$. In particular we have the following corollary.

Corollary 5.3. *There exists an isometric correspondence with phase $\theta = \frac{\pi}{2}$ between simply connected minimal surfaces in the Heisenberg space Nil_3 and simply connected CMC $\frac{1}{2}$ surfaces in $\mathbb{H}^2 \times \mathbb{R}$.*

The fact that $\theta = \frac{\pi}{2}$ suggests that this correspondence looks like the conjugate cousin correspondence between minimal surfaces in \mathbb{R}^3 and CMC 1 surfaces in \mathbb{H}^3 ([Bry87], [UY93]). This correspondence has nice geometric properties, and is useful to construct CMC 1 surfaces in \mathbb{H}^3 with some prescribed geometric properties starting from a solution of a Plateau problem in \mathbb{R}^3 (see for example [Kar01], [Dan03]).

In particular, if a minimal surface Σ_1 in Nil_3 contains an ambient geodesic γ , then the normal curvature of γ vanishes, and so

$$0 = \langle \gamma', S_1 \gamma' \rangle = \langle \gamma', -JS_2 \gamma' + \frac{1}{2} J \gamma' \rangle = -\langle \gamma', JS_2 \gamma' \rangle.$$

This means that $S\gamma'$ is colinear to γ' , i.e., γ is a geodesic line of curvature in the sister CMC $\frac{1}{2}$ surface in $\mathbb{H}^2 \times \mathbb{R}$.

We describe two examples of sister CMC $\frac{1}{2}$ surfaces in $\mathbb{H}^2 \times \mathbb{R}$ of minimal surfaces in Nil_3 . We will use the exponential coordinates given in section 2.3 (with $\tau = \frac{1}{2}$). We will denote between parentheses $()$ the coordinates of a vector in the coordinate frame $(\partial_x, \partial_y, \partial_z)$, and between brackets $[]$ the coordinates of a vector in the canonical frame (E_1, E_2, E_3) ; with these notations one has

$$\begin{pmatrix} a \\ b \\ c \end{pmatrix} = \begin{bmatrix} a \\ b \\ \frac{1}{2}(ya - xb) + c \end{bmatrix}.$$

Example 5.4 (vertical plane). A vertical plane \mathcal{P} in Nil_3 is a flat minimal surface (but not totally geodesic). A conformal parametrisation is

$$\varphi : (u, v) \mapsto \begin{pmatrix} v \\ 0 \\ u \end{pmatrix}.$$

We have

$$\varphi_u = E_3, \quad \varphi_v = E_1, \quad N = E_2,$$

and so

$$\begin{aligned}\nu &= 0, \\ \langle T, \partial_u \rangle &= \langle \xi, \varphi_u \rangle = 1, \\ \langle T, \partial_v \rangle &= \langle \xi, \varphi_v \rangle = 0,\end{aligned}$$

i.e.,

$$T = \partial_u.$$

We also have

$$\bar{\nabla}_{\varphi_u} N = \frac{1}{2}E_1 = \frac{1}{2}\varphi_u, \quad \bar{\nabla}_{\varphi_v} N = \frac{1}{2}E_3 = \frac{1}{2}\varphi_v,$$

so in the direct orthonormal frame (∂_u, ∂_v) we have

$$S = -\frac{1}{2} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

We now show that the CMC $\frac{1}{2}$ sister in $\mathbb{H}^2 \times \mathbb{R}$ of \mathcal{P} is the product $\mathcal{H} \times \mathbb{R}$ where \mathcal{H} is a horocycle in \mathbb{H}^2 . We will use the upper half-plane model for \mathbb{H}^2 . Then $\mathbb{H}^2 \times \mathbb{R} = \{(x, y, z) \in \mathbb{R}^3; y > 0\}$ and the metric is $ds^2 = \frac{1}{y^2}(dx^2 + dy^2) + dz^2$. We consider the direct orthonormal frame (E_1, E_2, E_3) defined by $E_1 = y\partial_x$, $E_2 = y\partial_y$, $E_3 = \partial_z$; it satisfies $\bar{\nabla}_{E_1} E_1 = E_2$, $\bar{\nabla}_{E_1} E_2 = -E_1$, and the other derivatives vanish. For \mathcal{H} , we can choose the curve of equation $y = 1$ in \mathbb{H}^2 . A conformal parametrization of $\mathcal{H} \times \mathbb{R}$ is

$$\tilde{\varphi} : (u, v) \mapsto \begin{pmatrix} -u \\ 1 \\ v \end{pmatrix}.$$

We have

$$\tilde{\varphi}_u = -E_1, \quad \tilde{\varphi}_v = E_3, \quad N = E_2,$$

and so

$$\tilde{\nu} = 0, \quad \tilde{T} = \partial_v.$$

We also have

$$\bar{\nabla}_{\tilde{\varphi}_u} N = E_1 = -\tilde{\varphi}_u, \quad \bar{\nabla}_{\tilde{\varphi}_v} N = 0,$$

so in the direct orthonormal frame (∂_u, ∂_v) we have

$$\tilde{S} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}.$$

Hence, $\tilde{\varphi}$ induces on \mathbb{R}^2 the same metric as φ , and we have $\tilde{\nu} = \nu$, $\tilde{T} = JT$ and $\tilde{S} = JS + \frac{1}{2}I$, so $\tilde{\varphi}$ is the sister immersion of φ . The vertical lines in \mathcal{P} are mapped to horizontal horocycles in $\mathcal{H} \times \mathbb{R}$, and horizontal lines in \mathcal{P} are mapped to vertical lines in $\mathcal{H} \times \mathbb{R}$.

Example 5.5 (surface of equation $z = 0$). The surface \mathcal{A} of equation $z = 0$ in the exponential coordinates is a minimal surface in Nil_3 which is invariant by rotation about the z -axis (but it is not invariant by any translation; see [FMP99]). We consider the following parametrisation:

$$\varphi : (u, v) \mapsto \begin{pmatrix} u \cos v \\ u \sin v \\ 0 \end{pmatrix},$$

for $u > 0$ (the origin in \mathcal{A} is excluded). We have

$$\begin{aligned} \varphi_u &= \begin{pmatrix} \cos v \\ \sin v \\ 0 \end{pmatrix} = \begin{bmatrix} \cos v \\ \sin v \\ 0 \end{bmatrix}, \\ \varphi_v &= \begin{pmatrix} -u \sin v \\ u \cos v \\ 0 \end{pmatrix} = \begin{bmatrix} -u \sin v \\ u \cos v \\ -\frac{1}{2}u^2 \end{bmatrix}, \end{aligned}$$

so

$$\begin{aligned} \langle \varphi_u, \varphi_u \rangle &= 1, \\ \langle \varphi_v, \varphi_v \rangle &= u^2 \left(1 + \frac{u^2}{4} \right), \\ \langle \varphi_u, \varphi_v \rangle &= 0. \end{aligned}$$

The unit normal vector is $N = \frac{\varphi_u \times \varphi_v}{\|\varphi_u \times \varphi_v\|}$; we compute that

$$\nu = \frac{1}{\sqrt{1 + \frac{u^2}{4}}}.$$

A direct orthonormal frame (e_1, e_2) is given by

$$e_1 = \partial_u, \quad e_2 = \frac{1}{u\sqrt{1 + \frac{u^2}{4}}} \partial_v.$$

We compute that

$$T = -\frac{u}{2\sqrt{1 + \frac{u^2}{4}}} \partial_v.$$

We now show that the CMC $\frac{1}{2}$ sister in $\mathbb{H}^2 \times \mathbb{R}$ of \mathcal{A} is the CMC $\frac{1}{2}$ graph \mathcal{B} of theorem D in [NR04]. This surface \mathcal{B} is also invariant by rotation about a vertical axis. If we take for \mathbb{H}^2 the Poincaré unit disk model, then \mathcal{B} is the graph of the function $(x, y) \mapsto \frac{2}{\sqrt{1-x^2-y^2}}$. We will use the Lorentzian for $\mathbb{H}^2 \times \mathbb{R}$, i.e.,

$$\mathbb{H}^2 \times \mathbb{R} = \{(x^0, x^1, x^2, x^3) \in \mathbb{L}^3 \times \mathbb{R}; -(x^0)^2 + (x^1)^2 + (x^2)^2 = -1, x^0 > 0\}$$

with the restriction of the quadratic form $-(dx^0)^2 + (dx^1)^2 + (dx^2)^2 + (dx^3)^2$. In this model, we consider the map

$$\tilde{\varphi} : (u, v) \mapsto \begin{pmatrix} 1 + \frac{u^2}{2} \\ u\sqrt{1 + \frac{u^2}{4}} \cos v \\ u\sqrt{1 + \frac{u^2}{4}} \sin v \\ 2\sqrt{1 + \frac{u^2}{4}} \end{pmatrix},$$

for $u > 0$. We can check that it is a parametrization of \mathcal{B} minus the origin (using that the correspondence between the Poincaré model and the Lorentzian model is given by $x + iy = \frac{x^1 + ix^2}{1 + x^0}$, $z = x^3$). We have

$$\tilde{\varphi}_u = \frac{1}{\sqrt{1 + \frac{u^2}{4}}} \begin{pmatrix} u\sqrt{1 + \frac{u^2}{4}} \\ 1 + \frac{u^2}{2} \cos v \\ 1 + \frac{u^2}{2} \sin v \\ \frac{u}{2} \end{pmatrix}, \quad \tilde{\varphi}_v = \begin{pmatrix} 0 \\ -u\sqrt{1 + \frac{u^2}{4}} \sin v \\ u\sqrt{1 + \frac{u^2}{4}} \cos v \\ 0 \end{pmatrix},$$

so

$$\begin{aligned} \langle \tilde{\varphi}_u, \tilde{\varphi}_u \rangle &= 1, \\ \langle \tilde{\varphi}_v, \tilde{\varphi}_v \rangle &= u^2 \left(1 + \frac{u^2}{4} \right), \\ \langle \tilde{\varphi}_u, \tilde{\varphi}_v \rangle &= 0, \end{aligned}$$

so $\tilde{\varphi}$ induces the same metric as φ . We compute that

$$\tilde{T} = \frac{u}{2\sqrt{1 + \frac{u^2}{4}}} e_1 = JT.$$

Thus we also have $\tilde{\nu}^2 = \nu^2$. Moreover, $\tilde{\varphi}_u$ points outwards and $\tilde{\varphi}_v$ points in the counter-clockwise direction, so the normal \tilde{N} points up, i.e., $\tilde{\nu} > 0$. So we get

$$\tilde{\nu} = \nu.$$

It remains to check that $\tilde{S} = JS + \frac{1}{2}I$. Since $\nu \neq 0$, the compatibility equations (10) for φ and $\tilde{\varphi}$ imply that $\tilde{S} = J(S - \frac{1}{2}J) = JS + \frac{1}{2}I$. Hence $\tilde{\varphi}$ is the sister immersion of φ .

The straight lines in \mathcal{A} passing through the origin are mapped to the generatrices of \mathcal{B} , which are lines of curvatures lying in vertical planes. Thus the symmetries of \mathcal{B} with respect to these vertical planes correspond to the symmetries of \mathcal{A} with respect to the straight lines passing through the origin.

Example 5.6 (CMC rotational spheres). The sister of the CMC H_1 rotational sphere in Nil_3 is the CMC $\sqrt{H_1^2 + \frac{1}{4}}$ rotational sphere in $\mathbb{H}^2 \times \mathbb{R}$. Indeed, the sister of this sphere is a possibly immersed CMC sphere in $\mathbb{H}^2 \times \mathbb{R}$, which is necessarily rotational by a theorem of Abresch and Rosenberg ([AR03]).

Remark 5.7. CMC H surfaces in $\mathbb{H}^2 \times \mathbb{R}$ have very different properties when $H \leq \frac{1}{2}$ and when $H > \frac{1}{2}$; for example compact embedded CMC H surfaces exist only for $H > \frac{1}{2}$. The reader can refer for example to [NR04]. An explanation is that CMC H surfaces in $\mathbb{H}^2 \times \mathbb{R}$ arise from minimal surfaces in a Berger sphere when $H > \frac{1}{2}$, in Nil_3 when $H = \frac{1}{2}$, and in a space $\widetilde{\text{PSL}}_2(\mathbb{R})$ when $H < \frac{1}{2}$.

Remark 5.8. When $\kappa - 4\tau^2 = 0$, the sister relation is the composition of the classical cousin relation between the round 3-spheres and \mathbb{R}^3 and of the conjugation by a phase θ in the associate family. The hyperbolic 3-space does not appear in this classification since it is not a fibration over a 2-manifold of constant curvature.

Remark 5.9. A classical problem in the theory of minimal surfaces is the question of the existence of minimal isometric deformations of a given minimal surface. The compatibility equations show that an associated family of a given minimal surface (i.e., a one-parameter family of minimal isometric deformation of this surface obtained by rotating the shape operator) in a homogeneous 3-manifold \mathbb{E} when $\tau \neq 0$ cannot be obtained in a simple way as in \mathbb{S}^3 , \mathbb{R}^3 , \mathbb{H}^3 , $\mathbb{S}^2 \times \mathbb{R}$ or $\mathbb{H}^2 \times \mathbb{R}$ (see [Dan04]). Indeed, if the quadruple (ds^2, S, T, ν) satisfies the compatibility equations for \mathbb{E} , then, in general, the quadruple $(ds^2, e^{\theta J}S, e^{\theta J}T, \nu)$ where $\theta \in \mathbb{R} \setminus 2\pi\mathbb{Z}$ does not. The question of the existence of the associate family for minimal surfaces in \mathbb{E} when $\tau \neq 0$ remains open.

5.2. Twin immersions. In this section we will study the special case of sister immersions lying in the same homogeneous 3-manifold. They necessarily have opposite mean curvatures.

Theorem 5.10. *Let \mathbb{E} be a homogeneous 3-manifold with a 4-dimensional isometry group, of base curvature κ and bundle curvature τ . Let ξ be its vertical vector field.*

Let Σ be a simply connected Riemann surface and let $x : \Sigma \rightarrow \mathbb{E}$ be a conformal constant mean curvature $H \neq 0$ immersion. Let N be the induced normal (compatible with the orientation of Σ). Let S be the symmetric operator on Σ induced by the shape operator of $x(\Sigma)$ associated to the normal N . Let T be the vector field on Σ such that $dx(T)$ is the projection of ξ onto $T(x(\Sigma))$. Let $\nu = \langle N, \xi \rangle$. Let

$$\theta = -2 \arctan \frac{H}{\tau}.$$

Then there exists a unique conformal immersion $\hat{x} : \Sigma \rightarrow \mathbb{E}$ such that:

- (1) *the metrics induced on Σ by x and \hat{x} are the same,*
- (2) *the symmetric operator on Σ induced by the shape operator of $\hat{x}(\Sigma)$ is $\tilde{S} = e^{\theta J}(S - HI) - HI = e^{\theta J}(S - \tau J) + \tau J$,*
- (3) *$\xi = d\hat{x}(e^{\theta J}T) + \nu \hat{N}$ where \hat{N} is the unit normal to \hat{x} .*

Moreover, this immersion \hat{x} is unique up to isometries of \mathbb{E} preserving the orientations of both the fibers and the base of the fibration, and it has constant mean curvature $-H$.

It is called the twin immersion of the immersion x .

Proof. This is a particular case of theorem 5.2 with $\mathbb{E}_1 = \mathbb{E}_2 = \mathbb{E}$, $\tau_1 = -\tau_2 = \tau$, $H_1 = -H_2 = H$. It suffices to check that the phase θ satisfies $\tau - iH = e^{i\theta}(\tau + iH)$.

The equivalence of the two expressions of \tilde{S} is a consequence of (14). \square

We notice that when $\tau \rightarrow 0$, then $\theta \rightarrow \pi$, i.e., $\tilde{T} \rightarrow -T$, and also $\tilde{S} \rightarrow -S$. This limit corresponds to the image of the initial surface by a horizontal symmetry in $\mathbb{M}^2(\kappa) \times \mathbb{R}$.

Moreover, we notice that the twin surface of a multigraph (over a part of the base of the fibration) is also a multigraph (since a surface is a multigraph if and only if ν does not vanish).

This suggests that the twin surface could be used to get an Alexandrov reflection-type principle in homogeneous manifolds with non-vanishing bundle curvature, since there is no Alexandrov reflection principle (see [Ale62]) in these manifolds (the horizontal and vertical ‘‘symmetries’’ are not isometries). Such an Alexandrov reflection principle would be very useful for the theory of CMC surfaces in homogeneous manifolds, in particular for proving that any closed embedded CMC surface in the Heisenberg space or in $\widetilde{\text{PSL}}_2(\mathbb{R})$ is a rotational sphere (this was proved for CMC surfaces in \mathbb{R}^3 , \mathbb{H}^3 , a 3-hemisphere, $\mathbb{H}^2 \times \mathbb{R}$ and a 2-hemisphere cross \mathbb{R} using the Alexandrov reflection principle).

We now give some examples of twin surfaces in the Heisenberg space Nil_3 with its standard metric (i.e., $\kappa = 0$, $\tau = \frac{1}{2}$). We will use the exponential coordinates described in section 2.3. Figueroa, Mercuri and Pedrosa classified CMC surfaces in Nil_3 invariant by a one-parameter family of translations or rotations (see [FMP99]; note that in their article the mean curvature is defined as the trace of the shape operator, whereas in this paper it is defined as the half of the trace). We will compute the twin surfaces of these examples. We will denote between parentheses $(\)$ the coordinates of a vector in the coordinate frame $(\partial_x, \partial_y, \partial_z)$, and between brackets $[\]$ the coordinates of a vector in the canonical frame (E_1, E_2, E_3) .

Example 5.11 (translational tubes). Let $H > 0$. The map

$$\varphi : (u, v) \mapsto \begin{pmatrix} u \\ \frac{\cos v}{2H} \\ u \frac{\cos v}{4H} + \frac{1}{4H} f(v) \end{pmatrix},$$

with

$$f(v) = \sqrt{1 + \frac{\cos^2 v}{4H^2}} \sin v + \frac{1 + 4H^2}{2H} \arcsin \left(\frac{\sin v}{\sqrt{1 + 4H^2}} \right),$$

for $(u, v) \in \mathbb{R}^2$, is a CMC H immersion defining a surface which is invariant by horizontal translations in the x -direction. This surface is an annulus, and it is a bigraph over a part of the minimal surface of equation $z = \frac{xy}{2}$; moreover it is “symmetric” with respect to this minimal surface.

We have

$$\begin{aligned}\varphi_u &= \begin{pmatrix} 1 \\ 0 \\ \frac{\cos v}{4H} \end{pmatrix} = \begin{bmatrix} 1 \\ 0 \\ \frac{\cos v}{2H} \end{bmatrix}, \\ \varphi_v &= \begin{pmatrix} 0 \\ -\frac{\sin v}{2H} \\ -u\frac{\sin v}{4H} + \frac{1}{4H}f'(v) \end{pmatrix} = \begin{bmatrix} 0 \\ -\frac{\sin v}{2H} \\ \frac{1}{4H}f'(v) \end{bmatrix}, \\ f'(v) &= 2 \cos v \sqrt{1 + \frac{\cos^2 v}{4H^2}},\end{aligned}$$

and so

$$\begin{aligned}\langle \varphi_u, \varphi_u \rangle &= 1 + \frac{\cos^2 v}{4H^2}, \\ \langle \varphi_v, \varphi_v \rangle &= \frac{1}{4H^2} \left(1 + \frac{\cos^4 v}{4H^2} \right), \\ \langle \varphi_u, \varphi_v \rangle &= \frac{\cos^2 v}{4H^2} \sqrt{1 + \frac{\cos^2 v}{4H^2}}.\end{aligned}$$

The unit normal vector is given by $N = \frac{\varphi_u \times \varphi_v}{\|\varphi_u \times \varphi_v\|}$; we compute that

$$\nu = -\frac{\sin v}{\sqrt{1 + \frac{\cos^4 v}{4H^2}}}.$$

We have

$$\begin{aligned}\langle T, \partial_u \rangle &= \langle \xi, \varphi_u \rangle = \frac{\cos v}{2H}, \\ \langle T, \partial_v \rangle &= \langle \xi, \varphi_v \rangle = \frac{\cos v}{2H} \sqrt{1 + \frac{\cos^2 v}{4H^2}},\end{aligned}$$

We notice that $\nu(u_1, -v) = -\nu(u_2, v)$ for all (u_1, u_2, v) . This indicates that the twin immersion could be an orientation-reversing reparametrization of the surface. For this reason we set

$$\tilde{\varphi} : (u, v) \mapsto \varphi(u + h(v), -v) = \begin{pmatrix} u + h(v) \\ \frac{\cos v}{2H} \\ (u + h(v))\frac{\cos v}{4H} - \frac{1}{4H}f(v) \end{pmatrix}$$

where h is a function. This is a CMC $-H$ immersion defining globally the same surface as φ . We compute that

$$\tilde{\varphi}_u = \begin{bmatrix} 1 \\ 0 \\ \frac{\cos v}{2H} \end{bmatrix}, \quad \tilde{\varphi}_v = \begin{bmatrix} h'(v) \\ -\frac{\sin v}{2H} \\ h'(v)\frac{\cos v}{2H} - \frac{1}{4H}f'(v) \end{bmatrix},$$

and so

$$\begin{aligned}\langle \tilde{\varphi}_u, \tilde{\varphi}_u \rangle &= 1 + \frac{\cos^2 v}{4H^2}, \\ \langle \tilde{\varphi}_v, \tilde{\varphi}_v \rangle &= \left(1 + \frac{\cos^2 v}{4H^2}\right) h'(v)^2 - \frac{\cos^2 v}{2H^2} h'(v) \sqrt{1 + \frac{\cos^2 v}{4H^2}} \\ &\quad + \frac{1}{4H^2} \left(1 + \frac{\cos^4 v}{4H^2}\right), \\ \langle \tilde{\varphi}_u, \tilde{\varphi}_v \rangle &= \left(1 + \frac{\cos^2 v}{4H^2}\right) h'(v) - \frac{\cos^2 v}{4H^2} \sqrt{1 + \frac{\cos^2 v}{4H^2}}.\end{aligned}$$

Thus $\tilde{\varphi}$ induces on \mathbb{R}^2 the same metric as φ if and only if

$$h'(v) = \frac{\cos^2 v}{2H^2 \sqrt{1 + \frac{\cos^2 v}{4H^2}}}.$$

We now assume that this condition is satisfied; we can also assume that $h(0) = 0$. The function h is increasing. We have

$$\begin{aligned}\tilde{\nu} &= \nu, \\ \langle \tilde{T}, \partial_u \rangle &= \langle \xi, \tilde{\varphi}_u \rangle = \frac{\cos v}{2H}, \\ \langle \tilde{T}, \partial_v \rangle &= \langle \xi, \tilde{\varphi}_v \rangle = \frac{\cos v}{2H \sqrt{1 + \frac{\cos^2 v}{4H^2}}} \left(\frac{\cos^2 v}{4H^2} - 1 \right).\end{aligned}$$

The direct orthonormal frame (e_1, e_2) obtained from the frame (∂_u, ∂_v) by the Gram-Schmidt process satisfies

$$\begin{aligned}e_1 &= \frac{\partial_u}{\|\partial_u\|}, \\ e_2 &= \frac{-\langle \partial_u, \partial_v \rangle \partial_u + \|\partial_u\|^2 \partial_v}{\|\partial_u\| \sqrt{\|\partial_u\|^2 \|\partial_v\|^2 - \langle \partial_u, \partial_v \rangle^2}}.\end{aligned}$$

A computation gives

$$\|\partial_u\|^2 \|\partial_v\|^2 - \langle \partial_u, \partial_v \rangle^2 = \frac{1}{4H^2} \left(1 + \frac{\cos^2 v}{4H^2}\right).$$

Thus we get

$$\begin{aligned}e_1 &= \frac{1}{\sqrt{1 + \frac{\cos^2 v}{4H^2}}} \partial_u, \\ e_2 &= -\frac{\cos^2 v}{2H \sqrt{1 + \frac{\cos^2 v}{4H^2}}} \partial_u + 2H \partial_v.\end{aligned}$$

So we have

$$T = \frac{\cos v}{\sqrt{1 + \frac{\cos^2 v}{4H^2}}} \left(\frac{1}{2H} e_1 + e_2 \right),$$

$$\tilde{T} = \frac{\cos v}{\sqrt{1 + \frac{\cos^2 v}{4H^2}}} \left(\frac{1}{2H} e_1 - e_2 \right).$$

Let $\theta = -2 \arctan(2H)$. Then we have

$$\cos \theta = \frac{1 - 4H^2}{1 + 4H^2}, \quad \sin \theta = -\frac{4H}{1 + 4H^2}.$$

Since $Je_1 = e_2$ and $Je_2 = -e_1$, we get

$$e^{\theta J} T = \tilde{T}.$$

Finally, the compatibility equation (10) implies that

$$\tilde{S} = e^{\theta J} (S - \tau J) + \tau J$$

at points where $\nu \neq 0$; and by continuity this identity holds everywhere. This proves that $\tilde{\varphi}$ is the twin immersion of φ .

Thus the translational tube is *globally* invariant by the twin relation, but it is *not pointwise* invariant: the correspondence is

$$\varphi(u, v) \mapsto \varphi(u + h(v), -v).$$

Geometrically, this correspondence maps a point of the tube to the other point of the tube lying in the same fiber and then translates it by $h(v)$ in the x -direction. In particular, the closed curve $v \mapsto \varphi(u_0, v)$ is mapped to the curve $v \mapsto \varphi(u_0 + h(v), -v)$, which is *not* closed.

Example 5.12 (rotational spheres). Let $H > 0$. The map

$$\varphi : (u, v) \mapsto \begin{pmatrix} \frac{1}{H} \cos u \cos v \\ \frac{1}{H} \sin u \cos v \\ \frac{1}{2H} f(v) \end{pmatrix},$$

with f as in example 5.11, for $(u, v) \in \mathbb{R} \times (-\frac{\pi}{2}, \frac{\pi}{2})$, is a CMC $-H$ immersion defining a rotational sphere minus the top and bottom points (the normal of the immersion points outside whereas the mean curvature vector points inside). It is a bigraph over a part of the minimal surface of equation $z = 0$; moreover it is “symmetric” with respect to this minimal surface.

We have

$$\varphi_u = \frac{1}{H} \begin{bmatrix} -\sin u \cos v \\ \cos u \cos v \\ -\frac{1}{2H} \cos^2 v \end{bmatrix}, \quad \varphi_v = \frac{1}{H} \begin{bmatrix} -\cos u \sin v \\ -\sin u \sin v \\ \frac{1}{2} f'(v) \end{bmatrix},$$

and so

$$\begin{aligned} \langle \varphi_u, \varphi_u \rangle &= \frac{\cos^2 v}{H^2} \left(1 + \frac{\cos^2 v}{4H^2} \right), \\ \langle \varphi_v, \varphi_v \rangle &= \frac{1}{H^2} \left(1 + \frac{\cos^2 v}{4H^2} \right), \\ \langle \varphi_u, \varphi_v \rangle &= -\frac{\cos^3 v}{2H^3} \sqrt{1 + \frac{\cos^2 v}{4H^2}}. \end{aligned}$$

The unit normal vector is given by $N = \frac{\varphi_u \times \varphi_v}{\|\varphi_u \times \varphi_v\|}$; we compute that

$$\nu = \frac{\sin v}{\sqrt{1 + \frac{\cos^4 v}{4H^2}}}.$$

We have

$$\begin{aligned} \langle T, \partial_u \rangle &= \langle \xi, \varphi_u \rangle = -\frac{\cos^2 v}{2H^2}, \\ \langle T, \partial_v \rangle &= \langle \xi, \varphi_v \rangle = \frac{\cos v}{H} \sqrt{1 + \frac{\cos^2 v}{4H^2}}. \end{aligned}$$

Let

$$\tilde{\varphi} : (u, v) \mapsto \varphi(u + g(v), -v) = \begin{pmatrix} \frac{1}{H} \cos(u + g(v)) \cos v \\ \frac{1}{H} \sin(u + g(v)) \cos v \\ -\frac{1}{2H} f(v) \end{pmatrix}$$

where g is a function. This is a CMC H immersion defining globally the same surface as φ . We compute that

$$\begin{aligned} \tilde{\varphi}_u &= \frac{1}{H} \begin{bmatrix} -\sin(u + g(v)) \cos v \\ \cos(u + g(v)) \cos v \\ -\frac{1}{2H} \cos^2 v \end{bmatrix}, \\ \tilde{\varphi}_v &= \frac{1}{H} \begin{bmatrix} -\cos(u + g(v)) \sin v - g'(v) \sin(u + g(v)) \cos v \\ -\sin(u + g(v)) \sin v + g'(v) \cos(u + g(v)) \cos v \\ -\frac{1}{2} f'(v) - \frac{1}{2H} g'(v) \cos^2 v \end{bmatrix}, \end{aligned}$$

and thus $\tilde{\varphi}$ induces on $\mathbb{R} \times (\frac{\pi}{2}, \frac{\pi}{2})$ the same metric as φ if and only if

$$g'(v) = -\frac{\cos v}{H \sqrt{1 + \frac{\cos^2 v}{4H^2}}}.$$

We now assume that this condition is satisfied; we can also assume that $g(0) = 0$. The function g is odd and 2π -periodic. We have

$$\begin{aligned} \tilde{\nu} &= \nu, \\ \langle \tilde{T}, \partial_u \rangle &= \langle \xi, \tilde{\varphi}_u \rangle = -\frac{\cos^2 v}{2H^2}, \\ \langle \tilde{T}, \partial_v \rangle &= \langle \xi, \tilde{\varphi}_v \rangle = \frac{\cos v}{H \sqrt{1 + \frac{\cos^2 v}{4H^2}}} \left(\frac{\cos^2 v}{4H^2} - 1 \right). \end{aligned}$$

The direct orthonormal frame (e_1, e_2) obtained from the frame (∂_u, ∂_v) by the Gram-Schmidt process satisfies

$$\begin{aligned} e_1 &= \frac{H}{\cos v \sqrt{1 + \frac{\cos^2 v}{4H^2}}} \partial_u, \\ e_2 &= -\frac{\cos v}{2\sqrt{1 + \frac{\cos^2 v}{4H^2}}} \partial_u + H \partial_v. \end{aligned}$$

So we have

$$T = \frac{\cos v}{\sqrt{1 + \frac{\cos^2 v}{4H^2}}} \left(-\frac{1}{2H} e_1 + e_2 \right),$$

$$\tilde{T} = \frac{\cos v}{\sqrt{1 + \frac{\cos^2 v}{4H^2}}} \left(-\frac{1}{2H} e_1 - e_2 \right).$$

Let $\theta = 2 \arctan(2H)$. We check as in example 5.11 that

$$e^{\theta J} T = \tilde{T},$$

$$\tilde{S} = e^{\theta J} (S - \tau J) + \tau J.$$

This proves that $\tilde{\varphi}$ is the twin immersion of φ .

Thus the rotational sphere is *globally* invariant by the twin relation, but it is *not pointwise* invariant: the correspondence is

$$\varphi(u, v) \mapsto \varphi(u + g(v), -v).$$

Geometrically, this correspondence maps a point of the sphere to the other point of the sphere lying in the same fiber and then rotates it by the angle $g(v)$ about the z -axis. In particular, the circle $v \mapsto \varphi(u_0, v)$ lying in a vertical plane is mapped to the curve $v \mapsto \varphi(u_0 + g(v), -v)$, which is closed but not contained in a vertical plane.

REFERENCES

- [Abr04] U. Abresch. Generalized Hopf differentials. To appear in Proceedings of the 13th School of Differential Geometry, *Mat. Contemp.*, 2004.
- [Ale62] A. D. Alexandrov. A characteristic property of spheres. *Ann. Mat. Pura Appl.*, 58:303–315, 1962.
- [AR03] U. Abresch and H. Rosenberg. The Hopf differential for constant mean curvature surfaces in $\mathbb{S}^2 \times \mathbb{R}$ and $\mathbb{H}^2 \times \mathbb{R}$. To appear in *Acta Math.*, 2003.
- [Bry87] R. Bryant. Surfaces of mean curvature one in hyperbolic space. *Astérisque*, 154–155:321–347, 1987.
- [Car92] M. do Carmo. *Riemannian Geometry*. Birkhäuser, 1992.
- [Dan03] B. Daniel. Minimal disks bounded by three straight lines in Euclidean space and trinions in hyperbolic space. Preprint, math.DG/0307066, 2003.
- [Dan04] B. Daniel. Isometric immersions into $\mathbb{S}^n \times \mathbb{R}$ and $\mathbb{H}^n \times \mathbb{R}$ and applications to minimal surfaces. Preprint, math.DG/0406426, 2004.
- [FMP99] C. Figueroa, F. Mercuri, and R. Pedrosa. Invariant surfaces of the Heisenberg groups. *Ann. Mat. Pura Appl. (4)*, 177:173–194, 1999.
- [Kar01] H. Karcher. Hyperbolic constant mean curvature one surfaces with compact fundamental domains. Preprint, 2001.
- [NR04] B. Nelli and H. Rosenberg. Global properties of constant mean curvature surfaces in $\mathbb{H}^2 \times \mathbb{R}$. To appear in *Pacific J. Math.*, 2004.
- [Pet98] P. Petersen. *Riemannian Geometry*. Springer-Verlag, 1998.
- [Sco83] P. Scott. The geometries of 3-manifolds. *Bull. London Math. Soc.*, 15(5):401–487, 1983.
- [Ten71] K. Tenenblat. On isometric immersions of Riemannian manifolds. *Bol. Soc. Brasil. Mat.*, 2(2):23–36, 1971.
- [UY93] M. Umehara and K. Yamada. Complete surfaces of constant mean curvature 1 in the hyperbolic 3-space. *Ann. of Math. (2)*, 137:611–638, 1993.

IMPA, ESTRADA DONA CASTORINA 110, 22460-320 RIO DE JANEIRO - RJ, BRAZIL

E-mail address: `bdaniel@impa.br`

URL: `http://www.math.jussieu.fr/~daniel`