

**ON THE LIMIT OF FAMILIES OF ALGEBRAIC SUBVARIETIES
WITH UNBOUNDED VOLUME
(DRAFT)**

CÉSAR CAMACHO AND LUIZ HENRIQUE DE FIGUEIREDO

ABSTRACT. We prove that the limit of a sequence of generic semi-algebraic sets given by a finite number of formulas always exists and is a semi-algebraic set that can be explicitly given as a Boolean expression involving the primitives of the additive forms of the formulas.

1. INTRODUCTION

Bishop [2] proved that the limit set of a sequence of complex purely k -dimensional algebraic subvarieties whose real volumes are uniformly bounded is again a purely k -dimensional algebraic subvariety. On the other hand, there are many reasons why one should be interested in analyzing the limit sets of algebraic subvarieties with unbounded volume. One reason is the existence of families of algebraic curves of increasing degree that are integrals of families of polynomials differential equations on the plane with bounded degree, a badly understood phenomenon related to the sixteenth Hilbert Problem (see [3], for instance). Another reason is that, despite the existence of topologically complicated limit sets of curves with unbounded volume (see [5], for instance), much can be said about the limit sets of algebraic subvarieties which lie in a family of subvarieties with finite complexity (see [4] for a definition of this concept).

In this paper we consider the limit sets of one-parameter families of algebraic subvarieties, indexed by a natural number n , defined by a finite number of equations, each one defined by a *formula*. Associated to each formula there is a *height*, which is the maximum number of nested n -th powers that appear in it. Here is the formal definition:

Definition 1. Formulas and their heights are defined recursively as follows:

- (1) Every $F \in \mathbb{C}[X_1, \dots, X_m]$ is a formula of height zero.
- (2) If F_1 and F_2 are formulas then $F_1 + F_2$ and $F_1 F_2$ are formulas of height $\max(h_1, h_2)$, where h_i is the height of F_i .
- (3) If F is a formula of height h , then F^n is a formula of height $h + 1$.

A formula of height zero is also called a *primitive* formula; it is simply a complex polynomial.

The height is a measure of the complexity of the formula: it measures how the degree increases with n . A formula of height h has degree proportional to n^h . An example of a formula of height 3 is

$$(((x^2 - y + 1)^n - 1)^n + x)^n + (xy)^n + (y^n - 1)^2 + 1.$$

2000 *Mathematics Subject Classification.* Primary 14P10; Secondary 14P15.

Note that the degree is $2n^3$.

The same polynomial family may be given by different formulas. For instance,

$$(x^n + y)^2 = (x^n)^2 + 2x^n y + y^2.$$

So, we shall need a normal form for formulas. A formula is in *additive form* when it can be expressed as

$$Q_1 A_1^n + Q_2 A_2^n + \cdots + Q_l A_l^n - P,$$

where Q_1, \dots, Q_l , and P are primitive formulas and A_1, \dots, A_l are arbitrary subformulas (necessarily of smaller height than the original formula). Additive forms are normal forms, as the next lemma shows:

Lemma 1. *Every formula can be written in additive form.*

Proof. The proof is by induction on the number of operations required to obtain the formula according to Definition 1. If F is a primitive formula, then we can take $l = 0$ and $P = -F$. If $F = A^n$, then F is already in additive form because we can take $l = 1$, $Q_1 = 1$, $A_1 = A$, and $P = 0$. If $F = A + B$, then by induction A and B can be expressed in additive form, whose combination gives an additive form for F . If $F = AB$, then again by induction A and B can be expressed in additive form. By performing the multiplication AB on their additive forms, we get an additive form for F . \square

As an example, $(x^n + y)^2$ can be written in additive form as $(x^2)^n + (2y)x^n + y^2$.

Definition 2. The *limit* (as $n \rightarrow \infty$) of a sequence (Ω_n) of subsets of \mathbb{C}^m is the set $\lim \Omega_n$ of points that are limits of sequences of points lying in a subsequence of (Ω_n) . More precisely,

$$\lim \Omega_n = \{ z \in \mathbb{C}^m : \exists(z_n), z_n \rightarrow z, \exists(k_n), k_n \rightarrow \infty, z_n \in \Omega_{k_n} \text{ for sufficiently large } n \}.$$

Thus, according to our definition, the family of real curves $x^{2n} + y^{2n} = 1$ converges to the border of the unit square given by $x^2 \leq 1$, $y^2 \leq 1$. Actually, this definition applies to the curves $x^n + y^n = 1$ (note that we now allow both even and odd exponents). These curves converge to the union of the border of the unit square with the two rays given by $x = -y$, $x^2 \geq 1$ (the curves actually alternate between two limit sets, but our definition of limit covers this). Considered as a family of complex curves, the family $x^n + y^n = 1$ has as limit set the subset of \mathbb{C}^2 given by $\partial(|x| < 1 \cap |y| < 1) \cup [|x| = |y| > 1]$, as it is easy to verify.

We shall consider two situations: limit sets in \mathbb{R}^k of families of algebraic subvarieties given by a finite number of formulas and limit sets in \mathbb{C}^k of families of complex algebraic subvarieties.

In the real case it turns out that it is easier to describe the limits of semi-algebraic subsets, instead of algebraic subsets. An *algebraic subvariety* of codimension 1 is the set of points that satisfy a polynomial equation $f(z) = 0$. For simplicity, we shall write this set as $[f = 0]$. We shall also deal with *basic closed semi-algebraic subsets*, which are the solutions of a system of polynomial inequalities: $[f_1 \geq 0, \dots, f_k \geq 0]$, and with *basic open semi-algebraic subsets*, which are given by strict inequalities: $[f_1 > 0, \dots, f_k > 0]$.

One main difficulty in the theory of semi-algebraic sets is that the closure of a basic open semi-algebraic set is not necessarily the corresponding basic closed semi-algebraic set obtained by relaxing the strict inequalities. Nor is the interior of

a closed semi-algebraic set equal to the corresponding basic open semi-algebraic set obtained by restricting the inequalities. However, these are true generically, in two senses: (i) they are true if we perturb the polynomials slightly, and (ii) relaxing or restricting the inequalities only adds or removes lower dimensional components. So, we say that a basic closed semi-algebraic set is *generic* when it coincides with the closure of the corresponding basic open semi-algebraic set obtained by restricting the inequalities. In other words, a basic closed semi-algebraic set given by $[f_1 \geq 0, \dots, f_k \geq 0]$ is generic when $[f_1 \geq 0, \dots, f_k \geq 0] = \text{closure}[f_1 > 0, \dots, f_k > 0]$. A *generic algebraic* set is, by definition, the boundary of a generic semi-algebraic subset.

Our main result is the following:

Theorem 1. *The limit of a sequence of generic semi-algebraic sets given by a finite number of formulas always exists and is a semi-algebraic set that can be explicitly given as a Boolean expression involving the primitives of the additive forms of the formulas.*

The corresponding algebraic version is also valid:

Theorem 2. *The limit of a sequence of generic algebraic sets given by a finite number of formulas always exists and is an algebraic set that can be explicitly given as a Boolean expression involving the primitives of the additive forms of the formulas.*

In the complex case the limit set of a family of algebraic sets given by a finite number of formulas has also an underlying semi-algebraic structure in the sense that it projects, by means of a rational map, onto a proper real semi-algebraic subset defined by expressions involving the absolute values of the primitives of the formulas. More precisely, we have the following result:

Theorem 3. *The limit of a sequence of generic algebraic subsets given by a finite number of formulas with complex coefficients always exists; it is a subset with a complex structure obtained by means of a rational pull-back on semi-algebraic subsets defined explicitly in terms of Boolean expressions involving the absolute values of the primitives of the formulas.*

Here is an example of this situation, which generalizes the $x^n + y^n = 1$ example given above. Let A_1 , A_2 , and P be polynomials. Then

$$\lim[A_1^n + A_2^n = P] = \partial([|A_1| < 1] \cap [|A_2| < 1] \cap [P \neq 0]) \cup [|A_1| = |A_2| > 1]$$

This limit can be also understood as the pull-back by the polynomial map

$$(A_1, A_2): \mathbb{C}^2 \rightarrow \mathbb{C}^2$$

of the Reinhardt preimage of the semi-algebraic subset of \mathbb{R}^2 given by the second member of the equation above, where the axes of \mathbb{R}^2 are taken as $|A_1|$ and $|A_2|$.

2. THE REAL CASE

We start with the simplest cases. *We assume that all semi-algebraic sets are generic.*

Let A and P be real polynomials. We want to describe the limit of the algebraic subsets $[A^{2n} = P]$. As mentioned before, it is simpler to state the results for the semi-algebraic sets $\Omega_n = [A^{2n} \leq P]$. Therefore, we shall describe $\Omega_\infty = \lim \Omega_n$.

Lemma 2. $\lim[A^{2n} \leq P] = [A^2 \leq 1, P \geq 0]$.

Proof. Let $\Omega_n = [A^{2n} \leq P]$, $\Omega_\infty = \lim \Omega_n$, and $\Omega = [A^2 \leq 1, P \geq 0]$. We shall show that $\Omega_\infty = \Omega$.

To show that $\Omega_\infty \subseteq \Omega$, take $z \in \Omega_\infty$. Then, by definition of Ω_∞ , we have $z = \lim z_n$ and $k_n \rightarrow \infty$, with $z_n \in \Omega_{k_n}$, that is, $A^{2k_n}(z_n) \leq P(z_n)$. Since $A^{2k_n}(z_n) \geq 0$, we get $P(z_n) \geq 0$ and hence $P(z) \geq 0$, because P is continuous. Moreover, $(P(z_n))$ is a bounded sequence because it converges, and so $P(z_n) \leq L$ for some $L \geq 0$. This implies that $A(z_n)^2 \leq P(z_n)^{1/k_n} \leq L^{1/k_n}$. Since (L^{1/k_n}) converges to either 0 or 1 (according to whether $L = 0$ or $L \neq 0$), we conclude that $A(z)^2 = \lim A(z_n)^2 \leq \lim L^{1/k_n} \leq 1$. Hence, $z \in \Omega$.

To show that $\Omega \subseteq \Omega_\infty$, take $z \in \Omega$. Since Ω is generic, we have that $z = \lim z_n$, with $z_n \in [A^2 < 1, P > 0]$. From $A(z_n)^2 < 1$ we get that $A(z_n)^{2k} \rightarrow 0$ as $k \rightarrow \infty$. Since $P(z_n) > 0$, there is k_n such that $A(z_n)^{2k_n} < P(z_n)$, that is, $z_n \in \Omega_{k_n}$. By definition, this means that $z \in \Omega_\infty$. (Increase k_n beyond n if necessary to get $k_n \rightarrow \infty$, as required.) \square

The genericity hypothesis is essential to the lemma as stated. Although the proof of the lemma shows that $\Omega_\infty \subseteq \Omega$ in all cases, the reverse inclusion is not always true because things are more complicated in the general case. We give here an example just to give a taste of this complication. Let $A = y(y-1)^2 + 1$ and $P = x(x+1)$. Then $\lim[A^n \leq P]$ is shown in Figure 1. Note that $[P \geq 0]$ is not the closure of $[P > 0]$ because $[P \geq 0]$ contains the line $[x = 1]$, which is not in the closure of $[P > 0]$ since P is negative near $x = -1$ (). Note that $[A = 1, P \geq 0]$ is not completely contained in $\lim[A^n \leq P]$; only $[A = 1, P \geq 1]$ is part of the limit set.

In general, $\lim[A^n \leq P]$ is equal to $\lim[A^{2n} \leq P]$, except that $P \geq 1$ when $A = 1^+$ and $A = 0$ when $P = 0^-$ ().

*** Figure 1 here

The next lemma generalizes Lemma 2:

Lemma 3. *Let A_1, \dots, A_k and P be polynomials. If $[A_1^2 \leq 1, \dots, A_k^2 \leq 1, P \geq 0]$ is generic, then it is equal to $\lim[A_1^{2n} + \dots + A_k^{2n} \leq P]$.*

Proof. The proof is essentially the same as that of Lemma 2. To show that $\Omega_\infty \subseteq \Omega$, just note that $A_i(z_n)^{2n} \leq A_1(z_n)^{2n} + \dots + A_k(z_n)^{2n} \leq P(z_n)$. To show that $\Omega \subseteq \Omega_\infty$, just note that $A_i(z_n)^{2r} \rightarrow 0$ as $r \rightarrow \infty$ implies that $A_i(z_n)^{2r} < P(z_n)/k$ for sufficiently large r . The rest of the proof follows as before. \square

The next lemma generalizes Lemma 2 in a different direction:

Lemma 4. *Let A_n be a formula of positive height and P be a polynomial. Then $\lim[A_n^{2n} \leq P] = \lim[A_n^2 \leq 1] \cap [P \geq 0]$.*

Proof. Take $z \in \lim[A_n^{2n} \leq P]$. By definition, there are sequences $z_n \rightarrow z$ and $k_n \rightarrow \infty$ such that $A_{k_n}(z_n)^{2k_n} \leq P(z_n)$. Clearly, $P(z) = \lim P(z_n) \geq 0$. As before, since $(P(z_n))$ is bounded, we have $A_{k_n}^{2k_n}(z_n) \leq L$ for some $L > 0$.

This implies $A_{k_n}^2(z_n) \leq L^{1/k_n}$ and since $\lim[A_n^2 \leq 1] = \lim[L^{-1/n} A_n^2 \leq 1]$ we obtain $z \in \lim[A_n^2 \leq 1] \cap [P \geq 0]$. Suppose now that $z \in \lim[A_n^2 \leq 1] \cap [P \geq 0]$. Since $[P \geq 0]$ is generic, we may assume that $P > 0$. Then there are sequences $z_n \rightarrow z$ and $k_n \rightarrow \infty$ such that $A_{k_n}(z_n)^2 \leq 1$ and $P(z_n) > 0$. Since $(A_{k_n}^{2k_n}/P(z_n))$ is bounded we have $A_{k_n}(z_n)^{2k_n} \leq LP(z_n)$, for some $L > 0$. Since $\lim[A_n^{2n} \leq LP] = \lim[A_n^{2n} \leq P]$ we conclude that $z \in \lim[A_n^{2n} \leq P]$. \square

Lemma 5. *Suppose that A_n is a formula of positive height h and P is a primitive formula. Then, $\lim[A_n^{2n} \geq P] = [P \leq 0] \cup [\lim[A_n^2 \geq 1], P \geq 0]$*

Proof. Take $z \in \lim[A_n^{2n} \geq P]$. Then $z = \lim z_n$ and there is $k_n \rightarrow \infty$ such that $A_{k_n}^{2k_n}(z_n) \geq P(z_n)$. So, either $P(z) \leq 0$, or $P(z) > 0$ and $A_{k_n}^2 \geq P(z_n)^{1/k_n}$. Since $\lim[P^{-1/n} A_n^2 \geq 1] = \lim[A_n^2 \geq 1]$, we obtain $z \in [P \leq 0] \cup [\lim[A_n^2 \geq 1], [P \geq 0]]$. Conversely, take $z \in [\lim[A_n^2 \geq 1], [P \geq 0]]$. Since $[P \geq 0]$ is generic, we may assume $P(z_n) > 0$ and that there are sequences $z_n \rightarrow z$, and $k_n \rightarrow \infty$ such that $A_{k_n}(z_n)^2 \geq 1$ and $P(z_n) > 0$. The sequence $(A_{k_n}(z_n)^{2k_n}/P(z_n))$ is bounded below by $L > 0$, i.e., $A_{k_n}^{2k_n}(z_n) \geq LP(z_n)$. Since $\lim[A_n^{2n} \geq LP] = \lim[A_n^{2n} \geq P]$, we conclude that $z \in \lim[A_n^{2n} \geq P]$. \square

Lemma 6. *Suppose that A_n is a formula of positive height and P and Q are primitive formulas. Then*

$$\lim[QA_n^{2n} \leq P] = ([Q > 0] \cap \lim[A_n^{2n} \leq P]) \cup ([Q < 0] \cap \lim[A_n \geq -P]) \cup [Q = 0, P \geq 0].$$

Proof. If $Q(z) > 0$ and $z \in \lim[QA_n^{2n} \leq P]$, then there are sequences $z_n \rightarrow z$ and $k_n \rightarrow \infty$ such that $Q(z_n)A_{k_n}(z_n)^{2k_n} \leq P(z_n) \leq L$, with $L > 0$. Since $\lim[(QL^{-1})^{1/n} A_n^2 \leq 1] = \lim[A_n^2 \leq 1]$ and $P(z) \geq 0$, we obtain that $z \in [Q > 0] \cap \lim[A_n^{2n} \leq P]$. If $Q(z) < 0$, there are sequences $z_n \rightarrow z$, $k_n \rightarrow \infty$ such that $A_{k_n}(z_n)^{2k_n} \geq -P(z_n)/Q(z_n)$. By Lemma 5, either $-P(z)/Q(z) \leq 0$ or $z \in \lim[A_n^2 \geq 1, -P/Q \geq 0]$, or equivalently $P(z) \leq 0$ or $\lim[A_n^2 \geq 1, P(z) \geq 0]$, i.e., $z \in \lim[A_n^{2n} \geq -P]$. \square

By setting $Q = A$ in the limit above, we get an expression for $\lim[A^{2n+1} \leq P]$, and from this an expression for $\lim[A^n \leq P]$, which should convince the reader that restricting to even powers is a good thing.

Lemma 7. *Suppose that A_n and B_n are formulas of positive height and let P be a primitive formula. Then*

$$\lim[A_n^{2n} \leq P + B_n^{2n}] = \lim[B_n^2 < 1] \cap \lim[A_n^{2n} \leq P] \cup \lim[B_n^2 \geq 1] \cap \lim[A_n^{2n} \leq B_n^{2n}]$$

Proof. Take $z \in \lim[A_n^{2n} \leq P + B_n^{2n}]$. Then, there are sequences $z_n \rightarrow z$ and $k_n \rightarrow \infty$ such that $A_{k_n}(z_n)^{2k_n} \leq P(z_n) + B_{k_n}(z_n)^{2k_n}$. If $\lim B_{k_n}^2 < 1$, then $B_{k_n}(z_n)^{2k_n} \rightarrow 0$ and we have $P(z) \geq 0$ and $A_{k_n}(z_n)^{2k_n} \leq L$, where L is a constant. Thus $A_{k_n}(z_n)^2 \leq L^{1/k_n}$ and so $z \in [P \geq 0] \cap \lim[A_n^2 \leq 1] = \lim[A_n^{2n} \leq P]$ by Lemma 4. So we get $z \in \lim[B_n^2 \leq 1] \cap \lim[A_n^{2n} \leq P]$. If $\lim B_{k_n}(z_n)^2 \geq 1$, then for n large $P(z_n) \leq KB_{k_n}(z_n)^{2k_n}$ for some constant $K > 0$. Thus $A_{k_n}(z_n)^{2k_n} \leq (K+1)B_{k_n}(z_n)^{2k_n}$, so $z \in \lim[A_n^{2n} \leq B_n^{2n}]$. On the other hand, if $z \in \lim[B_n^2 < 1] \cap \lim[A_n^{2n} \leq P]$, then there are sequences $z_n \rightarrow z, k_n \rightarrow \infty$ such that $A_{k_n}(z_n)^{2k_n} \leq P(z_n) \leq P(z_n) + B_{k_n}(z_n)^{2k_n}$. Moreover, if $z \in \lim[B_n^2 \geq 1] \cap \lim[A_n^{2n} \leq B_n^{2n}]$, then we have two possibilities: either $P(z) > 0$ and then $A_{k_n}(z_n)^{2k_n} \leq B_{k_n}(z_n)^{2k_n} \leq B_{k_n}(z_n)^{2k_n} + P(z_n)$ or $P(z) \leq 0$ and then for n large $B_{k_n}(z_n)^{2k_n} > -2P(z_n)$. Since $\lim[A_n^{2n} \leq B_n^{2n}] = \lim[2A_n^{2n} \leq B_n^{2n}]$ we can write $2A_{k_n}(z_n)^{2k_n} \leq B_{k_n}(z_n)^{2k_n} = 2B_{k_n}(z_n)^{2k_n} - B_{k_n}(z_n)^{2k_n} < 2B_{k_n}(z_n)^{2k_n} + 2P(z_n)$, i.e., $A_{k_n}(z_n)^{2k_n} \leq B_{k_n}(z_n)^{2k_n} + P(z_n)$. \square

Lemma 8. *Suppose that $A_1, \dots, A_k, B_1, \dots, B_l$ are formulas of positive height. Then*

$$\begin{aligned} \lim[A_1^{2n} + \dots + A_k^{2n} \leq P + B_1^{2n} + \dots + B_l^{2n}] &= \\ &= \bigcap_{i=1}^k \lim[A_i^{2n} \leq P + B_1^{2n} + \dots + B_l^{2n}] \\ &= \bigcap_{i=1}^k \bigcup_{j=1}^l \lim[A_i^{2n} \leq P + B_j^{2n}]. \end{aligned}$$

Proof. Define $P_1 := P + B_1^{2n} + \dots + B_l^{2n}$. We proceed to show that

$$\lim[A_1^{2n} + \dots + A_k^{2n} \leq P_1] = \bigcap_{i=1}^k \lim[A_i^{2n} \leq P_1].$$

Indeed, if $z \in \lim[A_1^{2n} + \dots + A_k^{2n} \leq P_1]$, since $A_i^{2n} \leq A_1^{2n} + \dots + A_k^{2n} \leq P_1$, we have $z \in \bigcap_{i=1}^k \lim[A_i^{2n} \leq P_1]$. On the other hand, if $z \in \bigcap_{i=1}^k \lim[A_i^{2n} \leq P_1]$, since $\lim[A_i^{2n} \leq P_1] = \lim[A_i^{2n} \leq (1/k)P_1]$, then $z \in \lim[A_1^{2n} + \dots + A_k^{2n} \leq P_1]$. We proceed to show now that

$$\lim[A^{2n} \leq P + B_1^{2n} + \dots + B_l^{2n}] = \bigcap_{j=1}^l \lim[A^{2n} \leq P + B_j^{2n}].$$

On one hand it is clear that $\lim[A^{2n} \leq P + B_j^{2n}] \subset \lim[A^{2n} \leq P + B_1^{2n} + \dots + B_l^{2n}]$. On the other hand, if $z \in \lim[A^{2n} \leq P + B_1^{2n} + \dots + B_l^{2n}]$, then we have a relation $B_{i_1}(z)^2 \leq \dots \leq B_{i_l}(z)^2$, where $i_1, \dots, i_l = 1, \dots, l$. Then $A(z_n)^{2k_n} \leq P(z_n) + B_1(z_n)^{2k_n} + \dots + B_l(z_n)^{2k_n} \leq P(z_n) + lB_{i_l}(z_n)^{2k_n}$, i.e., $z \in \lim[A^{2n} \leq P + lB_{i_l}^{2n}] = \lim[A^{2n} \leq P + B_{i_l}^{2n}]$ \square

The next lemma is similar to Lemma 6, and we leave its proof to the reader.

Lemma 9. *Suppose that $A_1, \dots, A_k, B_1, \dots, B_l$ are formulas of positive height and P is a primitive formula. Then*

$$\begin{aligned} \lim[Q_1 A_1^{2n} + \dots + Q_k A_k^{2n} \leq P + R_1 B_1^{2n} + \dots + R_l B_l^{2n}] \\ = \lim[A_1^{2n} + \dots + A_k^{2n} \leq P + B_1^{2n} + \dots + B_l^{2n}] \end{aligned}$$

provided that the primitive formulas $Q_1, \dots, Q_k, R_1, \dots, R_l$ are positive.

Proof of Theorem 1. By Lemma 1, a formula can be expressed in additive form and the question is reduced to determining

$$\lim[Q_1 A_1^{2n} + \dots + Q_k A_k^{2n} \leq P + R_1 B_1^{2n} + \dots + R_l B_l^{2n}],$$

where the Q_i 's and the R_j 's are positive, since the complete limit can be written as a finite union of expressions as above. On the other hand, by Lemma 9 it is enough to find $\lim[A_1^{2n} + \dots + A_k^{2n} \leq P + B_1^{2n} + \dots + B_l^{2n}]$, where $A_1^{2n} + \dots + A_k^{2n} - P - B_1^{2n} - \dots - B_l^{2n}$ is a formula of height $h \geq 1$. Since $\lim[A_1^{2n} + \dots + A_k^{2n} \leq P + B_1^{2n} + \dots + B_l^{2n}] = \bigcap_{i=1}^k \bigcup_{j=1}^l \lim[A_i^{2n} \leq P + B_j^{2n}]$, it is enough to find the limit of expressions of the type $\lim[A^{2n} \leq P + B^{2n}]$. Proceeding by induction on the height h of $A^n - B^n - P$, we have by Lemma 7 that for $h = 0$ $\lim[A^{2n} \leq P + B^{2n}] = [B^2 < 1] \cap [A^2 \leq 1] \cap [P \geq 0] \cup [B^2 \geq 1] \cap [A^2 \leq B^2]$ and

so this limit can be given by a Boolean expression in terms of the primitives of the formula. Again, by Lemma 7 if $h > 0$, then $\lim[A_n^{2^n} \leq P + B_n^{2^n}] = \lim[B_n^2 \leq 1] \cap \lim[[A_n^2 \leq 1] \cap [P \geq 0]] \cup \lim[B_n^2 \geq 1] \cap \lim[A_n^2 \leq B_n^2]$ is expressed in terms of limits of formulas of height smaller than h . Thus by induction hypothesis we conclude that $\lim[A^{2^n} \leq P + B^{2^n}]$ exists, has a semi-algebraic structure, and can be given in terms of a Boolean expression involving the primitives of the formula. \square

3. THE COMPLEX CASE

Consider now a formula of height $h \geq 1$ written in additive form: $Q_1 A_1^n + \dots + Q_l A_l^n - P$, where Q_1, \dots, Q_l , and P are complex polynomials in m variables and A_1, \dots, A_l are formulas of height $\leq h-1$. We wish to describe $\lim[Q_1 A_1^n + \dots + Q_l A_l^n = P]$. We start with the simplest situation, $\lim[A_1^n = P]$.

Lemma 10. *Let A, P be complex polynomials in m variables. Suppose that $P \neq 0$ and A and P are independent in the sense that $P \nmid dP \wedge dA$ in the region where $|A| < 1$. Then $\lim[A^n = P] = \partial(|A| < 1] \cup [P \neq 0]$.*

Proof. Let $z \in \lim[A^n = P]$. Then $z = \lim z_n$ and there is a sequence $k_n \rightarrow \infty$ such that $A(z_n)^{k_n} = P(z_n)$. There are two possibilities: $|A(z)| < 1$, then $|A(z_n)| < 1$ for large n and $P(z) = \lim P(z_n) = 0$, i.e., $z \in [|A| < 1, P = 0]$; and $|A(z)| = 1$, then $z \in [|A| = 1] = \overline{[|A| = 1] \cup [P \neq 0]}$. Since $\partial(|A| < 1] \cap [P \neq 0] = \overline{[|A| < 1]} \cap [P = 0] \cup [|A| = 1] \cap \overline{[P \neq 0]}$, we obtain that $z \in \partial(|A| < 1] \cap [P \neq 0]$. Conversely, we wish to prove that $\overline{[|A| < 1]} \cap [P = 0] \cup [|A| = 1] \cap \overline{[P \neq 0]} \subset \lim[A^n = P]$. Since $\lim[A^n = P]$ is closed, it is enough to show that $[|A| < 1] \cap [P = 0] \cup [|A| = 1] \cap [P \neq 0] \subset \lim[A^n = P]$. First take $z \in [|A| < 1] \cap [P = 0]$. Then $|A(z)| < 1$ and $P(z) = 0$. In the plane (A, P) the graph G_{k_n} of the map $P = A^{k_n}$ approaches any point $(A, 0)$ with $|A| < 1$ as $k_n \rightarrow \infty$. Thus given $\epsilon > 0$ there is N such that for each $n \geq N$ the point $(A(z), \xi_n) \in G_{k_n}$ satisfies $|\xi_n| < \epsilon$. Since $S := A^{-1}(A(z)) \cap P^{-1}(P(z))$ is an algebraic subvariety of codimension ≥ 2 , there is a 1-disc $z \in U_\epsilon \subset A^{-1}(A(z))$, in general position with S such that $P|_{U_\epsilon}$ is a covering map of U_ϵ over a neighborhood of $0 \in \mathbb{C}$. Thus for k_n large enough there is $w_n \in U_\epsilon$ such that $P(w_n) = \xi_n$. Since $A(w_n) = A(z)$ and $(A(z), \xi_n) \in G_{k_n}$ we obtain that $P(w_n) = A(w_n)^{k_n}$. Clearly $w_n \rightarrow z$ and so $z \in \lim[A^n = P]$.

Figure 2

Suppose now that $z \in [|A| = 1] \cap [P \neq 0]$. Then $|A(z)| = 1$ and $P(z) \neq 0$. In the plane (A, P) the horizontal line through the point $(0, P(z))$ intersects the graph G_{k_n} of the map $P = A^{k_n}$ in k_n points over the points $\mathfrak{A}_n = \{P(z)^{1/k_n}\}$ in the A -axis. For each of the points $w \in A^{-1}(\mathfrak{A}_n)$ we have $P(z) = P(w) = A(w)^{k_n}$. Since $|P(z)|^{1/k_n} \rightarrow 1$, the graph G_{k_n} approaches the set $|A| = 1$, thus the set \mathfrak{A}_n tends to fill the unitary circle. Therefore for each n we can find $w_n \in A^{-1}(\mathfrak{A}_n)$, $w_n \rightarrow z$, such that $P(w_n) = A(w_n)^{k_n}$. \square

Lemma 11. $\lim(|A|^n = |P|) = \partial(|A| < 1] \cap [|P| \neq 0]) = \lim[A^n = P]$.

Proof. Same as above. \square

Lemma 12. *Suppose P and Q are polynomials, not identically zero, and let A_n be a formula of positive height h . Assuming for n large that $P \nmid dP \wedge dA_n$ and $Q \nmid dQ \wedge dA_n$, we have*

$$\lim[QA_n = P] = \partial(\lim[|A_n| < 1] \cap [P \neq 0]) \cup \partial(\lim[|A_n| > 1] \cap [Q \neq 0])$$

Proof. Take $z \in \lim[QA_n^n = P]$. Then there are sequences $z_n \rightarrow z$ and $k_n \rightarrow \infty$ such that $Q(z_n)A_{k_n}(z_n)^{k_n} = P(z_n)$. We have the following possibilities:

- $\lim |A_{k_n}| < 1$. Then for n large $|A_{k_n}(z_n)| < 1$ and $P(z) = \lim P(z_n) = 0$. Thus $z \in \lim[|A_n| < 1] \cap [P = 0]$.
- $\lim |A_{k_n}(z_n)| = 1$. Then $z \in \lim[|A_n| = 1] = \lim[|A_n| = 1] \cap \overline{[P \neq 0]} = \lim[|A_n| = 1] \cap \overline{[Q \neq 0]}$
- $\lim |A_{k_n}(z_n)| > 1$. Then for n large $|A_{k_n}| > 1$ and $Q(z) = \lim Q(z_n) = \lim P(z_n)A_{k_n}^{-k_n} = 0$.

Reciprocally, if $z \in \lim[|A_n| < 1] \cap [P = 0]$ then there are sequences $z_n \rightarrow z$ and $k_n \rightarrow \infty$ such that $P(z) = 0$ and $\lim |A_{k_n}(z_n)| < 1$. Assume that $Q(z) \neq 0$. Let $\mathfrak{D} = \{w : |A_{k_n}(w)| \leq 1, n \geq 1\}$. Then $\mathfrak{D} \neq \emptyset$ and since (A_{k_n}) is bounded in \mathfrak{D} , it is a normal family. Then there is a subsequence, say (A_{k_n}) , which converges to a holomorphic function A , i.e., $\lim A_{k_n}(w) = A(w)$. Since $|A(z)| < 1$, we have $\xi_n = A(z)^{l_n} \rightarrow 0$ as $l_n \rightarrow \infty$. As by hypothesis $S_n := A_{k_n}^{-1}(A(z)) \cap (P/Q)^{-1}((P/Q)(z))$ is a codimension 2 algebraic subvariety for n large, then there is a neighborhood $z \in U$ such that $(P/Q)|_{U \cap A_{k_n}^{-1}(A(z))}$ projects onto a neighborhood of $0 \in \mathbb{C}$. Thus, there is $w_n \in U \cap A_{k_n}^{-1}(A(z))$, such that $(P/Q)(w_n) = \xi_n$. Therefore $P(w_n) = Q(w_n)\xi_n = Q(w_n)A(z)^{l_n} = Q(w_n)A_{k_n}(w_n)^{l_n}$. Clearly, $w_n \rightarrow z$ and so $z \in \lim[QA_n^n = P]$. Similarly, if $z \in \lim[|A_n| = 1]$ we have that $z \in \lim[QA_n^n]$. On the other hand, if $z \in \lim[|A_n| > 1] \cap [Q = 0]$ then $Q(z) = 0$ and $\lim |A_{k_n}(z_n)| > 1$. If $P = 0$ then $z \in \lim[QA_n^n = P]$. We assume $P(z) \neq 0$. Define the domain $\tilde{\mathfrak{D}} = \{w : |A_{k_n}(w)^{-1}| < 1, n > 1\}$. On $\tilde{\mathfrak{D}}$ the sequence $(A_{k_n}^{-1})$ is normal and converges to a holomorphic function B , i.e., $\lim A_{k_n}(w)^{-1} = B(w)^{-1}$. Thus $|B(z)| > 1$ and $\eta_n = B(z)^{-l_n} \rightarrow 0$ as $l_n \rightarrow \infty$. By hypothesis $A_{k_n}^{-1}(B(z)) \cap Q^{-1}(0)$ is a codimension 2 algebraic subvariety for n large. Then, since $P(z) \neq 0$, there is a neighborhood $z \in U$ such that $Q/P|_{U \cap A_{k_n}^{-1}(B(z))}$ projects over a neighborhood of $0 \in \mathbb{C}$. Thus there is $w_n \in A_{k_n}^{-1}(B(z)) \cap U$ such that $(Q/P)(w_n) = \eta_n = B(z)^{-l_n} = A_{k_n}(w_n)^{-l_n}$ or $Q(w_n)A_{k_n}(w_n)^{l_n} = P(w_n)$. Clearly, $w_n \rightarrow z$ and so $z \in \lim[QA_n^n = P]$. \square

Example. Let us compute $\lim[(A^n + P_1)^n = Q]$.

$$\begin{aligned} \lim[(A^n + P_1)^n = Q] &= \partial(\lim[|A^n + P_1| < 1] \cap [Q \neq 0]) \\ &= \overline{\lim[|A^n + P_1| < 1] \cap [Q = 0]} \cup (\lim[|A^n| = 1] \cap \overline{[Q \neq 0]}) \\ \lim[|A^n + P_1| < 1] &= \lim[|A|^n < 1 + |P_1|] \\ &= [|A|^2 < 1] \\ \lim[|A|^n = 1] &= [|A| = 1] \end{aligned}$$

Thus,

$$\begin{aligned} \lim[(A^n + P_1)^n = Q] &= [|A|^2 \leq 1] \cap [Q = 0] \cup \overline{[|A| = 1] \cap [Q \neq 0]} \\ &= [|A|^2 \leq 1] \cap [Q = 0] \cup [|A| = 1]. \end{aligned}$$

Figure 3

Thus this limit is the pull back by a rational map of a Reinhardt variety over a semi-algebraic subset of \mathbb{R}^2 . This example reflects pretty well the general picture described in Theorem 3.

Suppose that $Q_1, \dots, Q_l, P \in \mathbb{C}[x_1, \dots, x_m]$. In what follows we will write

$$\begin{aligned} \mathcal{Q}A^n &= [Q_1A_1^n + \dots + Q_lA_l^n = P] \\ \mathcal{Q}A^n(\hat{i}) &= [Q_1A_1^n + \dots + \widehat{Q_iA_i^n} + \dots + Q_lA_l^n = P] \end{aligned}$$

Lemma 13. *Suppose that $Q_1, \dots, Q_l, P \in \mathbb{C}[x_1, \dots, x_m]$ and assume that $Z_{Q_i}, Z_{Q_j}, Z_{Q_k}$ intersect in general position if $i \neq j \neq k \neq i$. Let A_1, \dots, A_l be formulas of positive height h . Then,*

- (1) $\lim[Q_1A_1^n + \dots + Q_lA_l^n = P] = \bigcup_{i,j=1}^l \lim[|A_i| = |A_j| > 1] \cup (Z_{Q_i} \cap Z_{Q_j}) \bigcup_{i=1}^l \lim[|A_i| < 1] \cap \lim[Q_1A_1^n + \dots + \widehat{Q_iA_i^n} + \dots + Q_lA_l^n = P]$
- (2) $\lim[Q_1A_1^n + \dots + Q_lA_l^n = P] = \bigcup_{i,j=1}^l \lim[|A_i| = |A_j| > 1] \cup (Z_{Q_i} \cap Z_{Q_j}) \cup \partial(\lim[|A_1| < 1] \cap \dots \cap \lim[|A_l| < 1] \cap [P \neq 0])$

Proof. Write $\mathcal{R}_1 := \bigcup_{i=1}^l \lim[|A_i| < 1]$, $\mathcal{R}_2 := \bigcap_{i=1}^l \lim[|A_i| \geq 1]$. Then Lemma 13 follows from the next two lemmas. \square

Lemma 14.

$$\begin{aligned} &\lim[Q_1A_1^n + \dots + Q_lA_l^n = P] \cap \mathcal{R}_1 \\ &= \bigcup_{i=1}^l \lim[|A_i| < 1] \cap \lim[Q_1A_1^n + \dots + \widehat{Q_iA_i^n} + \dots + Q_lA_l^n = P] \\ &= \partial(\lim[|A_1| < 1] \cap \dots \cap \lim[|A_l| < 1] \cap [P \neq 0]) \end{aligned}$$

Proof. Let $z \in \lim[Q_1A_1^n + \dots + Q_lA_l^n = P] \cap \mathcal{R}_1$. Then there are sequences $z_n \rightarrow z$, $k_n \rightarrow \infty$ such that $Q_1(z_n)A_{1k_n}(z_n)^{k_n} + \dots + Q_l(z_n)A_{lk_n}(z_n)^{k_n} = P(z_n)$. Suppose first that $\lim|A_{1k_n}(z_n)| < 1$. Then $\epsilon_n(z_n) = Q_1(z_n)A_{1k_n}(z_n)^{k_n} \rightarrow 0$ as $n \rightarrow \infty$ and if we define

$$f_n(z_n) := P(z_n) - Q_2(z_n)A_{2k_n}(z_n)^{k_n} - \dots - Q_l(z_n)A_{lk_n}(z_n)^{k_n}$$

we have $f_n(z_n) = \epsilon_n(z_n)$. Let $Z_n = f_n^{-1}(0)$ and $Z = \lim Z_n$. We claim that $z \in Z$. Indeed, if $z \notin Z$ then there are neighborhoods $z \in V$ and $Z \subset W$ with $V \cap W = \emptyset$. For n large $z_n \in V$ and $f_n^{-1}(\epsilon_n) \subset W$, a contradiction since $f_n(z_n) = \epsilon_n$ and $f_n(z_n) = \epsilon_n$ and $z_n \rightarrow z$. Then there is $w_n \in Z_n = f_n^{-1}(0)$, $w_n \rightarrow z$, i.e., $f_n(w_n) = 0$, $w_n \rightarrow z$. This means,

$$Q_2(w_n)A_{2k_n}(w_n)^{k_n} + \dots + Q_l(w_n)A_{lk_n}(w_n)^{k_n} = P(w_n)$$

or what is the same, $z \in \lim[|A_1| < 1] \cap \lim[Q_2A_2^n + \dots + Q_lA_l^n = P]$. Similarly, if $z \in \mathcal{R}_1$, then

$$z \in \bigcup_{i=1}^l \lim[|A_i| < 1] \cap \lim[Q_1A_1^n + \dots + \widehat{Q_iA_i^n} + \dots + Q_lA_l^n = P]$$

Conversely, suppose there are sequences $z_n \rightarrow z$ and $w_n \rightarrow z$ such that $f_n(w_n) = 0$ and $\epsilon_n(z_n) \rightarrow 0$, then denoting again by $Z_n = f_n^{-1}(0)$ then $Z_n \rightarrow Z$ and since $w_n \in Z_n$ and $w_n \rightarrow z$ then $z \in Z$. Therefore for any δ small positive there is $y_n \in \epsilon_n^{-1}(\delta) \cap f_n^{-1}(\delta) \neq \emptyset$, i.e., $\epsilon_n(y_n) = f_n(y_n)$. We will show now that for the

points in the region \mathcal{R}_1 we have,

$$\begin{aligned} & \partial([|A_1| < 1] \cap \cdots \cap [|A_l| < 1] \cap [P \neq 0]) = \\ &= \bigcup_{i=1}^l [|A_i| < 1] \cap \lim[Q_1 A_1^n + \cdots + \widehat{Q_i A_i^n} + \cdots + Q_l A_l^n = P] \quad (**) \\ &= \lim[Q_1 A_1^n + \cdots + Q_l A_l^n = P]. \end{aligned}$$

We proceed by induction on l . For $l = 2$ we have by Lemma 12,

$$\begin{aligned} & \partial([|A_1| < 1] \cap [|A_2| < 1] \cap [P \neq 0]) = \\ &= [|A_1| < 1] \cap \partial([|A_2| < 1] \cap [P \neq 0]) \cup [|A_2| < 1] \cap \partial([|A_1| < 1] \cap [P \neq 0]) \\ &= [|A_1| < 1] \cap \lim[Q_2 A_2^n = P] \cup [|A_2| < 1] \cap \lim[Q_1 A_1^n = P] \\ &= \lim[Q_1 A_1^n + Q_2 A_2^n = P]. \end{aligned}$$

For $l > 2$, we have

$$\begin{aligned} & \partial([|A_1| < 1]) \cap \cdots \cap (|A_l| < 1) \cap [P \neq 0] = \\ &= \bigcup_{i=1}^l (|A_i| < 1) \cap \partial([|A_1| < 1] \cap \cdots \cap \widehat{(|A_i|)} \cap \cdots \cap (|A_l| < 1) \cap [P \neq 0]) \\ &= \bigcup_{i=1}^l (|A_i| < 1) \cap \lim[Q_1 A_1^n + \cdots + \widehat{Q_i A_i^n} + \cdots + Q_l A_l^n = P] \\ &= \lim[Q_1 A_1^n + \cdots + Q_l A_l^n = P] = \lim[Q_1 A_1^n + \cdots + Q_l A_l^n = P], \end{aligned}$$

where the last two equalities are derived by induction hypothesis on (**). \square

Lemma 15.

$$\lim[Q_1 A_1^n + \cdots + Q_l A_l^n = P] \cap \mathcal{R}_2 = \bigcup_{i,j=1}^l \lim[|A_i| = |A_j| > 1] \cup (Z_{Q_i} \cap Z_{Q_j})$$

Proof. Suppose now that

$$z \in \lim[|A_1| > 1] \cap \cdots \cap \lim[|A_q| > 1] \cup \lim[|A_{q+1}| = 1] \cap \cdots \cap \lim[|A_l| = 1]$$

Then, $q \neq 1$ and,

$$Q_1(z_n)(A_1)_{k_n}(z_n)^{k_n} + \cdots + Q_q(z_n)(A_q)_{k_n}(z_n)^{k_n} = R(z_n),$$

where

$$R(z_n) := P(z_n) - Q_{q+1}(z_n)(A_{q+1})_{k_n}(z_n)^{k_n} - \cdots - Q_l(z_n)(A_l)_{k_n}(z_n)^{k_n}$$

is locally bounded at z . For any $i, j = 1, \dots, q$, $i \neq j$, we can write the next inequality where, for simplicity, we wrote $i = 1$ and $j = 2$:

$$|Q_1(z_n)| |(A_1)_{k_n}(z_n)|^{k_n} - |Q_2(z_n)| |(A_2)_{k_n}(z_n)|^{k_n} \leq \sum_{t=3}^q |Q_t(z_n)| |(A_t)_{k_n}(z_n)|^{k_n} + |R(z_n)|$$

Thus, dividing both members of this expression by $\prod_{t=3}^q |A_t(z_n)|^{k_n}$ we obtain a left member locally bounded at z . Then there is a bounded sequence $\{\lambda_n\}$ such that

$$|Q_1(z_n)| |A_1(z_n)|^{k_n} / \prod_{t=3}^q |A_t(z_n)|^{k_n} = \lambda_n |Q_2(z_n)| |A_2(z_n)|^{k_n} / \prod_{t=3}^q |A_t(z_n)|^{k_n},$$

i.e.,

$$|Q_1(z_n)||A_1(z_n)|^{k_n} = \lambda_n |Q_2(z_n)||A_2(z_n)|^{k_n}$$

Thus, either $z \in Z_{Q_1} \cap Z_{Q_2}$, or $|A_1(z_n)| = (\lambda_n |Q_2(z_n)| / |Q_1(z_n)|)^{1/k_n} |A_2(z_n)|$. Therefore, $\lim[|A_{1k_n}(z_n)|] = \lim[|A_{2k_n}(z_n)|]$. Thus, $z \in \bigcup_{i,j=1}^q [\lim[|A_i|] = \lim[|A_j|]] \cup (Z_{Q_i} \cap Z_{Q_j})$. This shows that for $l > 1$,

$$\lim[Q_1 A_1^n + \cdots + Q_l A_l^n = P] \cap \mathcal{R}_2 \subset [\lim[|A_i|] = \lim[|A_j|]] \cup (Z_{Q_i} \cap Z_{Q_j}) \quad (*)$$

We proceed to show now the converse to (*). Suppose $z \in [|A_i| = |A_j| > 1] \cup (Z_{Q_i} \cap Z_{Q_j})$. For simplicity take $i = 1, j = 2$ and $z \in [|A_1| = |A_2| > 1]$, $|A_i(z)| > 1, i = 1, \dots, k, |A_j(z)| \leq 1, j = k+1, \dots, l$. Consider the expression

$$a_n := Q_1(A_1/A_3 \dots A_k)^n + Q_2(A_2/A_3 \dots A_k)^n$$

We claim that the curve $a_n = 0$ approaches z as $n \rightarrow \infty$. Indeed, from $a_n(w) = 0$ we obtain

$$(A_1/A_2)^n(w) = -(Q_2/Q_1)(w).$$

For any w close to z such that $\arg(-Q_2/Q_1)(w)$ is irrational we have that $(-Q_2/Q_1)(w)^{1/n}$ approaches the circle of center $0 \in \mathbb{C}$ and radius one as $n \rightarrow \infty$. Therefore $(A_1/A_2)(z)$ is in the closure of the sequence $((-Q_2/Q_1(w))^{1/n})_n$. On the other hand, if

$$b_n := P/(A_3 \cdots A_k)^n - 1/(A_3 \cdots A_k)^n \sum_{j=3}^l Q_j A_j^n$$

then $b_n(z) \rightarrow 0$ as $n \rightarrow \infty$. Therefore the curve $b_n = 0$ approaches z as $n \rightarrow \infty$. Thus there is $z_n \in [a_n = b_n], z_n \rightarrow z$, i.e.,

$$(Q_1 A_1^n / (A_3 \cdots A_k)^n + Q_2 A_2^n / (A_3 \cdots A_k)^n)(z_n) = 1 / (A_3 \cdots A_k)^n (P - \sum_{j=3}^l Q_j A_j^n)(z_n)$$

$$\text{or } Q_1(z_n)A_1(z_n)^n + \cdots + Q_l(z_n)A_l(z_n)^n = P(z_n). \quad \square$$

Lemma 16. *Suppose that $Q_1 A_1^n + \cdots + Q_l A_l^n$ is a formula of positive height h . Then,*

$$\begin{aligned} & \lim[|Q_1 A_1^n + \cdots + Q_l A_l^n| < 1] \\ &= \bigcup_{i=1}^l \lim[|A_i| < 1] \cap \lim[|Q_1 A_1^n + \cdots + \widehat{Q_i A_i^n} + \cdots + Q_l A_l^n| < 1] \bigcup_{i,j=1, i \neq j}^l [\lim |A_i| ???] \\ &= \lim |A_j| > 1 \cup (Z_{Q_i} \cap Z_{Q_j}). \end{aligned}$$

Proof. Let $z \in \lim[|Q_1 A_1^n + \cdots + Q_l A_l^n| \leq 1]$. There is $z_n \rightarrow z$ and $k_n \rightarrow \infty$ such that $|Q_1(z_n)A_1(z_n)^{k_n} + \cdots + Q_l(z_n)A_l(z_n)^{k_n}| < 1$. Suppose that $\lim |A_{1k_n}(z_n)| < 1$, then $\epsilon_n = |Q_1 A_1(z_n)^{k_n}| \rightarrow 0$ and there is $w_n \rightarrow z$ such that $|Q_2(w_n)A_2(w_n)^{k_n} + \cdots + Q_l(w_n)A_l(w_n)^{k_n}| < 1$. Therefore, $z \in \bigcup_{i=1}^l \lim[|A_i| < 1]$, then $z \in \bigcup_{i=1}^l \lim[|A_i|] \cap \lim[|Q_1 A_1^n + \cdots + \widehat{Q_i A_i^n} + \cdots + Q_l A_l^n| < 1]$. On the other hand, if $z \in \bigcap_{i=1}^q \lim[|A_i| > 1] \cap \bigcap_{j=q+1}^l \lim[|A_j|]$, then $Q_1(z_n)A_{1k_n}(z_n)^{k_n} + \cdots + Q_q(z_n)A_{qk_n}(z_n)^{k_n} \leq 1 + S(z_n)$, where $S(z_n) = |\sum_{j=q+1}^l Q_j(z_n)A_j(z_n)^{k_n}|$, is locally bounded at z . Proceeding as in Lemma 13 we obtain that for any $i, j = 1, \dots, q$ either $z \in (\lim |A_i| = \lim |A_j|)$ or $z \in Z_{Q_i} \cap Z_{Q_j}$. The proof of the converse follows the same line of arguments of Lemma 13 \square

Proof of Theorem 2. In order to describe $\lim[Q_1A_1^n + \cdots + Q_lA_l^n = P]$ we first use induction on l by means of Lemma 13 which reduces the problem to describe $\lim[QA^n = P]$ and $\lim[|A| < 1]$ where $QA^n - P$ has height $h \geq 1$. Then we proceed by induction on h . For $h = 1$ Lemma 12 gives $\lim[QA^n = P] = \partial(\lim[|A_n| < 1] \cap (P \neq 0)) \cup \partial(\lim[|A_n| > 1] \cap [Q \neq 0])$ which reduces the problem to height $h - 1$. It only remains to find $\lim[|A_n| < 1]$ and this follows from Lemma 14.

Thus we have shown that this limit can be expressed by algebraic relations between $|A_1|, \dots, |A_l|$ and $|P|$. \square

4. ALGEBRAIC CURVES AS INTEGRALS OF DIFFERENTIAL EQUATIONS

Lemma 17. *Given polynomials A and P , there is a family (\mathcal{X}_n) of polynomial vector fields of fixed degree such that $[A^{2n} = P]$ is an integral curve of \mathcal{X}_n .*

Proof. Let \mathcal{X}_n be the field corresponding to the following differential equation:

$$\dot{x} = -2nPA_y + P_yA, \quad \dot{y} = 2nPA_x - P_xA.$$

Let $f = A^{2n} - P$. Then

$$\dot{x}f_x + \dot{y}f_y = 2n(P_yA_x - P_xA_y)f,$$

as can be easily verified. This shows that $[f = 0]$ is an integral curve of \mathcal{X}_n . \square

Thus, we have curves of increasing degree that are integral curves of polynomial fields of fixed degree. The next lemma says that in this case the field is essentially unique. The following proof is essentially due to B. Scárdua.

Lemma 18. *Suppose that $[f_n = 0]$ is a family of polynomial curves indexed by their degree. Assume that each curve is an integral curve of two differential equations of bounded degree: $\omega_n = 0$ and $\Omega_n = 0$. Then, for n large enough, $\omega_n = 0$ and $\Omega_n = 0$ define the same foliation.*

Proof. Forget the indices, for simplicity.

The hypotheses imply that

$$\begin{aligned} df \wedge \omega &= f \ell dx \wedge dy \\ df \wedge \Omega &= f L dx \wedge dy, \end{aligned}$$

where ℓ and L are polynomials.

Assume that $\omega \wedge \Omega \neq 0$.

If $df \wedge \Omega \neq 0$, then we can write

$$\omega = \alpha df + \beta \Omega.$$

The coefficients α and β are determined as follows:

$$\begin{aligned} \omega \wedge \Omega = \alpha df \wedge \Omega &\Rightarrow \alpha = \frac{\omega \wedge \Omega}{df \wedge \Omega} \\ df \wedge \omega = \beta df \wedge \Omega &\Rightarrow \beta = \frac{df \wedge \omega}{df \wedge \Omega}. \end{aligned}$$

Therefore

$$\beta = \frac{\ell}{L}, \quad \alpha = \frac{\omega \wedge \Omega}{f L dx \wedge dy}$$

and so

$$\omega = \frac{\omega \wedge \Omega}{dx \wedge dy} \cdot \frac{df}{fL} + \frac{\ell}{L} \Omega,$$

or

$$L\omega = \frac{\omega \wedge \Omega}{dx \wedge dy} \cdot \frac{df}{f} + \ell\Omega.$$

Assume that f is irreducible. Since $L\omega - \ell\Omega$ has bounded degree, we must have that $f dx \wedge dy$ divides $\omega \wedge \Omega$, that is,

$$\omega \wedge \Omega = f\mu dx \wedge dy,$$

for some polynomial μ . Hence, $L\omega = \mu df + \ell\Omega$.

Now $\partial\ell = \partial\omega - 1$ and $\partial L = \partial\Omega - 1$, and so μdf has bounded degree. Since $df_n \rightarrow \infty$ we conclude that $\mu_n = 0$ for large n .

If $df \wedge \Omega = 0$, then we take $df \wedge \omega \neq 0$. If both expressions vanish identically, then ω , Ω , and df define the same foliation. \square

Moreover, as the next lemma indicates, formulas that are more complicated than $A^{2n} = P$ are not likely to be integral curves of fields of fixed degree.

Lemma 19. *Let A , B , and P be bivariate polynomials such that $A(0,0) = 0 = B(0,0)$ and $(A,B) = 1$. Then, the curves in the family $A^n + B^n = P$ are not integral curves of a family of polynomial fields of degree 2.*

Proof. Suppose that A and B have degree k and P has degree j . Let $f = A^n + B^n - P$. Suppose that f is an integral curve of the 1-form

$$\omega = adx + bdy,$$

with a and b polynomials of degree 2. Then,

$$df \wedge \omega = fLdx \wedge dy,$$

with L a polynomial of degree 1. This equation is equivalent to

$$(nA^{n-1}A_x + nB^{n-1}B_x - P_x)b - (nA^{n-1}A_y + nB^{n-1}B_y - P_y)a = (A^n + B^n - P)L.$$

For n large, because $A(0,0) = 0 = B(0,0)$, we obtain

$$\begin{aligned} (1) \quad & P_x b - P_y a = PL \\ (2) \quad & nA^{n-1}(A_x b - A_y a) + nB^{n-1}(B_x b - B_y a) = (A^n + B^n)L. \end{aligned}$$

Because $(A,B) = 1$, we get

$$\begin{aligned} (3) \quad & n(A_x b - A_y a) = AL \\ & n(B_x b - B_y a) = BL \end{aligned}$$

(The proof is at the end.)

Suppose that P is homogeneous of degree j , A and B are homogeneous of degree k , and a and b are homogeneous of degree 2, in equations (1) and (3). This is not a restriction because it suffices to compare the homogeneous parts of highest degree in these equations.

Equation (3) can be written as

$$n \begin{pmatrix} -A_y & A_x \\ -B_y & B_x \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = L \begin{pmatrix} A \\ B \end{pmatrix}$$

If $\Delta = A_x B_y - A_y B_x$, then

$$n \begin{pmatrix} a \\ b \end{pmatrix} = \frac{L}{\Delta} \begin{pmatrix} B_x & -A_x \\ B_y & -A_y \end{pmatrix} \begin{pmatrix} A \\ B \end{pmatrix}$$

Since, by Euler's formula, $kA = A_x x + A_y y$ and $kB = B_x x + B_y y$, we get

$$\begin{aligned} n \begin{pmatrix} a \\ b \end{pmatrix} &= \frac{L}{k\Delta} \begin{pmatrix} B_x & -A_x \\ B_y & -A_y \end{pmatrix} \begin{pmatrix} A_x & A_y \\ B_x & B_y \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} \\ &= \frac{L}{k\Delta} \begin{pmatrix} 0 & -\Delta \\ \Delta & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} \\ &= \frac{L}{k} \begin{pmatrix} -y \\ x \end{pmatrix}, \end{aligned}$$

which implies that

$$a = -\frac{L}{kn}y, \quad b = \frac{L}{kn}x.$$

From (1), we get

$$P_x \frac{L}{kn}x + P_y \frac{L}{kn}y = PL,$$

that is,

$$P_x x + P_y y = nkP,$$

which implies that P is homogeneous of degree nk . Since n is arbitrarily large and P has a fixed degree, this is cannot happen. Therefore, f is not an integral curve of ω .

We still have to prove that (2) implies (3). In fact, let $\alpha = A_x b - A_y a$ and $\beta = B_x b - B_y a$. Then

$$nA^{n-1}\alpha + nB^{n-1}\beta = L(A^n + B^n),$$

that is,

$$A^{n-1}(n\alpha - LA) = B^{n-1}(-n\beta + LB).$$

Since $(A, B) = 1$, this implies that $A^{n-1} | (-n\beta + LB)$ and $B^{n-1} | (n\alpha - LA)$. Hence, there is a polynomial λ such that

$$\begin{aligned} \lambda A^{n-1} &= -n\beta + LB \\ \lambda B^{n-1} &= (n\alpha - LA) \end{aligned}$$

Comparing degrees, we get $\lambda = 0$ for large n . Therefore,

$$(-n\beta + LB) = 0 = (n\alpha - LA),$$

as claimed. □

Remark

Define the *length* of a formula as the minimum number of its primitives of degree ≥ 1 . So, for instance, the formula

$$(x+1)^{2n} + ((x-y-1)^n + y)^n + y^2 - 1$$

has length 4.

Suppose that \mathcal{C} is a family of curves given by the zeros of a formula of positive height. Let l be the length of the formula and assume that the curves defined by the zeros of its primitives intersect transversely in the complex domain. If \mathcal{V} is a family of vector fields of degree k such that the elements of \mathcal{C} are integral curves of the corresponding elements of \mathcal{V} , then $l \leq k^2 + k + 1$, as this last expression is the number of singular points of the elements of \mathcal{V} . In particular if $l > k^2 + k + 1$ the

elements in \mathcal{C} can not be integral curves of a family of polynomial vector fields of degree $\leq k$.

Theorem 4. *Every generic basic closed one-dimensional semi-algebraic set in the plane is the limit of an family of algebraic curves that are integral curves of a family of polynomial vector fields of fixed degree.*

Proof. Let Ω be a generic basic closed semi-algebraic set. It is known (but hard to prove) that every basic open semi-algebraic set in the plane can actually be given by *two* inequalities [?]. Since Ω is generic, this also applies to Ω and we can write $\Omega = [P \geq 0, Q \geq 0]$. We shall show that $\Omega = \lim[A_n^{2n} \leq P]$ for

$$A_n = \frac{Q}{n} - 1.$$

Indeed,

$$[A_n^2 \leq 1] = [(\frac{Q}{n} - 1)^2 \leq 1] = [0 \leq Q \leq 2n]$$

Hence,

$$\{z : A_n^2(z) \leq 1, \text{ for sufficiently large } n\} = [Q \geq 0].$$

and so

$$\{z : P(z) \geq 0, A_n^2(z) \leq 1, \text{ for sufficiently large } n\} = [P \geq 0, Q \geq 0].$$

Lemma ?? then says that

$$\lim[A_n^{2n} \leq P] = [P \geq 0, Q \geq 0] = \Omega,$$

if $[A_n^2 \leq 1, P \geq 0]$ is generic for sufficiently large n . (***)

As mentioned in Section 4, the curves $[A_n^{2n} = P]$ are integral curves of a family of polynomial vector fields of fixed degree. (Note that, although A_n has coefficients that depend on n , the vector fields are still of fixed degree.) \square

Acknowledgements. Figure 1 was obtained with *Grafeq*, which was instrumental in our research. The authors are partially supported by research grants from CNPq.

REFERENCES

- [1] R. Benedetti and J.-J. Risler. *Real algebraic and semi-algebraic sets*. Actualités Mathématiques. [Current Mathematical Topics]. Hermann, Paris, 1990.
- [2] E. Bishop. Conditions for the analyticity of certain sets. *Michigan Math. J.*, 11:289–304, 1964.
- [3] J. V. Pereira. On the Poincaré problem for foliations of general type. *Math. Ann.*, 323(2):217–226, 2002.
- [4] J.-J. Risler. Additive complexity and zeros of real polynomials. *SIAM J. Comput.*, 14(1):178–183, 1985.
- [5] G. Stolzenberg. *Volumes, limits, and extensions of analytic varieties*. Lecture Notes in Mathematics, No. 19. Springer-Verlag, Berlin, 1966.

IMPA—INSTITUTO NACIONAL DE MATEMÁTICA PURA E APLICADA, ESTRADA DONA CASTORINA 110, 22460-320 RIO DE JANEIRO, RJ, BRAZIL
E-mail address: camacho@impa.br, lhf@impa.br