

A New Tower Over Cubic Finite Fields

Alp Bassa, Arnaldo Garcia and Henning Stichtenoth

We present an explicit new tower of function fields $(F_n)_{n \geq 0}$ over the finite field with $\ell = q^3$ elements, where the limit of the ratios (number of rational places of F_n)/(genus of F_n) is bigger or equal to $2(q^2 - 1)/(q + 2)$. This tower contains as a subtower the tower which was introduced by Bezerra–Garcia–Stichtenoth (see [3]) and in the particular case $q = 2$ it coincides with the tower of van der Geer–van der Vlugt (see [12]). Many features of the new tower are very similar to those of the optimal wild tower in [8] over the quadratic field \mathbb{F}_{q^2} (whose modularity was shown in [6] by Elkies).

1 Introduction

Let F/\mathbb{F}_ℓ be an algebraic function field of one variable whose full constant field is the finite field \mathbb{F}_ℓ of cardinality ℓ . We denote by $g(F)$ the genus and by $N(F)$ the number of rational places (i.e., places of degree one) of F/\mathbb{F}_ℓ . The classical Hasse–Weil Theorem states that $N(F) \leq \ell + 1 + 2g(F)\sqrt{\ell}$.

Ihara [13] was the first to observe that this inequality can be improved substantially if the genus of F is large with respect to ℓ . He introduced the real number

$$A(\ell) := \limsup_{g(F) \rightarrow \infty} \frac{N(F)}{g(F)},$$

where F runs over all function fields over \mathbb{F}_ℓ . This number $A(\ell)$ is of fundamental importance to the theory of function fields over a finite field, since it gives information about how many rational places a function field F/\mathbb{F}_ℓ of large genus can have. While the Hasse–Weil Theorem gives that $A(\ell) \leq 2\sqrt{\ell}$, Ihara showed $A(\ell) \leq \sqrt{2\ell}$ for any ℓ and that $A(\ell) \geq \sqrt{\ell} - 1$ for ℓ a square. Later Drinfeld and Vladut [4] showed that

$$A(\ell) \leq \sqrt{\ell} - 1 \quad \text{for any } \ell. \tag{1}$$

Hence we have the equality (see also [5], [7], [16]) $A(\ell) = \sqrt{\ell} - 1$ for ℓ a square.

Much less is known if ℓ is not a square. One knows that for any ℓ (see [14])

$$A(\ell) \geq c \cdot \log \ell > 0, \quad \text{for some constant } c.$$

For $\ell = p^3$ (p a prime number), the best known lower bound for $A(\ell)$ is due to Zink [17]:

$$A(p^3) \geq \frac{2(p^2 - 1)}{p + 2}. \quad (2)$$

Zink obtained this result using degenerations of Shimura modular surfaces. Zink's bound was generalized by Bezerra, Garcia and Stichtenoth [3] who showed that

$$A(q^3) \geq \frac{2(q^2 - 1)}{q + 2} \quad (3)$$

holds for all prime powers q . For more information and references concerning Ihara's quantity $A(\ell)$ we refer to the recent survey article [11].

In order to obtain lower bounds for $A(\ell)$, it is natural to study towers of function fields; i.e., one considers sequences $\mathcal{G} = (G_0, G_1, G_2, \dots)$ of function fields G_i over \mathbb{F}_ℓ with $G_0 \subseteq G_1 \subseteq G_2 \subseteq \dots$ such that $g(G_i) \rightarrow \infty$. It is easy to see that the limit

$$\lambda(\mathcal{G}) := \lim_{i \rightarrow \infty} \frac{N(G_i)}{g(G_i)}$$

always exists (see [8]) and it is clear that $0 \leq \lambda(\mathcal{G}) \leq A(\ell)$.

A particularly interesting example of a tower $\mathcal{H} = (H_0, H_1, H_2, \dots)$ over the field \mathbb{F}_ℓ with $\ell = q^2$ is defined recursively as follows (see [8]): $H_0 = \mathbb{F}_\ell(u_0)$ is the rational function field, and for all $i \geq 0$ one considers the field $H_{i+1} = H_i(u_{i+1})$ with

$$u_{i+1}^q + u_{i+1} = \frac{u_i^q}{u_i^{q-1} + 1}. \quad (4)$$

This tower over \mathbb{F}_{q^2} has the limit $\lambda(\mathcal{H}) = q - 1 = \sqrt{\ell} - 1$ and therefore attains the Drinfel'd-Vlăduț bound (1). Elkies [6] has shown that \mathcal{H} is in fact a modular tower.

In [3] the following tower $\mathcal{E} = (E_0, E_1, E_2, \dots)$ over a cubic field \mathbb{F}_ℓ with $\ell = q^3$ is considered: again $E_0 = \mathbb{F}_\ell(v_0)$ is the rational function field, and for $i \geq 0$ one considers the field $E_{i+1} = E_i(v_{i+1})$ with

$$\frac{1 - v_{i+1}}{v_{i+1}^q} = \frac{v_i^q + v_i - 1}{v_i}. \quad (5)$$

The limit $\lambda(\mathcal{E})$ satisfies the inequality (thus proving Inequality (3)):

$$\lambda(\mathcal{E}) \geq \frac{2(q^2 - 1)}{q + 2}. \quad (6)$$

The tower \mathcal{H} over the quadratic field \mathbb{F}_ℓ with $\ell = q^2$ which is defined by Eqn. (4) has some nice features which allow a rather simple proof of the equality $\lambda(\mathcal{H}) = q - 1$, see [9]. The most important one is that all extensions H_{i+1}/H_i are Galois of degree q , and for all places $Q|P$ with ramification index $e = e(Q|P) > 1$ in H_{i+1}/H_i , the different exponent is $d(Q|P) = 2(e - 1)$.

In contrast, the tower \mathcal{E} over the cubic field \mathbb{F}_ℓ with $\ell = q^3$ which is defined by Eqn. (5) is much more complicated. Here (for $q \neq 2$) the extensions E_{i+1}/E_i are not even Galois, and there occurs tame and also wild ramification in E_{i+1}/E_i . The determination of the genus of E_n in [3] requires long and rather technical calculations. In [1] these calculations were replaced by a structural argument, thus obtaining a simpler proof of Inequality (6) without the explicit determination of $g(E_n)$.

In this paper we present a new tower \mathcal{F} over the cubic field \mathbb{F}_ℓ with $\ell = q^3$, whose limit also satisfies the inequality $\lambda(\mathcal{F}) \geq 2(q^2 - 1)/(q + 2)$ and which has nicer properties than the tower given by the recursion in Eqn. (5). This new tower $\mathcal{F} = (F_0, F_1, F_2, \dots)$ over \mathbb{F}_ℓ is defined as follows: $F_0 = \mathbb{F}_\ell(x_0)$ is the rational function field over \mathbb{F}_ℓ , and for $n \geq 0$ one sets $F_{n+1} = F_n(x_{n+1})$ with

$$(x_{n+1}^q - x_{n+1})^{q-1} + 1 = \frac{-x_n^{q(q-1)}}{(x_n^{q-1} - 1)^{q-1}}. \quad (7)$$

Our proof that the limit of this new tower satisfies the inequality $\lambda(\mathcal{F}) \geq 2(q^2 - 1)/(q + 2)$ is much easier, shorter and less computational than the proof in [3] for the tower \mathcal{E} .

Moreover, since we show that \mathcal{E} is a subtower of \mathcal{F} we get also a proof here of Inequality (6); in fact, it follows from [8] that $\lambda(\mathcal{E}) \geq \lambda(\mathcal{F})$ when \mathcal{E} is a subtower of \mathcal{F} .

Another remark is that while for the two towers over \mathbb{F}_{q^2} presented in [7] and [8] the subtower (i.e., the tower \mathcal{H} in [8]) was easier to handle (see [9]), for the two towers \mathcal{E} and \mathcal{F} over \mathbb{F}_{q^3} the supertower (i.e., the tower \mathcal{F}) turns out to be much easier to handle.

Finally we note that the tower \mathcal{F} coincides with the tower in [12] when $q = 2$ and also that the towers \mathcal{F} and \mathcal{H} have surprising similarities (see Section 8).

This paper is organized as follows: In Sec. 2 we introduce the sequence of function fields F_0, F_1, F_2, \dots over a field $K \supseteq \mathbb{F}_q$ recursively given by Eqn. (7) and we show in Theorem 2.2 that they define a tower \mathcal{F} over K (i.e., $F_0 \subsetneq F_1 \subsetneq F_2 \subsetneq \dots$, and K is the full constant field of all fields F_n). We note that Lemma 2.7 gives a remarkable property of the recursion in Eqn. (7). In Sec. 3 it is shown that for $K = \mathbb{F}_{q^3}$ there exist $q^3 - q$ rational places of F_0 which split completely in all extensions F_n/F_0 , thus providing many rational places of the function fields F_n/\mathbb{F}_{q^3} . In Sec. 4 and Sec. 5 we study ramification in the first steps $F_0 \subseteq F_1 \subseteq F_2$ of the tower. We note that the methods in Sec. 4 and Sec. 5 involve just simple calculations about ramification in certain Galois extensions $k(x)/k(w)$ of rational function fields. Section 6 is the core of this paper. The information from Sec. 4 and Sec. 5 is used in Sec. 6 to give an upper bound for the genus of the n -th function field F_n of the tower (see Theorem 6.5). The main tool here is Abhyankar's Lemma and a version of it (see Lemma 6.2) dealing with ramification in composites of certain wildly ramified extensions. Putting together these results we obtain in Sec. 7 the inequality $\lambda(\mathcal{F}) \geq 2(q^2 - 1)/(q + 2)$ for $K = \mathbb{F}_{q^3}$, which is the main result of the paper. Finally, in Sec. 8 we point out some surprising analogies between the tower \mathcal{F} over \mathbb{F}_{q^3} and the tower \mathcal{H} over \mathbb{F}_{q^2} which is defined by Eqn. (4). We also show that the above-mentioned tower \mathcal{E} is a subtower of \mathcal{F} .

NOTATIONS: We consider function fields F/K where K is the full constant field of F . In most cases K will be a finite field or the algebraic closure $\overline{\mathbb{F}}_q$ of a finite field.

We denote by $\mathbb{P}(F)$ the set of places of F/K . For $P \in \mathbb{P}(F)$, we will denote by v_P the corresponding discrete valuation of F/K and by \mathcal{O}_P the valuation ring of P . For $z \in \mathcal{O}_P$ we denote by $z(P)$ the residue class of z in \mathcal{O}_P/P . We denote by $\deg(P)$ the degree of P . In particular, if P is a place of degree one, then $z(P) \in K$.

For a finite separable extension E of F and a place $Q \in \mathbb{P}(E)$ we will denote by $Q|_F$ the restriction of Q to F . We write $Q|P$ if the place $Q \in \mathbb{P}(E)$ lies over the place $P \in \mathbb{P}(F)$. In this situation, we denote by $e(Q|P)$ and $d(Q|P)$ the ramification index and the different exponent of $Q|P$, respectively. The place $P \in \mathbb{P}(F)$ is said to be totally ramified in E/F if there is a place $Q \in \mathbb{P}(E)$ above P with $e(Q|P) = [E : F]$. It is said to be completely splitting in E/F if there are $n = [E : F]$ distinct places of E above P .

Let E/F be a Galois extension of function fields, let $P \in \mathbb{P}(F)$ and $Q \in \mathbb{P}(E)$ above the place P . We say that $Q|P$ is *weakly ramified* if the second ramification group $G_2(Q|P) = 1$; in other words, if $e(Q|P) = e_0 \cdot e_1$ where $(e_0, p) = 1$ and $e_1 = p^j$ is a power of the characteristic p of F , then $d(Q|P) = (e_0 e_1 - 1) + (e_1 - 1)$.

If $F = K(x)$ is a rational function field, we will write $(x = \alpha)$ for the place of F which is the zero of $x - \alpha$ (where $\alpha \in K$), and $(x = \infty)$ for the pole of x in $K(x)/K$.

2 The tower

Let K be a field of characteristic $p > 0$, let q be a power of p and assume that $\mathbb{F}_q \subseteq K$. We study the sequence $\mathcal{F} = (F_0, F_1, F_2, \dots)$ of function fields F_i/K which is defined recursively as follows: $F_0 = K(x_0)$ is the rational function field, and for $n \geq 0$ let $F_{n+1} = F_n(x_{n+1})$ where x_{n+1} satisfies the equation over F_n below:

$$(x_{n+1}^q - x_{n+1})^{q-1} + 1 = \frac{-x_n^{q(q-1)}}{(x_n^{q-1} - 1)^{q-1}}. \quad (8)$$

Remark 2.1. We set

$$f(T) := (T^q - T)^{q-1} + 1 \in K[T]. \quad (9)$$

Then Eqn. (8) can be written as

$$f(x_{n+1}) = \frac{1}{1 - f(1/x_n)}. \quad (10)$$

We also remark that $f(T) = (T^{q^2} - T)/(T^q - T)$, hence the roots of $f(T)$ are exactly the elements $\beta \in \mathbb{F}_{q^2} \setminus \mathbb{F}_q$. This property of the polynomial $f(T)$ will play an important role in Sections 3 and 4.

Theorem 2.2. *Let \mathcal{F} be the sequence of function fields F_n over K which is defined by Eqn. (8). Then \mathcal{F} is a tower over K and more precisely the following hold:*

- (i) *The extensions F_{n+1}/F_n are Galois for all $n \geq 0$.*
- (ii) *$[F_1 : F_0] = q(q-1)$ and $[F_{n+1} : F_n] = q$ for all $n \geq 1$.*
- (iii) *K is the full constant field of F_n , for all $n \geq 0$.*

The proof of Thm. 2.2 is given in several steps.

Lemma 2.3. F_{n+1}/F_n is Galois and $[F_{n+1} : F_n]$ divides $q(q-1)$, for all $n \geq 0$.

Proof. We set

$$u_n := \frac{-x_n^{q(q-1)}}{(x_n^{q-1} - 1)^{q-1}}. \quad (11)$$

Then x_{n+1} is a root of the polynomial $f_n(T) := (T^q - T)^{q-1} + 1 - u_n \in F_n[T]$. The other roots of $f_n(T)$ are the elements $ax_{n+1} + b$ with $a \in \mathbb{F}_q^\times$ and $b \in \mathbb{F}_q$. Therefore F_{n+1} is the splitting field of $f_n(T)$ over F_n and the extension F_{n+1}/F_n is Galois.

Let G_{n+1} be the Galois group of F_{n+1}/F_n . Every element $\sigma \in G_{n+1}$ acts on the function x_{n+1} as $\sigma(x_{n+1}) = a_\sigma x_{n+1} + b_\sigma$, and the map

$$\sigma \mapsto \begin{pmatrix} a_\sigma & 0 \\ b_\sigma & 1 \end{pmatrix}$$

is a monomorphism of G_{n+1} into the group of invertible 2×2 -matrices over \mathbb{F}_q of the form $\begin{pmatrix} a & 0 \\ b & 1 \end{pmatrix}$. This group has order $q(q-1)$, and hence $\text{ord}(G_{n+1})$ divides $q(q-1)$. \square

Lemma 2.4. Let $P_0 = (x_0 = \infty)$ be the pole of x_0 in F_0 and let P_n be a place of F_n above P_0 . For $i = 1, \dots, n$ we set $P_i := P_n|_{F_i}$ and $e_i := e(P_i|P_{i-1})$. Then the place P_i is a pole of x_i , $v_{P_i}(x_i)$ divides $(q-1)^i$ and $e_i \equiv 0 \pmod{q}$, for $1 \leq i \leq n$.

Proof. Let $u_i \in F_i$ be defined as in Eqn. (11). We prove the lemma by induction. For the case $i = 1$, we have $v_{P_1}(u_0) = e_1 \cdot v_{P_0}(u_0) = -e_1 \cdot (q-1)$. Since $(x_1^q - x_1)^{q-1} + 1 = u_0$, it follows that $v_{P_1}(x_1) < 0$ and therefore

$$v_{P_1}((x_1^q - x_1)^{q-1} + 1) = q \cdot (q-1) \cdot v_{P_1}(x_1).$$

We then conclude that $q \cdot v_{P_1}(x_1) = -e_1$. We finish this case since e_1 divides $[F_1 : F_0]$ and $[F_1 : F_0]$ divides $q(q-1)$ by Lemma 2.3, it follows that $v_{P_1}(x_1)$ divides $(q-1)$ and that $e_1 \equiv 0 \pmod{q}$.

Now we assume that $v_{P_i}(x_i) < 0$ and $v_{P_i}(x_i)$ divides $(q-1)^i$ for some $i \in \{1, \dots, n-1\}$. From Eqn. (11) we obtain $v_{P_i}(u_i) = (q-1) \cdot v_{P_i}(x_i)$, hence

$$v_{P_{i+1}}(u_i) = e_{i+1} \cdot (q-1) \cdot v_{P_i}(x_i) < 0.$$

Since $(x_{i+1}^q - x_{i+1})^{q-1} + 1 = u_i$, it follows that P_{i+1} is a pole of x_{i+1} and

$$q(q-1) \cdot v_{P_{i+1}}(x_{i+1}) = e_{i+1} \cdot (q-1) \cdot v_{P_i}(x_i).$$

Now we finish as in the case $i = 1$; one concludes that $e_{i+1} \equiv 0 \pmod{q}$ and that $v_{P_{i+1}}(x_{i+1})$ divides $(q-1)^{i+1}$. \square

Lemma 2.5. $[F_{n+1} : F_n] \equiv 0 \pmod{q}$ for all $n \geq 0$.

Proof. Follows directly from Lemmas 2.3 and 2.4. \square

Lemma 2.6. $[F_1 : F_0] = q(q-1)$, and K is the full constant field of F_1 .

Proof. By definition, $F_1 = K(x_0, x_1)$ with

$$(x_1^q - x_1)^{q-1} + 1 = \frac{-x_0^{q(q-1)}}{(x_0^{q-1} - 1)^{q-1}} = u_0. \quad (12)$$

It follows that

$$[K(x_0) : K(u_0)] = [K(x_1) : K(u_0)] = q(q-1). \quad (13)$$

From Eqn. (12) it is obvious that the place $(u_0 = 0)$ of $K(u_0)$ is totally ramified in the extension $K(x_0)/K(u_0)$. The place of $K(x_0)$ above $(u_0 = 0)$ is the place $(x_0 = 0)$, and we have $e((x_0 = 0)|(u_0 = 0)) = q(q-1)$.

However, in the extension $K(x_1)/K(u_0)$ the place $(u_0 = 0)$ is unramified, since the polynomial $(x_1^q - x_1)^{q-1} + 1$ does not have multiple roots. Let Q be a place of $K(x_1)$ lying above $(u_0 = 0)$ and let R be a place of $K(x_0, x_1)$ above Q . It follows that $e(R|Q) = q(q-1)$. Therefore $[K(x_0, x_1) : K(x_1)] = q(q-1)$, and K is algebraically closed in $K(x_0, x_1) = F_1$ (as there is a place which is totally ramified in $F_1/K(x_1)$). The assertion $[F_1 : F_0] = q(q-1)$ follows since $[F_1 : F_0] = [F_1 : K(x_1)]$ by Eqn. (13). \square

The next lemma shows a striking property of the recursion in Eqn.(8) for $n \geq 1$. It gives a simple Artin-Schreier equation for the extension F_{n+1}/F_n of degree q .

Lemma 2.7. For each $n \geq 1$ there is some $\mu \in \mathbb{F}_q^\times$ such that

$$x_{n+1}^q - x_{n+1} = \mu \cdot \frac{x_{n-1}^q}{(x_{n-1}^{q-1} - 1) \cdot (x_n^{q-1} - 1)}.$$

Proof. By Eqn. (8) we have

$$(x_{n+1}^q - x_{n+1})^{q-1} + 1 = \frac{-x_n^{q(q-1)}}{(x_n^{q-1} - 1)^{q-1}} \quad \text{and} \quad (x_n^q - x_n)^{q-1} + 1 = \frac{-x_{n-1}^{q(q-1)}}{(x_{n-1}^{q-1} - 1)^{q-1}}. \quad (14)$$

Hence we get

$$\begin{aligned} (x_{n+1}^q - x_{n+1})^{q-1} &= \frac{-x_n^{q(q-1)}}{(x_n^{q-1} - 1)^{q-1}} - 1 = \frac{-((x_n^q - x_n)^{q-1} + 1)}{(x_n^{q-1} - 1)^{q-1}} \\ &= \frac{x_n^{q(q-1)}}{(x_n^{q-1} - 1)^{q-1} \cdot (x_n^{q-1} - 1)^{q-1}} = \left(\frac{x_{n-1}^q}{(x_{n-1}^{q-1} - 1) \cdot (x_n^{q-1} - 1)} \right)^{q-1}. \end{aligned}$$

\square

Proof of Theorem 2.2. Putting together the results of the lemmas above, one gets the assertions in Theorem 2.2. \square

3 Splitting places in the tower over $K = \mathbb{F}_\ell$ for $\ell = q^3$

In this section we consider the tower $\mathcal{F} = (F_0, F_1, F_2, \dots)$ which was introduced in Sec. 2, over the field $K = \mathbb{F}_\ell$ with $\ell = q^3$. We will show that many rational places of the field $F_0 = \mathbb{F}_\ell(x_0)$ split completely in \mathcal{F} ; i.e., they split completely in all extensions F_n/F_0 . This means that the function fields F_n/\mathbb{F}_ℓ have “many” rational places. As in Sec. 2, let

$$f(T) = (T^q - T)^{q-1} + 1 \in \mathbb{F}_q[T]. \quad (15)$$

For $q = 2$ we have that $f(T) = c$ is separable for all elements $c \in \overline{\mathbb{F}}_2$.

Lemma 3.1. *Let $c \in \overline{\mathbb{F}}_q$ be an element of the algebraic closure of \mathbb{F}_q . Then*

$$f(T) = c \text{ is inseparable if and only if } c = 1 \text{ and } q \neq 2.$$

For an element $\beta \in \overline{\mathbb{F}}_q$ we have that $f(\beta) = 1$ if and only if β belongs to \mathbb{F}_q .

Proof. Just notice that the derivative of $f(T)$ satisfies $f'(T) = (T^q - T)^{q-2}$. □

Lemma 3.2. *For an element $\beta \in \overline{\mathbb{F}}_q$ we have that $f(\beta) = 0$ if and only if $\beta \in \mathbb{F}_{q^2} \setminus \mathbb{F}_q$.*

Proof. Just notice that we have (see Remark 2.1)

$$f(T) = (T^{q^2} - T)/(T^q - T). \quad (16)$$

□

Now we consider the recursive equation for the tower \mathcal{F} (see Equation (10)):

$$f(Y) = \frac{1}{1 - f(1/X)}. \quad (17)$$

We will show that if $X = \alpha$ belongs to $\mathbb{F}_{q^3} \setminus \mathbb{F}_q$ then all solutions $Y = \beta \in \overline{\mathbb{F}}_q$ of Equation (17) with $X = \alpha$ are such that $\beta \in \mathbb{F}_{q^3} \setminus \mathbb{F}_q$. The assertion that $\beta \notin \mathbb{F}_q$ follows directly from Equation (17) and the lemmas above.

Using Equation (16) we have:

$$\frac{1}{1 - f(T)} = \frac{T - T^q}{T^{q^2} - T^q}. \quad (18)$$

Lemma 3.3. *For an element $\beta \in \overline{\mathbb{F}}_q$ we have that*

$$f(\beta)^q = \frac{1}{1 - f(\beta)} \quad \text{if and only if} \quad \beta \in \mathbb{F}_{q^3} \setminus \mathbb{F}_q.$$

Proof. Straightforward using Equations (16) and (18). □

Equation (17) can also be written as below:

$$f\left(\frac{1}{X}\right) = 1 - \frac{1}{f(Y)}. \quad (19)$$

Consider now a solution (α, β) of Equation (17) with $\alpha \in \mathbb{F}_{q^3} \setminus \mathbb{F}_q$. Then $1/\alpha \in \mathbb{F}_{q^3} \setminus \mathbb{F}_q$. We have:

$$f(\beta) = \frac{1}{1 - f\left(\frac{1}{\alpha}\right)} = f\left(\frac{1}{\alpha}\right)^q = 1 - \frac{1}{f(\beta)^q}.$$

In the last two equalities above we have used Lemma 3.3 and Equation (19), respectively. Hence we obtained that $f(\beta)^q = 1/1 - f(\beta)$; i.e., $\beta \in \mathbb{F}_{q^3} \setminus \mathbb{F}_q$.

We have then proved the main result of this section:

Theorem 3.4. *Let $\mathcal{F} = (F_0, F_1, \dots)$ be the tower over \mathbb{F}_{q^3} given recursively by Equation (17). Then the places $(x_0 = \alpha)$ with $\alpha \in \mathbb{F}_{q^3} \setminus \mathbb{F}_q$ split completely in all extensions F_n/F_0 . In particular the number of \mathbb{F}_{q^3} -rational places satisfies:*

$$N(F_n) \geq (q^3 - q) \cdot [F_n : F_0] \quad \text{for all } n \in \mathbb{N}.$$

4 The extensions $K(x)/K(w)$ and $K(x)/K(u)$

Throughout this section, K is a field with $\mathbb{F}_{q^2} \subseteq K$. Let $K(x)/K$ be a rational function field over K . We will consider certain subfields $K(w) \subseteq K(x)$ and $K(u) \subseteq K(x)$ which are related to the recursive definition of the tower \mathcal{F} . Detailed information about ramification in $K(x)/K(w)$ and in $K(x)/K(u)$ will enable us to study in Sec. 5 and Sec. 6 the ramification behaviour in the tower \mathcal{F} .

As in Sec. 2 we consider the polynomial $f(T) = (T^q - T)^{q-1} + 1 \in K[T]$, and we set

$$w := f(x) = (x^q - x)^{q-1} + 1 \in K(x). \quad (20)$$

Lemma 4.1. (i) *The extension $K(x)/K(w)$ is Galois of degree $q(q-1)$.*

(ii) *The place $(w = \infty)$ of $K(w)$ is totally ramified in $K(x)/K(w)$; the place above it is the place $(x = \infty)$. We have $d((x = \infty)|(w = \infty)) = q^2 - 2$; i.e., $(x = \infty)|(w = \infty)$ is weakly ramified.*

(iii) *Above the place $(w = 1)$ there are the q places $(x = \theta)$ of $K(x)$ with $\theta \in \mathbb{F}_q$, with ramification index $e((x = \theta)|(w = 1)) = q - 1$.*

(iv) *All other places of $K(w)$ are unramified in $K(x)/K(w)$.*

(v) *The places above $(w = 0)$ are exactly the places $(x = \beta)$ with $\beta \in \mathbb{F}_{q^2} \setminus \mathbb{F}_q$.*

Proof. i) One checks easily that $K(w)$ is the fixed field of the following group H of automorphisms of $K(x)/K$:

$$H := \{\sigma \in \text{Aut}(K(x)/K) \mid \sigma(x) = ax + b, a \in \mathbb{F}_q^\times, b \in \mathbb{F}_q\}.$$




Figure 2: Ramification in $K(x)/K(u)$.

Proof. Note that $u = 1/(1 - f(1/x))$ by Rem. 2.1 and therefore $f(1/x) = (u - 1)/u$.

The result follows directly from Lemma 4.1 with the changes of variables

$$x \mapsto 1/x \quad \text{and} \quad w \mapsto (u - 1)/u.$$

□

5 The fields F_1 and F_2

In this section we assume again that $\mathbb{F}_{q^2} \subseteq K$. We want to study ramification in the first two steps of the tower \mathcal{F} over K . So we consider the fields $F_0 = K(x_0)$, $F_1 = K(x_0, x_1)$ and $F_2 = K(x_0, x_1, x_2)$ where

$$(x_1^q - x_1)^{q-1} + 1 = \frac{-x_0^{q(q-1)}}{(x_0^{q-1} - 1)^{q-1}} \quad \text{and} \quad (x_2^q - x_2)^{q-1} + 1 = \frac{-x_1^{q(q-1)}}{(x_1^{q-1} - 1)^{q-1}}. \quad (22)$$

Lemma 5.1. *The extensions $F_1/K(x_0)$ and $F_1/K(x_1)$ are both Galois of degree $q(q-1)$.*

Proof. We proved the assertion for $F_1/K(x_0)$ in Thm. 2.2. As in Eqn.(11) we set

$$u_0 := \frac{-x_0^{q(q-1)}}{(x_0^{q-1} - 1)^{q-1}}.$$

The field F_1 is the compositum of $K(x_0)$ and $K(x_1)$ over $K(u_0)$ as in Figure 3. By Lemma 4.2 the extension $K(x_0)/K(u_0)$ is Galois, hence $F_1/K(x_1)$ is Galois as well. □




Figure 3: The extension $F_1/K(u_0)$

Lemma 5.2. Let $\Omega := \mathbb{F}_{q^2} \cup \{\infty\}$.

(i) For a place $P \in \mathbb{P}(F_1)$ the following are equivalent:

a) $P|_{K(x_0)} = (x_0 = \omega)$ for some $\omega \in \Omega$.

b) $P|_{K(x_1)} = (x_1 = \omega')$ for some $\omega' \in \Omega$.

(ii) If a place $Q \in \mathbb{P}(F_1)$ does not lie above a place $(x_0 = \omega)$ with $\omega \in \Omega$ then Q is unramified over $K(x_0)$ and over $K(x_1)$.

(iii) The ramification indices of the places $(x_0 = \omega)$ and $(x_1 = \omega')$ with $\omega, \omega' \in \Omega$ in the extensions $F_1/K(x_0)$ and $F_1/K(x_1)$ are as depicted in Figure 4. All places of F_1 are weakly ramified over $K(x_0)$ and over $K(x_1)$.




Figure 4: Ramification in $F_1/K(x_0)$ and in $F_1/K(x_1)$.

Proof. According to the notations in Sec. 4 we write

$$u_0 := \frac{-x_0^{q(q-1)}}{(x_0^{q-1} - 1)^{q-1}} \quad \text{and} \quad w_1 := (x_1^q - x_1)^{q-1} + 1.$$

Hence $u_0 = w_1$ by Eqn. (22). We consider the diagram of fields in Figure 3 where all extensions are Galois of degree $q(q-1)$. By Lemma 4.1 and Lemma 4.2 we know that only the places $(u_0 = 0)$, $(u_0 = 1)$ and $(u_0 = \infty)$ are ramified in $K(x_0)/K(u_0)$ or

in $K(x_1)/K(u_0)$. We will consider here only the case ($u_0 = \infty$); the other two cases are similar (even easier). Denote by Q a place of F_1 above ($u_0 = \infty$). The situation is depicted in Figure 5. It follows from Abhyankar's Lemma (see [15, Prop. III.8.9]) that Q




Figure 5: Ramification in $F_1/K(u_0)$

is unramified over $K(x_1)$ and that the ramification index of Q over $K(x_0)$ is $e = q$. Since $(x_1 = \infty)|(u_0 = \infty)$ is weakly ramified by Lemma 4.1 it follows from the transitivity of different exponents in $F_1 \supseteq K(x_0) \supseteq K(u_0)$ that Q is weakly ramified over $K(x_0)$. \square

Lemma 5.3. *The extensions $F_2/K(x_0, x_1)$ and $F_2/K(x_1, x_2)$ are Galois extensions of degree q . All places that are ramified in $F_2/K(x_0, x_1)$ or in $F_2/K(x_1, x_2)$ are totally and weakly ramified.*

Proof. The field F_2 is the compositum of $K(x_0, x_1)$ and $K(x_1, x_2)$ over $K(x_1)$. Since the extensions $K(x_0, x_1)/K(x_1)$ and $K(x_1, x_2)/K(x_1)$ are Galois by Lemma 5.1, it is clear that $F_2/K(x_0, x_1)$ and $F_2/K(x_1, x_2)$ are Galois. The assertion about the degrees follows from Lemma 2.7. Now we consider a place $Q \in \mathbb{P}(F_2)$ which is ramified in $F_2/K(x_1, x_2)$. Then the place $P := Q|_{K(x_0, x_1)}$ is ramified over $K(x_1)$ and therefore $Q|_{K(x_1)} = (x_1 = \beta)$ with some $\beta \in \mathbb{F}_{q^2} \setminus \mathbb{F}_q$, by Lemma 5.2. So we have the situation depicted in Figure 6, where R denotes the restriction of Q to $K(x_1, x_2)$




Figure 6:

As in the proof of Lemma 5.2, use Abhyankar's lemma to get that $e(Q|R) = q$ and the transitivity of different exponents to get that $d(Q|R) = 2 \cdot (q - 1)$.

Now if Q is a place of F_2 which is ramified over F_1 , then one also concludes (and it is simpler) that it is totally and weakly ramified over F_1 . \square

6 The genus of F_n

In order to estimate the limit $\lambda(\mathcal{F})$ of the tower \mathcal{F} over \mathbb{F}_{q^3} we need an upper bound for the genus of the n -th function field F_n ; therefore one has to study ramification in the extension F_n/F_0 . Without changing the ramification behaviour (i.e., ramification index and different exponent) and the genus, we can extend the constant field such that it contains \mathbb{F}_{q^2} . So we assume in this section that $\mathbb{F}_{q^2} \subseteq K$ and denote $\text{char}(K) = p$.

A place $P \in \mathbb{P}(F_0)$ is said to be ramified in the tower \mathcal{F} if P is ramified in F_m/F_0 for some $m \geq 1$, and the ramification locus $V(\mathcal{F}/F_0)$ is defined as

$$V(\mathcal{F}/F_0) := \{P \in \mathbb{P}(F_0) \mid P \text{ is ramified in } \mathcal{F}\}.$$

Lemma 6.1. *The ramification locus of \mathcal{F} over F_0 satisfies*

$$V(\mathcal{F}/F_0) \subseteq \{(x_0 = \omega) \mid \omega \in \mathbb{F}_{q^2} \text{ or } \omega = \infty\}.$$

Proof. Assume that a place $Q \in \mathbb{P}(F_n)$ is ramified in F_{n+1}/F_n . Then the restriction $Q|_{K(x_n)}$ ramifies in the extension $K(x_n, x_{n+1})/K(x_n)$. We conclude from Lemma 5.2 ii) that $Q|_{K(x_n)} = (x_n = \omega')$ with $\omega' \in \mathbb{F}_{q^2} \cup \{\infty\}$. By induction it follows from Lemma 5.2 i) that $Q|_{F_0} = (x_0 = \omega)$ with $\omega \in \mathbb{F}_{q^2} \cup \{\infty\}$. This proves the lemma. We remark that in fact $V(\mathcal{F}/F_0) = \{(x_0 = \omega) \mid \omega \in \mathbb{F}_{q^2} \text{ or } \omega = \infty\}$ but we do not need this here. \square

In the proof of Lemma 6.3 below, the following result is essential

Lemma 6.2. *Consider an extension E/F of function fields over K such that $E = E_1 \cdot E_2$ is the composite field of two intermediate fields $F \subseteq E_1, E_2 \subseteq E$ and the extensions E_1/F and E_2/F are Galois p -extensions. Let Q be a place of E , and let $Q_i := Q|_{E_i}$ and $P := Q|_F$ be the restrictions of Q . Suppose that $Q_1|P$ and $Q_2|P$ are weakly ramified. Then $Q|Q_1$ and $Q|Q_2$ are also weakly ramified.*

Proof. See [10, Prop. 1.10] and also [9, Lemma 1]. \square

A Galois extension E/F is *weakly ramified* if all places are weakly ramified in E/F .

Lemma 6.3. *Let $n \geq 1$. Then the extension F_{n+1}/F_n is weakly ramified.*

Proof. For $0 \leq i \leq j \leq n+1$ we define the subfield $E_{i,j} \subseteq F_{n+1}$ by

$$E_{i,j} := K(x_i, x_{i+1}, \dots, x_j).$$

The extensions $E_{i,i+2}/E_{i,i+1}$ and $E_{i,i+2}/E_{i+1,i+2}$ are weakly ramified Galois p -extensions by Lemma 5.3 (see Figure 7). By induction it follows for all $j \geq i+2$ that $E_{i,j}/E_{i,j-1}$ and $E_{i,j}/E_{i+1,j}$ are weakly ramified Galois p -extensions (using Lemma 6.2). Since $F_n = E_{0,n}$ and $F_{n+1} = E_{0,n+1}$, the assertion of Lemma 6.3 follows. \square




Figure 7: Bold face means Galois weakly ramified p -extensions, by Lemma 5.3.

Lemma 6.4. *Let E_1/F be a Galois extension of function fields over K and let E/E_1 be a finite and separable extension. Let Q be a place of the field E and denote by P_1 and P the restrictions of Q to E_1 and F , respectively. Suppose that we have:*

- (i) $e(Q|P_1)$ is a power of $p = \text{char}(K)$ and $d(Q|P_1) = 2e(Q|P_1) - 2$.
- (ii) The place P_1 is weakly ramified over P .

Then the different exponent $d(Q|P)$ satisfies

$$d(Q|P) = (e_0e_1 - 1) + (e_1 - 1) < e(Q|P) \cdot \left(1 + \frac{1}{e_0}\right)$$

where $e(Q|P) = e_0e_1$ with $(p, e_0) = 1$ and e_1 a p -power.

Proof. Straightforward, using transitivity of different exponents. \square

Theorem 6.5. *The genus of the n -th function field of the tower $\mathcal{F} = (F_0, F_1, F_2, \dots)$ defined by Eqn.(8), satisfies*

$$g(F_n) \leq \frac{q^2 + 2q}{2} \cdot [F_n : F_0].$$

Proof. Let $n \geq 1$. First we observe that for a place $Q \in \mathbb{P}(F_n)$ and the restriction $P_1 := Q|_{F_1}$ of Q to F_1 we have that:

$$e(Q|P_1) \text{ is a } p\text{-power and } d(Q|P_1) = 2e(Q|P_1) - 2.$$

This follows from Lemma 6.3 and repeated applications of Lemma 6.4. In fact one takes $E = F_n$, $E_1 = F_j$ and $F = F_{j-1}$ with $j = n - 1, n - 2, \dots, 2$.

Now we consider the places $P \in \mathbb{P}(F_0)$ which are in the ramification locus $V(\mathcal{F}/F_0)$. According to item (iii) of Lemma 5.2 we distinguish 2 cases:

Case 1: $P = (x_0 = \theta)$ with $\theta \in \mathbb{F}_q$ or $P = (x_0 = \infty)$.

By Lemma 5.2 and Lemma 6.4 we obtain

$$\sum_{\substack{Q \in \mathbb{P}(F_n) \\ Q|P}} d(Q|P) \cdot \deg Q < \sum_{\substack{Q \in \mathbb{P}(F_n) \\ Q|P}} 2e(Q|P) \cdot \deg Q = 2[F_n : F_0]. \quad (23)$$

Case 2: $P = (x_0 = \beta)$ with $\beta \in \mathbb{F}_{q^2} \setminus \mathbb{F}_q$.

In this case, Lemma 5.2 and Lemma 6.4 yield

$$\sum_{\substack{Q \in \mathbb{P}(F_n) \\ Q|P}} d(Q|P) \cdot \deg Q < \sum_{\substack{Q \in \mathbb{P}(F_n) \\ Q|P}} \left(1 + \frac{1}{q-1}\right) e(Q|P) \cdot \deg Q = \frac{q}{q-1} [F_n : F_0]. \quad (24)$$

There are $q + 1$ places $P \in \mathbb{P}(F_0)$ as in Case 1, and $q^2 - q$ places as in Case 2. By Hurwitz genus formula for the extension F_n/F_0 we obtain

$$\begin{aligned} 2g(F_n) &\leq -2[F_n : F_0] + (q + 1) \cdot 2[F_n : F_0] + (q^2 - q) \cdot \frac{q}{q-1} [F_n : F_0] \\ &= (q^2 + 2q)[F_n : F_0]. \end{aligned}$$

□

7 The limit of the tower over $K = \mathbb{F}_\ell$ with $\ell = q^3$

Putting together the results of the previous sections we obtain our main result:

Theorem 7.1. *Let $K = \mathbb{F}_\ell$ with $\ell = q^3$, and let $\mathcal{F} = (F_0, F_1, F_2, \dots)$ be the tower over K which is recursively defined by $F_0 = K(x_0)$ and $F_{n+1} = F_n(x_{n+1})$, where*

$$(x_{n+1}^q - x_{n+1})^{q-1} + 1 = \frac{-x_n^{q(q-1)}}{(x_n^{q-1} - 1)^{q-1}} \quad \text{for all } n \geq 0.$$

Then the limit $\lambda(\mathcal{F}) = \lim_{n \rightarrow \infty} N(F_n)/g(F_n)$ satisfies $\lambda(\mathcal{F}) \geq 2(q^2 - 1)/(q + 2)$.

Proof. By Thm. 3.4 and Thm. 6.5 we have

$$N(F_n) \geq (q^3 - q) \cdot [F_n : F_0] \quad \text{and} \quad g(F_n) \leq \frac{q^2 + 2q}{2} \cdot [F_n : F_0].$$

Hence

$$\frac{N(F_n)}{g(F_n)} \geq \frac{(q^3 - q) \cdot 2}{q^2 + 2q} = \frac{2(q^2 - 1)}{q + 2} \quad \text{for all } n \geq 0.$$

□

8 Remarks

We finish this paper with a few remarks.

Remark 8.1. Our tower $\mathcal{F} = (F_0, F_1, F_2, \dots)$ over $K = \mathbb{F}_{q^3}$ bears remarkable analogy to the tower $\mathcal{H} = (H_0, H_1, H_2, \dots)$ over the quadratic field $K = \mathbb{F}_{q^2}$ which is defined recursively by the equation

$$u_{i+1}^q + u_{i+1} = \frac{u_i^q}{u_i^{q-1} + 1}$$

and which attains the Drinfel'd-Vlăduț bound (1). The analogies between \mathcal{H} and \mathcal{F} become even more evident if we substitute $u_i = \xi y_i$ with $\xi^{q-1} = -1$; then the above equation becomes $y_{i+1}^q - y_{i+1} = -y_i^q / (y_i^{q-1} - 1)$. We now compare some features of the towers \mathcal{F} over \mathbb{F}_{q^3} and \mathcal{H} over \mathbb{F}_{q^2} :

- 1) The tower $\mathcal{H} = (H_0, H_1, H_2, \dots)$ is defined recursively over the field $K = \mathbb{F}_{q^2}$ by $H_0 = K(y_0)$ and $H_{i+1} = H_i(y_{i+1})$, where

$$y_{i+1}^q - y_{i+1} = \frac{-y_i^q}{y_i^{q-1} - 1} \quad \text{for all } i \geq 0. \quad (25)$$

- 2) Setting $h(T) := T^q - T$, Eqn. (25) can be written as

$$h(y_{i+1}) = \frac{1}{h(1/y_i)}. \quad (26)$$

- 3) The extensions H_{i+1}/H_i (for $i \geq 0$) are weakly ramified and Galois of degree q .
- 4) The ramification locus of \mathcal{H} over H_0 is

$$V(\mathcal{H}/H_0) = \{(y_0 = \omega) \mid \omega \in \mathbb{F}_q \cup \{\infty\}\}.$$

- 5) The places $(y_0 = \alpha)$ with $\alpha \in \mathbb{F}_{q^2} \setminus \mathbb{F}_q$ are completely splitting in all the extensions H_n/H_0 , for all $n \geq 0$.

The analogous properties of the tower \mathcal{F} are:

- 1*) The tower $\mathcal{F} = (F_0, F_1, F_2, \dots)$ is defined recursively over the field $K = \mathbb{F}_{q^3}$ by $F_0 = K(x_0)$ and $F_{i+1} = F_i(x_{i+1})$, where

$$(x_{i+1}^q - x_{i+1})^{q-1} + 1 = \frac{-x_i^{q(q-1)}}{(x_i^{q-1} - 1)^{q-1}} \quad \text{for all } i \geq 0. \quad (27)$$

- 2*) Setting $f(T) := (T^q - T)^{q-1} + 1$, Eqn. (27) can be written as

$$f(x_{i+1}) = \frac{1}{1 - f(1/x_i)}. \quad (28)$$

3*) The extensions F_{i+1}/F_i (for $i \geq 1$) are weakly ramified and Galois of degree q .

4*) The ramification locus of \mathcal{F} over F_0 is

$$V(\mathcal{F}/F_0) = \{(x_0 = \omega) \mid \omega \in \mathbb{F}_{q^2} \cup \{\infty\}\}.$$

5*) The places $(x_0 = \alpha)$ with $\alpha \in \mathbb{F}_{q^3} \setminus \mathbb{F}_q$ are completely splitting in all the extensions F_n/F_0 , for all $n \geq 0$.

We also note that the polynomials $h(T)$ and $f(T)$ in Eqns.(26) and (28) are defined in a very similar manner:

6) The polynomial $h(T) \in \mathbb{F}_q[T]$ generates the fixed field of $K(T)$ under the group of automorphisms

$$G = \{\sigma : K(T) \rightarrow K(T) \mid \sigma(T) = T + b \text{ with } b \in \mathbb{F}_q\}.$$

6*) The polynomial $f(T) \in \mathbb{F}_q[T]$ generates the fixed field of $K(T)$ under the group of automorphisms

$$G^* = \{\sigma : K(T) \rightarrow K(T) \mid \sigma(T) = aT + b \text{ with } a \in \mathbb{F}_q^\times \text{ and } b \in \mathbb{F}_q\}.$$

Another interesting observation is that the generators x_i of the tower \mathcal{F} satisfy

$$x_{i+2}^q - x_{i+2} = \frac{-x_i^q}{(x_i^{q-1} - 1)(x_{i+1}^{q-1} - 1)} \quad (29)$$

for all $i \geq 0$ (with an appropriate choice of the roots x_{i+1}, x_{i+2} of Eqn. (27), see Lemma 2.7). Compare with Eqn. (25).

Remark 8.2. The first explicit tower over a field with cubic cardinality $\ell = q^3$ which attains the Zink bound (Inequality (2)) was found by van der Geer–van der Vlugt [12]. It is a tower over the field \mathbb{F}_{p^3} with $p = 2$, recursively defined by the equation

$$x_{i+1}^2 + x_{i+1} = x_i + 1 + \frac{1}{x_i}. \quad (30)$$

This is the special case $q = 2$ of Eqn. (27) (after the change of variables $x_i \rightarrow x_i + 1$).

Remark 8.3. Again we consider the tower $\mathcal{F} = (F_0, F_1, F_2, \dots)$ over $K = \mathbb{F}_{q^3}$. We set

$$v_i := -\frac{1}{x_i^{q-1} - 1} \quad \text{for all } i \geq 0. \quad (31)$$

It follows by straightforward calculation from Eqn. (27) that

$$\frac{1 - v_{i+1}}{v_{i+1}^q} = \frac{v_i^q + v_i - 1}{v_i}, \quad \text{for all } i \geq 0. \quad (32)$$

This means that \mathcal{F} contains as a subtower the tower $\mathcal{E} = (E_0, E_1, E_2, \dots)$ (see [3]) with $E_0 = K(v_0)$ and $E_{i+1} = E_i(v_{i+1})$, where v_{i+1} satisfies Eqn. (32) over E_i . Since the limit of a subtower is at least as big as the limit of the tower itself (see [8]), we obtain that

$$\lambda(\mathcal{E}) \geq \lambda(\mathcal{F}) \geq \frac{2(q^2 - 1)}{q + 2}.$$

This gives another (in fact, much simpler) proof of the main result of [3].

Here is another striking analogy between \mathcal{F} and \mathcal{H} ; again we consider the tower $\mathcal{H} = (H_0, H_1, H_2, \dots)$ over $K = \mathbb{F}_{q^2}$ given recursively by

$$u_{i+1}^q + u_{i+1} = \frac{u_i^q}{u_i^{q-1} + 1}. \quad (33)$$

Performing the analogous change of variables as in Eqn.(31); i.e., setting

$$w_i := -\frac{1}{u_i^{q-1} + 1} \quad \text{for all } i \geq 0,$$

it follows by straightforward calculation from Eqn.(33) that

$$\frac{w_{i+1} + 1}{w_{i+1}^q} = \frac{w_i^q + 1}{w_i}, \quad \text{for all } i \geq 0. \quad (34)$$

The subtower \mathcal{G} of \mathcal{H} given recursively by Eqn.(34) was studied in [2].

Remark 8.4. We end up this paper with a closer look on the relations between the towers \mathcal{F} and \mathcal{E} given by Eqns. (27) and (32), respectively. One can show that F_1/E_1 is a Galois extension of degree $(q-1)^2$ with group $\mathbb{F}_q^\times \times \mathbb{F}_q^\times$; in fact the automorphisms of $F_1 = \mathbb{F}_{q^3}(x_0, x_1)$ over the subfield $E_1 = \mathbb{F}_{q^3}(v_0, v_1)$ are given by:

$$x_0 \mapsto ax_0 \text{ and } x_1 \mapsto bx_1, \text{ with } a, b \in \mathbb{F}_q^\times.$$

Moreover the n -th field F_n of the tower \mathcal{F} is the compositum with F_1 of the n -th field E_n of the tower \mathcal{E} ; i.e., we have

$$F_n = E_n \cdot F_1, \quad \text{for all } n \geq 1.$$

The assertions above follow from Eqns. (31) and (29).

Acknowledgment

We would like to thank Y. Ihara for his interest in and helpful discussions about splitting places in the tower \mathcal{E} (defined by Eqn. (32)).

References

- [1] A. Bassa, H. Stichtenoth, *A simplified proof for the limit of a tower over a cubic finite field*, J. Number Theory **123**, 2007, 154-169.
- [2] J. Bezerra, A. Garcia, *A tower with non-Galois steps which attains the Drinfeld-Vladut bound*, J. Number Theory **106**, 2004, 142-154.
- [3] J. Bezerra, A. Garcia, H. Stichtenoth, *An explicit tower of function fields over cubic finite fields and Zink's lower bound*, J. Reine Angew. Math. **589**, 2005, 159-199.
- [4] V. G. Drinfel'd, S. G. Vlăduț, *The number of points of an algebraic curve*, Func. Anal. **17**, 1983, 53-54.
- [5] N. Elkies, *Explicit modular towers*, Proceedings of the 35th Annual Allerton Conference on Communication, Control and Computing (eds. T. Basar et al.) Urbana IL, 1997, 23-32.
- [6] N. Elkies, *Explicit towers of Drinfeld modular curves*, European Congress of Math. (Barcelona, 2000) Vol. II, 189-198, Progr. Math., **202**, Birkhäuser, Basel, 2001.
- [7] A. Garcia, H. Stichtenoth, *A tower of Artin-Schreier extensions of function fields attaining the Drinfeld-Vladut bound*, Inventiones Math. **121**, 1995, 211-222.
- [8] A. Garcia, H. Stichtenoth, *On the asymptotic behaviour of some towers of function fields over finite fields*, J. Number Theory **61**, 1996, 248-273.
- [9] A. Garcia, H. Stichtenoth, *Some Artin-Schreier towers are easy*, Mosc. Math. J. **5**, 2005, 767-774.
- [10] A. Garcia, H. Stichtenoth, *On the Galois closure of towers*, Recent Trends in Coding Theory and its Applications (W. Li, ed.), to appear.
- [11] A. Garcia, H. Stichtenoth, *Explicit towers of function fields over finite fields*, Topics in geometry, coding theory and cryptography, 1-58, Algebr. Appl., **6**, Springer, Dordrecht, 2007.
- [12] G. van der Geer, M. van der Vlugt, *An asymptotically good tower of curves over the field with eight elements*, Bull. London Math. Soc. **34**, 2002, 291-300.
- [13] Y. Ihara, *Some remarks on the number of rational points of algebraic curves over finite fields*, J. Fac. Sci. Univ. Tokyo Sect. IA Math. **28**, 1981, 721-724.
- [14] J.-P. Serre, *Sur le nombre des points rationnels d'une courbe algébrique sur un corps fini*, C. R. Acad. Sci. Paris **296**, 1983, 397-402.
- [15] H. Stichtenoth, *Algebraic function fields and codes*, Springer Verlag, Berlin, 1993.
- [16] M. A. Tsfasman, S. G. Vlăduț, T. Zink, *Modular curves, Shimura curves, and Goppa codes, better than the Varshamov-Gilbert bound*, Math. Nachr. **109**, 1982, 21-28.
- [17] T. Zink, *Degeneration of Shimura surfaces and a problem in coding theory*, in Fundamentals of Computation Theory (L. Budach, ed.), Lecture Notes in Computer Science, Vol. **199**, Springer Verlag, Berlin, 1985, 503-511.