

# THE CHARACTERISTIC VARIETY OF A GENERIC FOLIATION

JORGE VITÓRIO PEREIRA

ABSTRACT. We confirm a conjecture of Bernstein-Lunts which predicts that the characteristic variety of a generic polynomial vector field has no homogeneous involutive subvarieties besides the zero section and subvarieties of fibers over singular points.

## CONTENTS

1. Introduction	1
2. Characteristic varieties and prolongations	3
3. Warm-up: Proof of Theorem 1 in dimension three	4
4. Prolongation versus holonomy	6
5. From invariant subvarieties to multi-distributions	8
6. From multi-distributions to invariant subvarieties	9
7. Proof of Theorem 1	10
8. Bernstein-Lunts Conjecture	11
References	11

## 1. INTRODUCTION

1.1. **Foliations.** Let  $\mathcal{F}$  be a one-dimensional singular holomorphic foliation on a smooth projective variety  $X$ . The **characteristic variety**  $\text{ch}(\mathcal{F})$  of  $\mathcal{F}$  is the irreducible subvariety of  $E(T^*X)$ , the total space of the cotangent bundle of  $X$ , with fiber over a non-singular point  $x \in X_0 = X \setminus \text{sing}(\mathcal{F})$  equal to the 1-forms at  $x$  which vanish on  $T_x\mathcal{F}$ . More succinctly, if  $N^*\mathcal{F}$  is the conormal sheaf of  $\mathcal{F}$  then its restriction at  $X_0$  is a vector sub-bundle of  $T^*X_0$  and we can write

$$\text{ch}(\mathcal{F}) = \overline{E(N^*\mathcal{F}|_{X_0})}$$

where the closure is taken in  $E(T^*X) \supset E(T^*X_0)$ .

Clearly  $\text{ch}(\mathcal{F})$  is a hypersurface of  $E(T^*X)$ . If  $\omega$  is the non-degenerate 2-form which induces the canonical symplectic structure on  $T^*X$  then its restriction to  $\text{ch}(\mathcal{F})$  induces a one-dimensional foliation  $\mathcal{F}^{(1)}$  on (the smooth locus of)  $\text{ch}(\mathcal{F})$  which will be called the **first prolongation of  $\mathcal{F}$** .

In this work we are interested in the subvarieties of  $\text{ch}(\mathcal{F})$  invariant by  $\mathcal{F}^{(1)}$  when  $\mathcal{F}$  is sufficiently general. For no matter which  $\mathcal{F}$  there is always at least one subvariety of  $\text{ch}(\mathcal{F})$  invariant by  $\mathcal{F}^{(1)}$ : the zero section of  $T^*X$ . If the singular set of

---

1991 *Mathematics Subject Classification.* Primary: 37F75, 16S32; Secondary: 37J30, 32C38.  
*Key words and phrases.* characteristic foliation, invariant variety,  $\mathcal{D}$ -modules.

$\mathcal{F}$  is non-empty but of dimension zero then the fibers over it, and some subvarieties of these fibers, are also left invariant by  $\mathcal{F}^{(1)}$ .

We will say that  $\text{ch}(\mathcal{F})$  is a **quasi-minimal characteristic variety** if (a)  $\mathcal{F}$  has isolated singularities; and (b) every irreducible homogeneous ( on the fibers of  $\text{ch}(\mathcal{F}) \rightarrow X$  ) subvariety of  $\text{ch}(\mathcal{F})$  left invariant by  $\mathcal{F}^{(1)}$  is either the zero section, or a subvariety of a fiber over the singular set of  $\mathcal{F}$ , or the whole  $\text{ch}(\mathcal{F})$ .

**Theorem 1.** *Let  $X$  be a smooth projective variety,  $\mathcal{L}$  an ample line bundle over it, and  $k \gg 0$  a sufficiently large integer. If  $\mathcal{F} \in \mathbb{P}H^0(X, TX \otimes \mathcal{L}^{\otimes k})$  is a very generic foliation then  $\text{ch}(\mathcal{F})$  is a quasi-minimal characteristic variety.*

In the statement of the theorem above and throughout, by a very generic point of a given variety we mean a point outside a countable union of Zariski closed subvarieties. The expression generic point will be reserved to points outside a finite union of Zariski closed subvarieties.

Although Theorem 1 can be thought as a natural development of Jouanolou's Theorem and its subsequent generalizations, see [6] and references therein, it is motivated by a problem coming from the representation theory of Weyl algebras that we briefly review below.

**1.2. Weyl algebras.** Let  $A_n$  be the  $n$ -th Weyl algebra over  $\mathbb{C}$ , that is  $A_n$  is the algebra of  $\mathbb{C}$ -linear differential operators on the polynomial ring  $\mathbb{C}[x_1, \dots, x_n]$ . A basic invariant of an irreducible  $A_n$ -module  $M$  is its Gelfand-Kirillov dimension  $GK \dim M$ . After Bernstein this invariant is subject to the inequality  $GK \dim M \geq n$  and equality holds true for important classes of irreducible  $A_n$ -modules, see [2]. If  $GK \dim M = n$  then  $M$  is, by definition, a holonomic  $A_n$ -module

For some time, some believed that every irreducible  $A_n$ -module  $M$  is holonomic. In 1985 Stafford came up with examples of  $A_n$ -modules of particularly simple form and having Gelfand-Kirillov dimension equal to  $2n - 1$ . His examples are of the form  $A_n/IA_n$  where  $I$  is a principal left ideal generated by an element of the form  $\xi + f$  where  $\xi$  is a polynomial vector field and  $f$  is polynomial, see [10]. For those not familiar with the Gelfand-Kirillov dimension it is useful to remark that when  $I$  is a principal maximal left ideal then  $GK \dim A_n/IA_n = 2n - 1$ , and the search of examples of non-holonomic  $A_n$ -modules can be reduced to search of principal maximal left ideals of  $A_n$ .

Stafford's examples are explicit and his arguments are purely algebraic. In [3], Bernstein and Lunts present two geometrically oriented approaches to construct principal maximal left ideals of  $A_n$ , and implement them for the second Weyl algebra. In rough terms, their strategy rely on the the study of a natural foliation defined on the characteristic varieties of the module. More specifically they relate the maximality of the ideal to the non-existence of proper invariant subvarieties of this foliation. To define a characteristic variety for a  $A_n$ -module, a filtration of  $A_n$  has to be fixed and their two approaches are determined by the choice of two different filtrations.

In the first approach they look at the Bernstein filtration of  $A_n$ , the  $i$ -th piece  $A_n^i$  consists of polynomials in  $\{x_1, \dots, x_n, \partial_{x_1}, \dots, \partial_{x_n}\}$  of degree at most  $i$ . The corresponding symbol maps are

$$\sigma_k : A_n^k \longrightarrow \frac{A_n^k}{A_n^{k-1}} \simeq \mathbb{C}_k[x_1, \dots, x_n, y_1, \dots, y_n].$$

They proved that if  $n = 2$ ,  $k \geq 4$  and  $P \in \mathbb{C}_k[x_1, \dots, x_n, y_1, \dots, y_n]$  is a very generic polynomial then each operator  $d \in A_n^k$  satisfying  $\sigma_k(d) = P$  generates a maximal left ideal of  $A_n^k$ . Still under the assumption that  $k \geq 4$ , Lunts extends the above result to arbitrary  $n \geq 2$  in [7]. For  $k = 3$  and  $n \geq 2$  the very same statement has been proved by McCune [8]. All these results, in contrast with Stafford's, do not exhibit explicit examples of non-holonomic  $A_n$ -modules but instead prove that they are generic in the above sense. For an algorithm to produce explicit examples of the above form for  $n = 2$  and its implementation see [1].

In their other approach, Bernstein and Lunts look at the standard filtration of  $A_n$ . Now the  $i$ -th piece corresponds to differential operators of order  $\leq i$ . If  $\xi$  is a polynomial vector field,  $f$  a polynomial and  $I = \langle \xi + f \rangle$  then the characteristic variety of  $A_n/IA_n$  coincides the characteristic variety of the foliation  $\mathcal{F}_\xi$  as defined in the previous section. If  $\mathcal{F}_\xi$  has a quasi-minimal characteristic variety then according to [3, Proposition 6] there exists  $f \in \mathbb{C}[x_1, \dots, x_n]$  for which  $I = \langle \xi + f \rangle$  is maximal. While they do show that a generic  $\xi$  of degree  $\geq 2$  on  $\mathbb{C}^2$  has this property, they leave the general case as a conjecture, see [3, §4.2].

**Conjecture (Bernstein-Lunts).** *Let  $n \geq 2$  and  $\xi$  be a very generic polynomial vector field on  $\mathbb{C}^n$  with coefficients of degree  $\geq 2$ . Then  $\text{ch}(\mathcal{F}_\xi)$  is a quasi-minimal characteristic variety.*

The three dimensional case of the conjecture has been proved recently by Coutinho [5]. In this paper we will settle the general case.

**Theorem 2.** *Bernstein-Lunts conjecture holds true.*

Even when specialized to  $n = 3$ , our proof is very different from the one of Coutinho.

**1.3. Acknowledgements.** I am grateful to S. C. Coutinho for, since 2003, bringing periodically to my attention Bernstein-Lunts conjecture.

## 2. CHARACTERISTIC VARIETIES AND PROLONGATIONS

**2.1. Characteristic variety.** Let  $X$  be a quasi-projective manifold and  $\mathcal{F}$  be a foliation on  $X$  with cotangent bundle  $\mathcal{L}$ , that is  $\mathcal{F} = [\xi] \in \mathbb{P}H^0(X, TX \otimes \mathcal{L})$  with the representative  $\xi$  having no divisorial components in its singular set. As in the introduction set  $X_0 = X \setminus \text{sing}(\mathcal{F})$ .

Contraction with the twisted vector field  $\xi$  determines a morphism of  $\mathcal{O}_X$ -modules

$$T^*X \longrightarrow \mathcal{L}$$

whose kernel is  $N^*\mathcal{F}$ , the conormal sheaf of  $\mathcal{F}$ . At points  $x \in X_0$  the sheaf  $N^*\mathcal{F}$  is clearly locally free, but it is not locally free in general. For example it is never locally free at an isolated singularity of  $\mathcal{F}$  as one can promptly verify. Nevertheless, the restriction of  $N^*\mathcal{F}$  at  $X_0$  determines a subbundle of  $T^*X_0$  of corank one. As mentioned in the introduction  $\text{ch}(\mathcal{F})$  is defined as the closure in  $E(T^*X)$  of  $E(N^*\mathcal{F}|_{X_0})$ . We will use  $\pi$  to denote the natural projection  $\pi : E(T^*X) \rightarrow X$  as well as its restriction  $\pi : \text{ch}(\mathcal{F}) \rightarrow X$ .

If  $(x_1, \dots, x_n)$  are local coordinates at a open subset  $U \subset X$  then the vector fields  $\{\partial_{x_i} = \frac{\partial}{\partial x_i}\}$  can be thought as linear coordinates on  $T^*U$ : the value of  $\partial_{x_i}$  at a 1-form  $\omega \in T^*U$  is given by the contraction  $\omega(\partial_{x_i})$ . Thus, if we set  $y_i = \partial_{x_i}$

then  $(x_1, \dots, x_n, y_1, \dots, y_n)$  are global coordinate functions for  $T^*U$ . In particular, if  $\xi = \sum a_i \partial_{x_i}$  then

$$\text{ch}(\mathcal{F})|_{\pi^{-1}(U)} = \left\{ \sum a_i y_i = 0 \right\}.$$

The singular set of  $\text{ch}(\mathcal{F})$  is contained in  $\pi^{-1}(\text{sing}(\mathcal{F}))$  and contains  $\pi^{-1}(\text{sing}(\mathcal{F})) \cap X$ , where  $X$  sits inside  $E(T^*X)$  as the zero section. Thus, unless  $\mathcal{F}$  is a smooth foliation,  $\text{ch}(\mathcal{F})$  is always singular. It follows promptly from the above local expression of  $\text{ch}(\mathcal{F})$  that its singular points away from the zero section and over a fiber  $\pi^{-1}(p)$  are the 1-forms at  $T_p^*X$  which annihilates the image of  $D\xi(p)$ . Thus, if the singular scheme of  $\mathcal{F}$  is reduced and of dimension zero then  $\text{ch}(\mathcal{F})$  is smooth away from the zero section.

**2.2. Prolongation.** Recall that  $T^*X$  is endowed with a canonical symplectic structure which, in the above local coordinates, is induced by the 2-form

$$\Omega = \sum dx_i \wedge dy_i.$$

If  $F$  is a holomorphic function on (an open subset of)  $T^*X$  then the hamiltonian of  $F$  is by definition the vector field  $\xi_F$  determined by the formula

$$dF(\cdot) = \Omega(\xi_F, \cdot).$$

Notice that the vector field  $\xi_F$  is tangent to the hypersurface determined by  $F$  since  $\xi_F(F) = 0$ . Leibniz's rule implies that  $\xi_{uF} = u\xi_F + F\xi_u$ . Consequently the restriction of the direction field determined by  $\xi_F$  to  $\{F = 0\}$  is the same as the one of determined by  $\xi_{uF}$  for an arbitrary unit  $u$ . Therefore, the symplectic structure determines a one-dimensional foliation on any reduced and irreducible hypersurface  $H \subset T^*X$ : one has just to factor out eventual divisorial components of the singular set of  $\xi_F|_H$  to end up with a foliation on  $H$ , usually called in the literature the characteristic foliation of  $H$ . When  $H = \text{ch}(\mathcal{F}) \subset T^*X$ , is the characteristic variety of a foliation  $\mathcal{F}$  on  $X$  we will denote its characteristic foliation by  $\mathcal{F}^{(1)}$  and call it the first prolongation of  $\mathcal{F}$ .

If  $U \subset X$  is an open set with coordinates as in §2.1 and  $\xi = \sum a_i \partial_{x_i}$  is a vector field inducing  $\mathcal{F}$  on  $U$  then the vector field

$$(2.1) \quad \hat{\xi} = \sum_{i=1}^n a_i \partial_{x_i} - \sum_{i,j=1}^n (\partial_{x_j} a_i) y_i \partial_{y_j}$$

is the hamiltonian vector field of  $\sum a_i y_i$ , and hence defines the prolongation of  $\mathcal{F}|_U$ .

### 3. WARM-UP: PROOF OF THEOREM 1 IN DIMENSION THREE

In this section we present a proof of Theorem 1 in dimension three. We believe this will make the general case easier to understand.

**3.1. Making sense of the  $\mathcal{F}^{(1)}$ -invariance.** We start by clarifying the meaning of  $\mathcal{F}^{(1)}$ -invariance. The first result is well-known and holds in arbitrary dimension.

**Lemma 3.1.** *If  $Y \subset \text{ch}(\mathcal{F})$  is  $\mathcal{F}^{(1)}$ -invariant then  $\pi(Y)$  is  $\mathcal{F}$ -invariant.*

*Proof.* If  $p$  is a smooth point of  $\text{ch}(\mathcal{F})$  then equation (2.1) makes clear that  $\pi$  sends  $T_p \mathcal{F}^{(1)}$  into  $T_{\pi(p)} \mathcal{F}$ , and that the restriction of  $\mathcal{F}^{(1)}$  to the zero section is nothing more than  $\mathcal{F}$ . Together these two facts promptly imply the lemma.  $\square$

Our next result holds only in dimension three, and it is the lack of a direct analogue in higher dimensions which will make the proof in the general case more involved.

**Proposition 3.2.** *Suppose  $n = 3$  and let  $Y \subsetneq \text{ch}(\mathcal{F})$  be a homogenous, and irreducible subvariety with dominant projection to  $X$ . If  $Y$  is  $\mathcal{F}^{(1)}$ -invariant then  $\mathcal{F}$  is tangent to a codimension one web  $\mathcal{W}_Y$  on  $X$ .*

*Proof.* Since we are in dimension three, over the smooth locus of  $\mathcal{F}$ ,  $\text{ch}(\mathcal{F})$  is a rank two vector subbundle of  $\Omega_X^1$ . A subvariety  $Y$  as in the statement, determines  $k$  distinct lines on  $N^*\mathcal{F}_x$  for generic points  $x \in X$ . Therefore  $Y$  can be seen as the graph of a rational section  $\varpi$  of  $\text{Sym}^k \Omega_X^1$ . Moreover, the foliation  $\mathcal{F}$  is tangent to the multi-distribution determined by  $\varpi$ . Notice that so far, we have not used the  $\mathcal{F}^{(1)}$ -invariance of  $Y$ , we just explored the fact that  $Y$  is contained in  $\text{ch}(\mathcal{F})$ .

It remains to prove the integrability of the multi-distribution determined by  $\varpi$ . For that sake we can place ourselves at a neighborhood of a point  $x \in X$  where  $\varpi$  is holomorphic and equal to the product of  $k$  pairwise distinct 1-forms, say  $\omega_1, \dots, \omega_k$ , and  $\mathcal{F}$  is smooth. Choose a local coordinate system  $(x_1, \dots, x_n)$  where  $\mathcal{F}$  is induced by the vector field  $\xi = \partial_{x_1}$ . Hence  $\mathcal{F}^{(1)}$  is still induced by  $\partial_{x_1}$  now seen as a vector field on the total space of  $N^*\mathcal{F}$ .

If  $\omega$  is any of the 1-forms  $\{\omega_i\}_{i \in \underline{k}}$  then  $\omega = adx_2 + bdx_3$  for suitable holomorphic functions  $a, b$ . Notice that  $\omega$  is integrable if and only if the quotient  $a/b$  does not depend on  $x_1$ . Finally, the  $\mathcal{F}^{(1)}$ -invariance of  $Y$  ensures that  $a/b$  is constant along the orbits of  $\hat{\xi}$  and thus  $\omega$  is integrable and so is the multi-distribution induced by  $\varpi$ .  $\square$

### 3.2. Invariant subvarieties from singular points.

**Proposition 3.3.** *Let  $\mathcal{F}$  be a foliation on  $X$  a smooth projective variety of dimension three. Suppose  $\mathcal{F}$  is tangent to a codimension one web  $\mathcal{W}$ . If  $p \in \text{sing}(\mathcal{F})$  is an isolated singularity then there exists an irreducible  $\mathcal{F}$ -invariant subvariety  $Y \subsetneq X$  of positive dimension containing  $p$ .*

*Proof.* Suppose  $\mathcal{W}$  is a  $k$ -web with  $k \geq 1$ . If  $k \geq 2$ , let  $\Delta(\mathcal{W}) \subset X$  be the discriminant of the web  $\mathcal{W}$ . By definition,  $\Delta(\mathcal{W})$  is the set where  $\mathcal{W}$  is not the product of  $k$  pairwise transverse foliations. The proof of Proposition 3.2 tell us that on a neighborhood of a smooth point of  $\mathcal{F}$ , the web  $\mathcal{W}$  is induced by a  $k$ -symmetric 1-form  $\varphi = \sum a_{ij} dx_2^i dx_3^j$ . Thus  $\Delta(\mathcal{W})$  is defined as the hypersurface cut out by the discriminant of  $\varphi$ , seen as a the binary form on the variables  $dx_2, dx_3$ . Notice that  $\Delta(\mathcal{W})$  is a  $\mathcal{F}$ -invariant hypersurface.

If  $p$  belongs to  $\Delta(\mathcal{W})$  we are done. Otherwise  $\mathcal{W}$ , at a neighborhood of  $p$ , can be written as the superposition of  $k$  foliations, that is  $\mathcal{W} = \mathcal{G}_1 \boxtimes \dots \boxtimes \mathcal{G}_k$ . So consider one foliation  $\mathcal{G}$  of codimension one in a neighborhood of  $p$  and suppose that  $\mathcal{F}$  is tangent to it.

Let  $\xi$  be holomorphic vector field inducing  $\mathcal{F}$  and  $\omega$  be a holomorphic 1-form inducing  $\mathcal{G}$  both defined on a neighborhood of  $p$  and without divisorial components in their zero sets. Since  $\mathcal{F}$  has an isolated singularity at  $p$  so does  $\xi$ . Consequently,  $\omega(\xi) = 0$  implies that  $\omega$  is also singular at  $p$ . At this point we can use an argument laid down by Cerveau in [4, page 46] that we now recall. As  $\xi$  has isolated singularities we can apply De Rham-Saito Lemma to ensure the existence of another vector field  $\zeta$  such that  $\omega = i_{\xi} i_{\zeta} dx \wedge dy \wedge dz$ . Therefore the zero set of  $\omega$  is formed

by the minors of a  $3 \times 2$  matrix and must be of codimension at least two. But if  $\mathcal{G}$  is one of the foliations  $\mathcal{G}_i$  then  $\text{sing}(\mathcal{G})$  is algebraic, and is the sought  $\mathcal{F}$ -invariant variety.  $\square$

**3.3. Conclusion of the proof.** To conclude the proof of Theorem 1 in dimension three we will make use of the following generalization of Jouanolou's Theorem proved in [6], see also [7, Theorem 2] for the very same statement on projective spaces.

**Theorem 3.4.** *Let  $X$  be a smooth projective variety,  $\mathcal{L}$  be an ample line bundle over it, and  $k \gg 0$  be a sufficiently large integer. If  $\mathcal{F} \in \mathbb{P}H^0(X, TX \otimes \mathcal{L}^{\otimes k})$  is a very generic foliation then, besides  $X$  itself, the only subvarieties left invariant by  $\mathcal{F}$  are its singular points.*

Let  $\mathcal{F} \in \mathbb{P}H^0(X, TX \otimes \mathcal{L}^{\otimes k})$  be a very generic foliation without invariant subvarieties. As its singular set has cardinality given by the top Chern class of  $TX \otimes \mathcal{L}^{\otimes k}$ , and this number is positive for  $k \gg 0$ , the singular set of  $\mathcal{F}$  is non-empty. Moreover we can assume the existence of an isolated singularity  $p \in \text{sing}(\mathcal{F})$ , see for instance [6, Proposition 2.4].

If the characteristic variety of  $\mathcal{F}$  is not quasi-minimal then Proposition 3.2 implies that  $\mathcal{F}$  is tangent to a codimension one web  $\mathcal{W}$ . Proposition 3.3 in its turn implies that  $\mathcal{F}$  has a invariant subvariety through  $p$ . This contradicts Theorem 3.4 and concludes the proof of Theorem 1 in dimension three.  $\square$

**3.4. Obstructions to generalize.** To generalize the argument above to deal with the general case one has to circumvent the following obstructions:

- (1) Proposition 3.2 does not generalize because irreducible components of  $ch(\mathcal{F})$  which are homogenous and dominate the base  $X$  are no longer graphs of multi-distributions as happens in the three dimensional case; and
- (2) Proposition 3.3 does not generalize since (multi)-distributions with infinitesimal automorphisms are not necessarily integrable.

To accomplish that we will take advantage of the structure of generic foliation singularities combined with the following reinterpretation of Theorem 3.4.

**Theorem 3.5.** *Let  $X$  be a smooth projective variety,  $\mathcal{L}$  an ample line bundle over it, and  $k \gg 0$  a sufficiently large integer. If  $\mathcal{F} \in \mathbb{P}H^0(X, TX \otimes \mathcal{L}^{\otimes k})$  is a very generic foliation then every leaf of  $\mathcal{F}$  is Zariski dense.*

#### 4. PROLONGATION VERSUS HOLONOMY

In this section  $\mathcal{F}$  will be a **smooth** foliation of dimension one on a complex manifold  $X$ .

**4.1. Holonomy.** To each leaf  $L$  of  $\mathcal{F}$ , once a point  $p \in L$  and a germ  $(\Sigma, p)$  of smooth hypersurface transverse to  $\mathcal{F}$  are fixed, one can associate a (anti)-representation

$$\text{hol}(L) : \pi_1(L, p) \longrightarrow \text{Diff}(\Sigma, p),$$

as follows. Given a closed path  $\gamma$  contained in  $L$  and centered at  $p$  one defines a germ diffeomorphism  $h_\gamma \in \text{Diff}(\Sigma, p)$  such that  $h_\gamma(x)$  is the end point of a lift of  $\gamma$  to the leaf of  $\mathcal{F}$  through  $x$ . The result does not depend on the choices involved in the process and is completely determined by the class of  $\gamma$  in  $\pi_1(L, p)$ . Thus one set  $\text{hol}(L)(\gamma) = h_\gamma$ . It is an anti-representation since  $h_{\gamma_1 \cdot \gamma_2} = h_{\gamma_2} \circ h_{\gamma_1}$ .

Of course, one can also consider the linear holonomy of  $L$  which is just the anti-representation

$$\begin{aligned} Dhol(L) : \pi_1(L, p) &\longrightarrow GL(T_p\Sigma) \\ [\gamma] &\longmapsto Dh_\gamma(p). \end{aligned}$$

Being an anti-representation of  $\pi_1(L)$  onto a general linear group it is natural to wonder if there is a natural connection on a natural vector bundle over  $L$  inducing. It is indeed the case, and even better, there is a partial connection along the tangent bundle  $T\mathcal{F}$  of  $\mathcal{F}$  on the normal bundle  $N\mathcal{F}$  which has monodromy along the leaves of  $\mathcal{F}$  equivalent to the linear holonomy.

**4.2. Bott's partial connection.** Let  $\rho : TX \rightarrow N\mathcal{F}$  be the natural projection. Of course  $\ker\rho = T\mathcal{F}$ . Bott's partial connection is defined as follows

$$\begin{aligned} \nabla : T\mathcal{F} &\longrightarrow Hom(N\mathcal{F}, N\mathcal{F}) \simeq N^*\mathcal{F} \otimes N\mathcal{F} \\ \xi &\longmapsto \{\vartheta \mapsto \rho([\hat{\vartheta}, \xi])\}, \end{aligned}$$

where  $\hat{\vartheta}$  stands for an arbitrary lift of  $\vartheta$  to  $TX$ . The involutiveness of  $T\mathcal{F}$  implies that  $\rho([\hat{\vartheta}, \xi])$  does not depend on the choice of the lift, and ensures that  $\nabla$  is well defined.

Let us now proceed to write explicitly the restriction of  $\nabla$  to a leaf  $L$  of  $\mathcal{F}$ . We will work in local coordinates  $(x_1, x_2, \dots, x_n)$  and will assume that  $L = \{x_2 = \dots = x_n = 0\}$ . Since  $L$  is invariant by  $\mathcal{F}$ , we can write a vector field  $\xi$  generating  $T\mathcal{F}$  in the following form

$$\xi = a(x)\partial_{x_1} + \sum_{i=2}^n \sum_{j=2}^n a_{ij}(x)x_i\partial_{x_j}.$$

Notice that vector fields  $\partial_{x_2}, \dots, \partial_{x_n}$  can be interpreted as a basis of  $N\mathcal{F}$ . Thus

$$\nabla(\xi)(\partial_{x_i}) = \rho \left( \partial_{x_i} a(x)\partial_{x_1} + \sum_{i=2}^n \sum_{j=2}^n (\partial_{x_i} a_{ij}(x))x_i\partial_{x_j} + \sum_{j=2}^n a_{ij}(x)\partial_{x_j} \right).$$

Hence, the induced connection  $\nabla|_L : TL \rightarrow N^*L \otimes NL$  is

$$\begin{aligned} \nabla|_L(\xi) &= \sum_{i=1}^{n-1} \sum_{j=2}^n a_{ij}(x_1, 0) dx_i \otimes \partial_{x_j} \\ (4.1) \quad &= (dx_2, \dots, dx_n) \cdot A(x_1, 0) \cdot (\partial_{x_2}, \dots, \partial_{x_n})^T. \end{aligned}$$

**4.3. Comparison with the prolongation.** In order to compare with Bott's connection, let us now write down the restriction to  $\pi^{-1}(L)$  of the lift of  $\xi$  to  $E(N^*\mathcal{F})$ . We will use the same system of coordinates used in Section 2.1, where  $y_i = \partial_{x_i}$ . Since in these coordinates  $\pi^{-1}(L) = \{y_1 = x_2 = \dots = x_n = 0\}$ , we can write

$$\hat{\xi}|_{\pi^{-1}(L)} = a(x_1, 0)\partial_{x_1} - \sum_{i,j=2}^n (a_{ji}(x_1, 0)) y_i \partial_{y_j}$$

which in matrix form is

$$\hat{\xi}|_{\pi^{-1}(L)} = a(x_1, 0)\partial_{x_1} - (y_2, \dots, y_n) \cdot A^T(x_1, 0) \cdot (\partial_{y_2}, \dots, \partial_{y_n})^T$$

with  $A(x_1, 0)$  being the same matrix as in (4.1). It is then clear, that in these coordinates, the leaves of  $\mathcal{F}^{(1)}$  restricted to  $\pi^{-1}(L)$  are flat sections of the connection

on  $N^*L$  having connection matrix  $-A^T$ , where  $A$  is the connection matrix on  $\nabla|_L$ . We have thus prove the following

**Proposition 4.1.** *The leaves of  $\mathcal{F}^{(1)}$  are flat sections of the partial connection dual to Bott's partial connection.*

## 5. FROM INVARIANT SUBVARIETIES TO MULTI-DISTRIBUTIONS

**5.1. Non-resonant singularities.** Let  $\mathcal{F}$  be a germ of one-dimensional foliation on  $(\mathbb{C}^n, 0)$ . Suppose that it has an isolated singularity at the origin. Suppose also that the linear part  $D\xi(0)$  of a vector field  $\xi$  inducing  $\mathcal{F}$  is invertible and its eigenvalues  $\lambda_1, \dots, \lambda_n$  generate a  $\mathbb{Z}$ -module of rank  $n$ . We will say that a singularity of this form is a **non-resonant singularity**.

**Lemma 5.1.** *There exists  $n$  germs of  $\mathcal{F}$ -invariant smooth curves  $\gamma_i : (\mathbb{C}, 0) \rightarrow (\mathbb{C}^n, 0)$  with tangents at zero determined by the eigenvectors of  $D\xi(0)$ .*

*Proof.* The Hadamard-Perron theorem for holomorphic flows [11, Chapter 2, Section 7] ensures the existence of a pair of invariant manifolds intersecting transversely at the origin and such that the restriction of the vector field to each of them has a non-resonant singularity in the Poincaré domain (Section 5 loc. cit.). Poincaré normalization Theorem (loc. cit.) implies that the corresponding restrictions of  $\xi$  are analytically linearizable. Since separatrices of the restrictions of  $\xi$  are also separatrices of  $\xi$ , the lemma follows.  $\square$

The linear holonomy along a positive oriented path around the origin contained in  $\gamma_i(\mathbb{C}, 0)$  is induced by a linearizable matrix  $A_i \in GL(n-1, \mathbb{C})$  with eigenvalues  $\{\exp(2\pi i \lambda_j / \lambda_i)\}_{j \neq i}$ . Moreover, the  $\mathbb{Z}$ -independence of the eigenvalues implies that the Zariski closure of the subgroup of  $GL(\mathbb{C}^{n-1})$  generated by  $A_i$  is a maximal torus  $\simeq (\mathbb{C}^*)^{n-1}$ .

**5.2. Singularities and the holonomy of separatrices.** Together with Proposition 6.2, the proposition below will replace Proposition 3.2 in the proof of the general case of Theorem 1. It guarantees that invariant subvarieties of  $\mathcal{F}^{(1)}$  correspond to multi-distributions tangent to  $\mathcal{F}$  as soon as  $\mathcal{F}$  has non-resonant singularities.

**Proposition 5.2.** *Let  $\mathcal{F}$  be a foliation on a smooth projective variety  $X$  and let  $Y \subsetneq \text{ch}(\mathcal{F})$  be an irreducible subvariety with dominant projection to  $X$  distinct from the zero section. Suppose  $\mathcal{F}$  has non-resonant singularity  $p$  and that at least one of its separatrices is Zariski dense. If  $Y$  is  $\mathcal{F}^{(1)}$  invariant then the fiber of  $Y$  over a generic point of  $X$  is a finite union of linear spaces of the same dimension. Consequently  $\mathcal{F}$  is tangent to a multi-distribution of codimension  $q = \dim Y - \dim X \leq \dim X - 2$ .*

*Proof.* First consider a point  $p_0 \in X$  in the Zariski dense separatrix through  $p$ , and let  $L$  be the leaf of  $\mathcal{F}$  through it. The fiber  $V$  of  $E(N^*\mathcal{F}) \simeq \text{ch}(\mathcal{F}) \rightarrow X$  over  $p$  is a vector space of dimension  $n-1$ . The intersection of  $Y$  with  $V$  is a subvariety of  $V$  invariant by the subgroup  $G \subset GL(V)$  image of the representation  $\pi_1(L) \rightarrow GL(V)$  dual to the linear holonomy of  $L$ . Since  $V \cap Y$  is algebraic, not only  $G$  but also its Zariski closure leaves  $V \cap Y$  invariant. By hypothesis,  $\overline{G} \simeq (\mathbb{C}^*)^{n-1}$  is a maximal torus in  $GL(V)$ . Consequently  $V \cap Y$  is a finite union of linear spaces for an arbitrary  $p \in L$ . To be a finite union of linear subspaces is clearly a Zariski closed condition. Thus the same will hold true for the fibers of  $Y$  over points in the Zariski closure of  $L$  which is, by assumption, equal to  $X$ .  $\square$



## 6. FROM MULTI-DISTRIBUTIONS TO INVARIANT SUBVARIETIES

We now proceed to establish the result which will replace Proposition 3.3. We start with a simple lemma.

**Lemma 6.1.** *Let  $\omega \in \Omega^q = \Omega^q(\mathbb{C}^n) \otimes \mathbb{C}[[x_1, \dots, x_n]]$  be a formal  $q$ -form. If  $\omega$  is invariant by the natural  $(\mathbb{C}^*)^n$ -action on  $\mathbb{C}^n$  then*

$$(6.1) \quad \omega = f \cdot \left( \sum_{I \in \{1, \dots, n\}^q} \lambda_I \frac{dx_I}{x_I} \right)$$

where  $f \in \mathbb{C}[[x_1, \dots, x_n]]$ ,  $\lambda_I \in \mathbb{C}$  and  $\frac{dx_I}{x_I} = \frac{dx_{i_1}}{x_{i_1}} \wedge \dots \wedge \frac{dx_{i_q}}{x_{i_q}}$ .

*Proof.* Write  $\omega = \sum_{i=i_0}^{\infty} \omega_i$ , where the coefficients of  $\omega_i$  are polynomials of degree  $i$  and  $\omega_{i_0} \neq 0$ . If  $\varphi_t(x) = t \cdot x$  then

$$\frac{(\varphi_t)^* \omega}{t^{i_0+q}} = \omega_{i_0} + \sum_{i=i_0+1}^{\infty} t^{i-t_0+q} \omega_i.$$

Since for arbitrary  $t$ ,  $\varphi_t^* \omega$  must be a multiple of  $\omega$  then after dividing by a suitable formal function we can assume that  $\omega$  is homogeneous.

Let  $x^J dx_I$  be a monomial appearing in  $\omega$ . Suppose  $x_1^{j_1}$  divides  $x^J$  but  $x_1^{j_1+1}$  does not. Consider the automorphism  $\varphi_t(x_1, x_2, \dots, x_n) = (tx_1, x_2, \dots, x_n)$ . Then  $\varphi_t^*(x^J dx_I) = t^{j_1+\epsilon} x^J dx_I$ , where  $\epsilon = 0$  if  $dx_1$  does not appear in  $dx_I$  and  $\epsilon = 1$  otherwise. If  $j_1 + \epsilon \geq 2$  then  $x_1$  divides all the other monomials appearing in  $\omega$ . Does after division we can assume  $j_1 + \epsilon = 1$  and the same will hold true for any other monomial appearing in  $\omega$ . Repeating the argument for the other coordinate functions makes clear the assertion of the lemma.  $\square$

**Proposition 6.2.** *Let  $\xi$  be a germ of holomorphic vector field on  $(\mathbb{C}^n, 0)$  with a non-resonant singularity at the origin. Suppose  $\xi$  is an infinitesimal automorphism of a distribution  $\mathcal{D}$  of codimension  $q \leq n-2$ . Then  $\mathcal{D}$  is integrable and the singular set of  $\mathcal{D}$  has positive dimension.*

*Proof.* Let  $\omega$  be a germ of holomorphic  $q$ -form,  $q = n - p$ , defining  $\mathcal{D}$ , that is  $\mathcal{D} = \{v \in T(\mathbb{C}^n, 0) \mid \omega(v) = 0\}$ . For further use let us recall that a  $q$ -form  $\omega$  defines a codimension  $q$  distribution if and only if

$$(i_v \omega) \wedge \omega = 0 \quad \text{for every } v \in \bigwedge^{q-1} \mathbb{C}^n,$$

and this distribution is integrable if and only if

$$(i_v \omega) \wedge \omega = (i_v \omega) \wedge d\omega = 0 \quad \text{for every } v \in \bigwedge^{q-1} \mathbb{C}^n,$$

see [9]. It follows that integrability is a formal condition, and as such can be verified in an arbitrary formal coordinate system.

Since the origin is a non-resonant singularity for  $\xi$ , we can choose formal coordinates such that

$$\xi = \sum_{i=1}^n \lambda_i x_i \partial_{x_i}$$

where  $\lambda_i \in \mathbb{C}$  are complex numbers. In exchange we can no longer assume that  $\omega$  is a holomorphic  $q$ -form, but it is certainly a formal  $q$ -form.

Since  $\xi$  is an infinitesimal automorphism of  $\mathcal{D}$ , its flow  $\varphi_t : (\mathbb{C}^n, 0) \rightarrow (\mathbb{C}^n, 0)$  preserves  $\omega$ . More precisely,

$$\varphi_t^* \omega = f(t, x) \omega$$

for a suitable formal function  $f \in \mathbb{C}[[t, x_1, \dots, x_n]]$ .

Consider now the subgroup  $G \subset (\mathbb{C}^*)^n \subset GL(\mathbb{C}^n)$  defined as

$$G = \{A \in (\mathbb{C}^*)^n \mid A^* \omega \wedge \omega = 0 \text{ in } \bigwedge^2 \Omega^q \otimes \mathbb{C}[[x_1, \dots, x_n]]\},$$

where  $(\mathbb{C}^*)^n$  acts on  $(\mathbb{C}^n, 0)$  through a diagonal linear map. The flow of  $\xi$  determines a non-closed one parameter subgroup of  $H \subset G$ . Since  $G$  is clearly an algebraic subgroup, it follows that the Zariski closure of  $H$  is also contained in  $G$ . But the dimension of the Zariski closure of  $H$  is nothing more than the rank of the  $\mathbb{Z}$ -module generated by  $\lambda_1, \dots, \lambda_n$ . It follows that  $\bar{H} = G = (\mathbb{C}^*)^n$ .

On the one hand, since  $\omega$  induces a distribution  $\iota_v \omega \wedge \omega = 0$ . On the other hand, Lemma 6.1 implies that  $\omega$  is a multiple of a closed  $q$ -form, and consequently  $\iota_v \omega \wedge d\omega = 0$ . This shows that  $\mathcal{D}$  is integrable.

It remains to prove that the singular set of  $\mathcal{D}$  has positive dimension. Looking at the expression (6.1) we realize that it must have at least two non-trivial summands. Indeed, if not,  $\mathcal{D}$  would be a smooth foliation tangent to  $\xi$ , what is clearly impossible. Therefore, if  $k$  is the cardinality of the set  $\bar{I} = \cup_{\lambda_I \neq 0} I$ , where the complex numbers  $\lambda_I$  are defined by (6.1), then  $k > q$ . Clearly, the coordinate hyperplanes with index in  $\bar{I}$  is invariant by  $\mathcal{D}$ . Consequently the intersection of any  $q + 1$  of these coordinate hyperplanes is also invariant by  $\mathcal{D}$ . Since  $\mathcal{D}$  has codimension  $q$ , this intersection must be contained in the singular locus of  $\mathcal{D}$ .  $\square$

**Remark 6.3.** Proposition 6.2 will be in the proof of the general case of Theorem 1 what Proposition 3.2 is in the proof of the three-dimensional case. The analogy is not perfect as we do not proved here the integrability of multi-distributions as we did there. Anyway, with some extra effort one can also prove the integrability of the multi-distribution. We will not pursue this here as the result above is sufficient for our purposes.

## 7. PROOF OF THEOREM 1

Let  $\mathcal{F} \in \mathbb{P}H^0(X, TX \otimes \mathcal{L}^{\otimes k})$  be a very generic foliation. We can assume, thanks to Theorem 3.5, that  $\mathcal{F}$  has isolated singularities, at least one non-resonant singularity, and every leaf of  $\mathcal{F}$  is Zariski dense.

Proposition 5.2 implies that  $\mathcal{F}$  is tangent to a multi-distribution  $\mathcal{D}$ . We can assume  $\mathcal{D}$  is irreducible without loss of generality.

If  $\mathcal{D}$  is locally decomposable around  $p$  then Proposition 6.2 implies the existence of a positive dimensional irreducible component  $Z$  of the singular set of  $\mathcal{D}$  through  $p$ . This set is clearly algebraic and invariant by  $\mathcal{F}$  since sections of  $T\mathcal{F}$  are infinitesimal automorphisms of  $\mathcal{D}$ . If  $\mathcal{D}$  is not locally decomposable at  $p$  then there exists a subvariety  $Z \subsetneq X$  where  $\mathcal{D}$  is not locally decomposable. As above, we conclude that  $Z$  is invariant by  $\mathcal{F}$ .

In both cases, we arrive at a contradiction with Theorem 3.5.  $\square$

## 8. BERNSTEIN-LUNTS CONJECTURE

Theorem 1 implies the existence of foliations, on arbitrary projective varieties, with quasi-minimal characteristic variety. Moreover, as the conclusion of Theorem 3.5 holds true for any foliation with ample cotangent bundle on  $\mathbb{P}^n$ , the existential part of Bernstein-Lunts Conjecture is settled. Nevertheless, there is still a detail to be dealt with in order to prove that a *very generic* polynomial vector field of degree  $d \geq 2$  has quasi-minimal characteristic variety.

**8.1. Projective versus affine degree.** The (projective) degree of a holomorphic foliation  $\mathcal{F}$  on  $\mathbb{P}^n$  is defined as the degree of the tangency divisor of  $\mathcal{F}$  with a generic hyperplane  $H$ . If  $\mathcal{F} \in \mathbb{P}H^0(\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(k))$  then the degree of  $\mathcal{F}$  is equal to  $k + 1$ .

If one starts with a polynomial vector field  $\xi$  of degree  $d$  on  $\mathbb{C}^n$  then it is natural to extend it to holomorphic foliation  $\mathcal{F}_\xi$  on  $\mathbb{P}^n$  such that  $H$  is not contained in the singular set of  $\mathcal{F}_\xi$ . We set the degree of  $\xi = \sum a_i \partial_i$  as the maximal degree of its coefficients  $a_i$ . In general the (projective) degree of  $\mathcal{F}_\xi$  is at most the (affine) degree of  $\xi$ . Moreover precisely,

$$\deg(\mathcal{F}_\xi) = \begin{cases} \deg(\xi) & \text{if } H \text{ is invariant by } \mathcal{F}_\xi, \\ \deg(\xi) - 1 & \text{if } H \text{ is not invariant by } \mathcal{F}_\xi. \end{cases}$$

If  $\mathcal{D}(n, d)$  is the set of polynomial vector fields of degree at most  $d$  then the generic element in it extends to a foliation of  $\mathbb{P}^n$  with singularities of codimension at least two which leaves the hyperplane at infinity invariant, see [12] for a through discussion. In more intrinsic terms, if  $T\mathbb{P}^n(-\log H)$  denotes the subsheaf of  $T\mathbb{P}^n$  constituted by germs of vector fields tangent to  $H$  then  $\mathcal{D}(n, d)$  can be identified with  $H^0(\mathbb{P}^n, T\mathbb{P}^n(-\log H) \otimes \mathcal{O}_{\mathbb{P}^n}(d-1))$ . Under this identification the extension which do not leave the hyperplane at infinity invariant will appear with a divisorial component in their singular set supported there.

**8.2. Relative version of Theorem 3.5.** The proof of Theorem 3.5 can be adapted to prove the following

**Theorem 8.1.** *Let  $X$  be a smooth projective variety and  $H \subset X$  a smooth hypersurface. Let also  $\mathcal{L}$  be an ample line bundle over  $X$ , and  $k \gg 0$  a sufficiently large integer. If  $\mathcal{F} \in \mathbb{P}H^0(X, TX(-\log H) \otimes \mathcal{L}^{\otimes k})$  is a very generic foliation then every leaf of  $\mathcal{F}$  not contained in  $H$  is Zariski dense.*

We will not detail its proof as the case of projective spaces (the one used in the proof of Theorem 2 below) is Theorem 4.2 of [5]. Moreover, there it is proved that it suffices to take  $k \geq 1$  when  $X = \mathbb{P}^n$  and  $\mathcal{L} = \mathcal{O}_{\mathbb{P}^n}(1)$ .

**8.3. Proof of Theorem 2.** According to Theorem 8.1 the leaves of a very generic vector field of degree  $d \geq 2$  are Zariski dense. Also a very generic vector field has at least one non-resonant singularity. Thus we can apply Propositions 5.2 and 6.2 to conclude that the characteristic variety of  $\mathcal{F}_\xi$  is quasi-minimal.  $\square$

## REFERENCES

- [1] L. C. O. ALMEIDA AND S. C. COUTINHO, *On homogenous minimal involutive varieties*. LMS J. Comput. Math. **8** (2005), 301–315
- [2] J. BERNSTEIN, *Analytic continuation of generalized functions with respect to a parameter*. Funkcional. Anal. i Priloen. **6** (1972), no. 4, 26–40.
- [3] J. BERNSTEIN AND V. LUNTS, *On nonholonomic irreducible  $D$ -modules*. Invent. Math. **94** (1988), no. 2, 223–243.

- [4] D. CERVEAU, *Feuilletages holomorphes de codimension 1. Réduction des singularités en petites dimensions et applications*. Dynamique et géométrie complexes, 11–47, Panor. Synthèses, **8**, Soc. Math. France, Paris, 1999.
- [5] S. C. COUTINHO, *Foliations of multiprojective spaces and a conjecture of Bernstein and Lunts*. To appear in Trans. of the AMS.
- [6] S. C. COUTINHO AND J. V. PEREIRA, *On the density of algebraic foliations without algebraic invariant sets*. J. Reine Angew. Math. **594** (2006), 117–135.
- [7] V. LUNTS, *Algebraic varieties preserved by generic flows*. Duke Math. J. **58** (1989), no. 3, 531–554.
- [8] T. C. MCCUNE, *A generic cubic surface contains no involutive curve*. J. Algebra **235** (2001), no. 1, 36–44.
- [9] A. S. DE MEDEIROS, *Singular foliations and differential p-forms*. Ann. Fac. Sci. Toulouse Math. (6) **9** (2000), no. 3, 451–466.
- [10] J. T. STAFFORD, *Nonholonomic modules over Weyl algebras and enveloping algebras*. Invent. Math. **79** (1985), no. 3, 619–638.
- [11] Y. ILYASHENKO AND S. YAKOVENKO, *Lectures on analytic differential equations*. Graduate Studies in Mathematics, **86**. Amer. Math. Soc., Providence, RI, 2008. xiv+625 pp.
- [12] S. ZAKERI, *Dynamics of singular holomorphic foliations on the complex projective plane*. Laminations and foliations in dynamics, geometry and topology, 179–233, Contemp. Math., **269**, Amer. Math. Soc., Providence, RI, 2001.

JORGE VITÓRIO PEREIRA, IMPA, ESTRADA DONA CASTORINA, 110, 22460-320, RIO DE JANEIRO, RJ, BRAZIL

*E-mail address:* jvp@impa.br