HOMOCLINIC BIFURCATIONS AND UNIFORM HYPERBOLICITY FOR THREE-DIMENSIONAL FLOWS

AUBIN ARROYO

ABSTRACT. In this paper we prove that any C^1 vector field defined on a three dimensional manifold can be approximated by one that is uniformly hyperbolic, or that exhibits either a homoclinic tangency or a singular cycle. This proves a statement of a conjecture of Palis in the context of flows in the C^1 topology. For that, we rely on the notion of dominated splitting for the associated linear Poincaré flow.

1. INTRODUCTION

The quest to understand the asymptotic behavior of orbits of a large set of evolution laws has been guiding one important strain of the theory of dynamical systems. In the sixties, the notion of structural stable systems, first introduced by Andronov and Pontryagin thirty years before, meaning systems that are equivalent up to a continous global change of coordinates to all nearby ones, was at the center of the attempts to provide a typical model inside the universe of dynamical systems. In that decade and in the early seventies, in a series of papers by Anosov, Palis and Smale, Robbin, de Melo and Robinson, they proved that uniform hyperbolicity and transversality of stable and unstable manifolds, imply stucutral stability. In the mid-eighties, Mañé obtained the converse for diffeomorphisms in a profund and very original work. Ten years later, Hayashi extended this result for flows, ending a remarkable page in the history of modern dynamics: The solution of the Stability Conjecture, [16].

Recall that an invariant set Λ of a flow X is hyperbolic if its tangent bundle splits into three DX-invariant sub-bundles, E^s, E^u and $[X]$, where vectors on E^s are uniformly contracted in the future, the same for E^u in the past, and $[X]$ is defined by the flow direction. A flow X is uniformly hyperbolic (or Axiom A) if its limit set is hyperbolic. For the case of diffeomorphisms this concept is similar by considering only the first two sub-bundles. For a complete description of uniform hyperbolicity we refeer the reader to [17] and [24]. However, at the

second half of the sixties it was already clear that uniform hyperbolicity could not be present for every system of a dense subset in the universe of all dynamics.

A key question was then: Is it possible to look for a general scenario for dynamics? The search for such an answer, in particular the work of Newhouse on the coexistence of infinitely many sinks (see [10]), draw the attention to homoclinic orbits. That is, orbits that in the past and the future converge to the same periodic orbit, first considered by Poincaré almost a century before. Altogether, the examples by Newhouse gave rise to a rich family of new dynamics. The creation-destruction of such orbits is, roughly speaking, what its meant by homoclinic bifurcations; see [17] for a formal definition in any dimensions.

Based on these and other subsequent developments, Palis formulated in the mid-nineties the following conjecture (see [17] and [14]):

Conjecture 1. *The diffeomorphisms exhibiting a homoclinic bifurcation are C^r dense in the complement of the closure of the hyperbolic ones* ($r \geqslant 1$).

There has been substantial advances in this direction. On [21], Pujals and Sambarino, provide a proof of this conjecture in the case of diffeomorphisms of compact surfaces, in the C^1 topology. In this setting, homoclinic bifurcations only arise from homoclinic tangencies. Recall that the stable and unstable manifolds of a hyperbolic periodic point p, $W^s(p)$ and $W^u(p)$, respectively, are inmersed submanifolds on the ambient space. A point $q \in W^s(p) \cap W^u(p)$ is a homoclinic point if $q \neq p$. If these submanifols meet tangentialy at a homoclinic point q, this is called a homoclinic tangency on q.

Then, it is natural to consider this conjecture in the context of flows; however, another homoclinic phenomenon involving singularities of the vector field must be considered: singular cycles. Recall that a singular point σ of a vector field X is hyperbolic if the derivative $D_\sigma X$ has no eigenvalues which are purely imaginary, and in this case we have stable and unstable manifolds of σ, denoted by $W^s(\sigma)$, and $W^u(\sigma)$, respectively. It may happen that $q \in W^s(\sigma) \cap W^u(\sigma)$ for $q \neq \sigma$. In this case we say the vector field has a *singular cycle*, a concept that was first considered by Labarca and Pacífico [7]. In dimension three, these two phenomena characterize all possible homoclinic biffurcations.

This work is intended to give a positive answer to a version of the conjecture above, in the context of three-dimensional vector fields and for the C^1-topology. To be more precise, let M be a compact manifold of dimension three without boundary. Denote by $\mathcal{X}^r(M)$ the space of vector fields of class C^r ($r \geqslant 1$) defined on M.

Theorem A. *Any vector field $X \in \mathcal{X}^1(M)$ can be approximated by another one $Y \in \mathcal{X}^1(M)$ showing one of the following phenomena:*

1. *Uniform Hyperbolicity with the no-cylces condition.*
2. *A homoclinic tangency.*
3. *A singular cycle.*

As in the case of surface diffeomorphisms, our methods rely on a relaxed form of hyperbolicity, first used by Mañé, Liao and Pliss in their attempts to characterize structural stable diffeomorphisms; it is called *dominated splitting* and it strictly includes the class partially hyperbolic systems (see [6], [1] and [25]). An invariant set Λ for a diffeomorphism f has a dominated splitting if its tangent bundle decomposes into two Df-invariant subbundles $T_\Lambda M = E \oplus F$ and there are constants $C > 0$ and $\lambda \in (0, 1)$ such that for all $x \in \Lambda$:

$$||Df^n|_{E(x)}||||Df^{-n}|_{F(f^n(x))}|| < C\lambda^n;\ \forall n \geqslant 0$$

There is a remarkable relationship between the absence of homoclinic tangencies and the existence of a dominated splitting. This fact is explored in [21] and more recently, the same authors provided a rather complete description of the dynamics underlying a dominated splitting for surface diffeomorphisms, see [22].

Still, the natural translation of this notion into the three-dimensional flows is more limited. That is, splittings of the kind: $T_\Lambda M = E \oplus [X] \oplus F$, where each sub-bundle is one-dimensional, or $T_\Lambda M = E \oplus F$, where one is one-dimensional, both exlude the scenarios depicted in the following example:

Consider an Anosov flow A on M and choose an infinite sequence of periodic orbits with periods arbitrarily large, say $\{p_n\}$. An application of Frank's Lemma (see [9] for a flow version) allows us to perturb (C^1) the flow on each p_n by pushing the strong-stable direction over the tangent direction of the orbit of p_n within the central stable space defined by $E^s + [X]$. In the same way, in another part of the orbit of p_n, we push the strong-unstable direction. Inductively, this process renders a sequence X_n of vector fields. Observe that no limit of $\{X_n\}$ can have an invariant splitting of any of the kind as above. Also it is not possible to find a nearby flow exhibiting a homoclinic tangency, since strong directions remain orthogonal to each other on each point.

Like hyperbolicity, the property of a dominated splitting is a condition on the action of the derivative on the tangent bundle which determines some dynamical behavior of the transformation itself. In our case, we shall consider a condition not on the whole tangent bundle but on its projection over a certain subbundle everywhere normal to

the flow direction; that is, the linear Poincaré flow associated. We shall be precise about this in the next section.

The notion of dominated splitting along the linear Poincaré flow allows us to obtain the following fact: Once we are far from homoclinic tangencies, we get equipped with a dominated splitting (see Proposition 3.2 for a precise statement).

On the other hand, Theorem A relies on a fundamental result related to the possible underlying dynamics of an invariant set provided with a dominated splitting. To explain this, let us call an embedded two-dimensional torus $\mathcal{T} \subset M$ an *irrational torus* if $\mathcal{T}$ is invariant by the action of X^t, it is normally hyperbolic and moreover, $X^t|_{\mathcal{T}}$ is conjugated to a linear irrational flow on $S^1 \times S^1$. Denote by $\mathrm{Per}(X)$ the set of periodic orbits of X, and $\mathrm{Sing}(X) = \{p \in M | X(p) = 0\}$, for a given $X \in \mathcal{X}^r(M)$, M being a closed three-manifold.

Theorem B. *Let Λ be a compact invariant set for $X \in \mathcal{X}^2(M)$ with a dominated splitting and such that $Sing(X) \cap \Lambda = \emptyset$ and $Per(X) \cap \Lambda$ are all hyperbolic saddles, then $\Lambda = \tilde{\Lambda} \cup \mathcal{T}$, where $\tilde{\Lambda}$ is hyperbolic and $\mathcal{T}$ is a finite union of irrational tori.*

The dominated splitting provides us with invariant foliations over the set Λ, formed by central stable and central unstable manifolds (See Lemma 4.2). *A priori*, such leaves do not have a dynamical meaning at all. However, a kind of Denjoy's property asserts that smoothness of the vector field does not allow the existence of wandering intervals; moreover, it provides us the asymptotic behavior of local central manifolds. This allows us to construct actual stable and unstable manifolds from the central ones. (See Lemma 5.2 and Lemma 5.14). For that, we must establish a way to control how fast orbits get farther apart as we iterate; not as two points at certain flow-time but as orbits of the linear Poincaré flow. In order to do that, it ougth to be necessary to consider a set of holonomy maps defined on transversal two-dimensional sections. These maps together play the role of a kind of global Poincaré transformation and allow us to proceed, at least locally, as in the case of a surface-diffeomorphism (see Section 2.2). Once we can use dynamically defined invariant manifolds, some modifications of the arguments of [21] are needed to conclude the proof of Theorem B. Indeed, [21] is a key reference thoughout this paper.

Further questions: Lorenz-like attractors (see [3]) are another kind of rich dynamics that can appear from singular cycles. They are characterized by the presence of infinitely many periodic orbits in any neighborhood of a singularity, in a robust way. The study of this situation

lead Morales, Pacífico and Pujals to define an adequate notion of hyperbolicity, where singularities are involved, called singular hyperbolicity (see [9]). Our methods together with the notions of [9] may lead to a stronger statement of Theorem A, remplacing Singular Cycle by Singular Hyperbolic, on item 3.

However, some work has to be done, in particular on the choice of the dominated splitting over an invariant set which contains a singularity, since the linear Poincaré flow it is not defined on equilibrium points. Nevertheless, it seems to be useful to consider also a splitting on the tangent bundle as in the way is done for singular hyperbolic sets.

Finally, this is a joint work with Federico Rodriguez Hertz.

Aknowledgements: The author is very thankful to Jacob Palis for his encouragement and guideance, to Enrique Pujals for many enlighting conversations and CNPq-Brazil and Conacyt-México for the financial support.

2. Dominated Splitting for Invariant Sets

This section is devoted to construct a useful notion of dominated splitting for invariant sets of flows related to the linear Poincaré flow associated, acting on a sub-bundle of the tangent space. At the end of this section, we will obtain a family of holonomy maps defined between pairs of two-dimensional sections, each one transversal to the flow. This family of maps acts on the manifold in a similar way the linear Poincaré flow acts on the normal bundle. On what follows M shall denote always a three-dimensional compact riemannian manifold without boundary. Given vector field $X \in \mathcal{X}^1(M)$ there is associated a family of diffeomorphisms on M, say, $\{X^t\}$ such that $X^s \circ X^t = X^{s+t}$ for all $s, t \in \mathbb{R}$, $X^0 = \mathrm{Id}$ and $dX^t(p)/dt|_{t=0} = X(p)$, for any $p \in M$. This is the flow associated to the vector field X.

2.1. **The Linear Poincaré Flow.** The vector field X induces two sub-bundles on $T_Q M$, say, $[X]_p = \{v \in T_pM | v = \alpha X(p), \alpha \in \mathbb{R}\}$ and $\tilde{\mathcal{N}_p} = [X]_p^{\perp}$, for any $p \notin \mathrm{Sing}(X)$. Denote by $\mathcal{N} = \frac{T_Q M}{[X]}$ the normal bundle of X over the regular points $Q = M \setminus \mathrm{Sing}(X)$. For each $t \in \mathbb{R}$ the tangent map of X^t restricted to $\mathcal{N} \subset T_Q M$ induces an automorphism on $\mathcal{N}$ that conmutes the following diagram:

$$\begin{array}{ccc} \mathcal{N} & \xrightarrow{L^t} & \mathcal{N} \\ \downarrow & & \downarrow \\ Q & \xrightarrow{X^t} & Q \end{array}$$

The flow of automorphisms $\{L^t\}_{t\in\mathbb{R}}$ is called the *linear Poincaré flow* of the normal bundle $\mathcal{N}$. We took this notion from the work of Doering, [2], and it allows us to define an adequate notion of *dominated splitting* for invariant sets of flows.

Definition 1. An invariant set Λ ($X^t(\Lambda)\subset\Lambda$, $\forall t\in\mathbb{R}$) has a dominated splitting if there exists a splitting of the normal bundle of $\tilde{\Lambda}=\Lambda\setminus \mathrm{Sing}(X)$, $\mathcal{N}_{\tilde{\Lambda}}=E\oplus F$ and there are numbers $C>0$ and $\lambda\in(0,1)$ such that: $L^t(E(x))=E(X^t(x))$, $L^t(F(x))=F(X^t(x))$ and

$$|L^t|_{E(x)}||L^{-t}|_{F(X^t(x))}|\leqslant C\lambda^t;\ \forall t\geqslant 0$$

Fix an invariant set Λ for X^t without singularities and take a splitting of $\mathcal{N}_\Lambda=E'\oplus F'$ not necessarily invariant. For any number $a\in(0,1]$ we can define the following two cone fields pointwise. On each $x\in\Lambda$ set:

$$C_a^E(x)=\{w\in\mathcal{N}_x | w=v_E+v_F,\ |v_F|\leqslant a|v_E|\}$$
$$C_a^F(x)=\{w\in\mathcal{N}_x | w=v_E+v_F,\ |v_E|\leqslant a|v_F|\}$$

As in the hyperbolic case, it is not difficult to see that if this cone field is invariant under the action of L^t we obtain a dominated splitting on Λ (see [12]). The following lemma precisely states this.

Lemma 2.1. *If there exist $\lambda<1$, $a>0$, and there is a splitting of $\mathcal{N}_\Lambda=E'\oplus F'$ such that*

$$L^t(C_a^{F'})\subset C^{F'}_{\lambda^t a}(X^t(x)) \text{ and } L^{-t}(C_a^{E'})\subset C^{E'}_{\lambda^t a}(X^{-t}(x))$$

for all $t\geqslant 0$, then Λ has a dominated splitting.

Moreover, the following Lemma due to Doering (see Proposition 1.1 in [2]), asserts that in the case of hyperbolicity, these two notions are coherent.

Lemma 2.2. *Let $\Lambda\subset M$ an invariant compact set for X^t. Λ has a hyperbolic structure for X^t if and only if the linear Poincaré flow restricted over Λ is hyperbolic.*

2.2. **Holonomy maps.** Let $\Lambda\subset M$ be a compact invariant subset with dominated splitting and such that $\Lambda\cap\mathrm{Sing}(X)=\emptyset$. Let U_a be a neighborhood of Λ in $\mathcal{N}$, for $a\in(0,1]$, where it is possible to find an invariant a-cone field. Denote by $\mathcal{N}_x(\varepsilon)$ the ball of radius $\varepsilon>0$ on $\mathcal{N}_x$. The set $\mathrm{Sing}(X)$ is closed, then there exist a neighborhood K such that

$$\Lambda\subset\mathrm{int}(K)\subset\mathrm{cl}(K)\subset U_1\subset M\setminus\mathrm{Sing}(X)$$

Hence, there is $\varepsilon^*>0$ such that the exponential map $\exp_x{:}\mathcal{N}_x(\varepsilon^*)\longrightarrow M$ is an isometry for any point $x\in K$. Call $N_x=\exp_x(\mathcal{N}_x(\varepsilon^*))$. Moreover,

$\forall x \in K$ there is $\eta_x > 0$ such that $X^t(N_x) \cap N_x = \emptyset$, $\forall t \in (-\eta_x, \eta_x)$, since there are no singularities in $\mathrm{cl}(K)$; and hence, the function $x \mapsto \eta_x$ has a lower bound, say $\eta_* > 0$. Using the Implicit Function Theorem define for any $x \in U_1$ and $s \in \mathbb{R}$ the map

$$\tau_x^s \colon D_x^s \subset N_x \longrightarrow \mathbb{R}$$

in such a way that $\tau_x^s(x) = s$ and $X^{\tau_x^s(y)}(y) \in N_{x_s}$, where $x_s = X^s(x)$. Whenever we speak about D_x^s, we are considering it as the maximal domain of definition. In spite of this, set

$$R_x^s \colon D_x^s \longrightarrow N_{x_s}$$

such that $R_x^s(y) := X^{\tau_x^s(y)}(y)$, $\forall y \in D_x^s$. Notice $D_x R_x^s = L_x^s$. Moreover, there exist a neighborhood $V_s \subset \mathcal{N}$ of the zero-section, where we can define a transformation $R^s \colon V_s \longrightarrow N$ that maps

$$V_s \ni (x, v) \overset{R^s}{\longmapsto} (X^s(x), \exp_{x_s}^{-1}(R_x^s(\exp_x(v))))$$

that covers the flow on M, *i.e.* the following diagram conmutes:

$$\begin{array}{ccc} V_s & \xrightarrow{R^s} & \mathcal{N} \\ \downarrow & & \downarrow \\ M & \xrightarrow{X^s} & M \end{array}$$

In fact, $[V_s]_x \supset \exp_x^{-1}(D_x^s)$, for all $x \in U_1$. Observe $\{R^t\}$ is a local flow on $\bigcap_{t\in\mathbb{R}} V_t$.

3. Proof of Theorem A

This section shall be devoted to give a proof of Theorem A assuming Theorem B. Consider this two subsets of the space of vector fields, $\mathcal{X}^1(M)$:

$$H = \{X \in \mathcal{X}^1(M) | \, \mathrm{Sing}(X) \cap \mathrm{cl}(\mathrm{Per}(X)) \neq \emptyset\}$$

$$Tg = \{X \in \mathcal{X}^1 | X \text{ has a homoclinic tangency } \}$$

and define $\mathcal{U} = \mathcal{X}^1(M) \setminus \mathrm{cl}(H \cup Tg)$; takeing the closure on the C^1-topology. The set $\mathcal{U}$ consist of all vector fields which are not C^1-approximated by a homoclinic bifurcation. Denote by $\mathrm{Per}_h(X)$ the set of hyperbolic periodic orbits of X^t of saddle type; we shall prove that for elements in $\mathcal{U}$, the closure of this set has a dominated splitting.

Recall the notion of angle between one dimensional subspaces of the plane: Let us assume $\mathbb{R}^2 = E \oplus F$, for two given linear subspaces E and F. The angle between them is defined by $\angle(E, F) := ||L||$, where $L \colon E \longrightarrow E^\perp$ is the operator such that $F = \mathrm{graph}(L) = \{u + Lu | \, u \in E\}$.

On the other hand, given $X \in \mathcal{X}^1(M)$ and any point $p \in \mathrm{Per}_h(X)$, let $\tilde{E}^s_p(X)$ and $\tilde{E}^u_p(X)$ be the stable and unstable 2-dimensional subspaces of T_pM, invariant for the derivative of the flow, respectively. We can define $E^s_p = \tilde{E}^s_p \cap \mathcal{N}_p$ and $E^u_p = \tilde{E}^u_p \cap \mathcal{N}_p$, since $X(p) \neq 0$ for any $p \in \mathrm{Per}_h(X)$.

Remark 1. For $p \in \mathrm{Per}_h(X)$ we have $L^{t_p}_p = D_p X^{t_p}|_{\mathcal{N}_p}$, where t_p denotes the period of p. Moreover, E^s_p and E^u_p are invariant by $L^{t_p}_p$ and also $R^{t_p}_p : N_p \longrightarrow N_p$ is a diffeomorphism. Actually, this is the usual Poincaré first return map on p.

The next lemma relates the angle between the stable and unstable spaces on hyperbolic periodic points of saddle type and the existence of a dominated splitting over the colsure of these points.

Lemma 3.1. *Let $X \in \mathcal{X}^1(M)$. If there exists $\gamma > 0$ and a neighborhood $\mathcal{V}(X)$ of X such that for any $Y \in \mathcal{V}(X)$ we have*

$$\angle(E^s_p, E^u_p) > \gamma; \forall p \in Per_h(Y)$$

then, $cl(Per_h(X))$ has a Dominated Splitting.

Proof. To show the existence of a dominated splitting on $\mathrm{cl}(\mathrm{Per}_h(X))$ one must proceed to verify that the splitting on $\mathrm{Per}_h(X)$ given by the stable and unstable spaces on periodic points is in fact dominated. Then, one can easily extend such splitting to the closure. To see that, we remark that the argument of the proof of Lemma 2.0.1 of [21] relies only on some estimates on the eigenvalues of the derivative on periodic points. In our case, all these estimates remain true for $R^{t_p}_p$ for each $p \in \mathrm{Per}_h(X)$, since each return map is a diffeomorphism (see Remark 1). Finally, in [4] we can see how to perturb the flow in order to obtain the desired Poincaré maps. □

Now we can state the fundamental relation between homoclinic tangencies and the existence of a dominated splitting.

Theorem 3.2. *If X is a Kupka Smale vector field in $\mathcal{U}$, then the $cl(Per_h(X))$ has a Dominated Splitting.*

It is not difficult to see that the proof of Theorem 3.2 reduces to prove the following lemma:

Lemma 3.3. *Let $X \in \mathcal{U}$ be a Kupka Smale vector field on M. Then there exists a neighborhood $\mathcal{V}(X)$ of X and a number $\gamma > 0$ such that $\forall Y \in \mathcal{V}(X)$ and $\forall p \in Per_h(Y)$ the angle*

$$\angle(E^s_p(Y), E^u_p(Y)) > \gamma.$$

Proof. See the proof of Lemma 2.2.2 in [21], and consider Remark 1. □

Let $\Omega(X)$ be the non wandering set of X and denote by $P_0(X)$ and $F_0(X)$ the sets of attracting and repelling periodic orbits, respectively. Let $\Omega_0(X) = \Omega(X) \setminus (P_0(X) \cup F_0(X) \cup \mathrm{Sing}(X))$.

Proposition 3.4. *There is an open and dense set $\mathcal{U}_1 \subset \mathcal{U}$ such that for any $Y \in \mathcal{U}_1$, $\Omega_0(Y)$ is closed and it has a dominated splitting.*

Proof. Observe that the function Γ that associates to each vector field X the set $\mathrm{cl}(\mathrm{Per}_h(X))$ is lower semi-continous; since hyperbolic periodic orbits cannot be destroyed by small perturbations.

Call G_1 the resiudal set of continuity points of Γ. Denote by G_2 the set of Kupka-Smale flows on M (*i.e.* periodic orbits and singularities are all hyperbolic and their stable and unstable manifolds are in general position) and set $G_3 = \{X | \Omega(X) = \mathrm{cl}(\mathrm{Per}(X)) \cup \mathrm{Sing}(X)\}$, which is also residual (see [19]). Hence, $G = G_1 \cap G_2 \cap G_3$ is a residual set.

Lemma 3.5. *For any $X \in G \cap \mathcal{U}$, we have $\Omega_0(X) = \Gamma(X)$.*

Proof of the Lemma. In order to see that $\Omega_0(X) \subset \Gamma(X)$, suppose there is $q \in \Omega_0(X) - \Gamma(X)$. Let U be an open set such that $\Gamma(X) \subset U$ and $q \notin \mathrm{cl}(U)$. Let $N \subset \mathcal{U}$ be a neighborhood of X where $\Gamma(Y) \subset U$ for any $Y \in N$. Since $X \in G_3$, there is a sequence $\{p_n\}$ of points of $\mathrm{Per}(X) \cup \mathrm{Sing}(X)$ such that $p_n \notin \mathrm{cl}(U)$ and $p_n \to q$. Since $\#\mathrm{Sing}(X) < \infty$ and $q \notin \Gamma(X)$ then we have $p_n \in P_0(X) \cup F_0(X)$ for all n. Without lost of generality we may think $p_n \in P_0(X)$ for all n. Let us recall the following theorem due to Pliss.

Theorem 3.6 (Pliss, [18])**.** *Let $X \in \mathcal{X}^1(M)$ and suppose there is an infinite sequence $\{p_n\} \subset P_0(X)$, then for any $\epsilon > 0$ there is $Y \in \mathcal{X}^1(M)$, ϵ-C^1-close to X and some $n \in \mathbb{N}$ such that p_n is a nonhyperbolic periodic orbit of Y.*

It is not difficult to see that if we take a very small perturbation of the map Y, we can assume p_n is an hyperbolic periodic orbit of saddle type. Hence we have found some $Y \in N$ with a hyperbolic periodic orbit of saddle type p_n outside U; contradicting the fact of $\Gamma(Y) \subset U$. Therefore $\Omega_0(X) \subset \Gamma(X)$. On the other hand, to prove $\Gamma(X) \subset \Omega_0(X)$ we claim that $\Omega_0(X)$ is closed. If it is not the case, it happens that $\mathrm{cl}(\Omega_0(X)) \cap \mathrm{Sing}(X) \neq \emptyset$. Since $\mathrm{cl}(\Omega_0(X)) \subset \Gamma(X)$, we have found a contradiction to the fact that $\Gamma(X) \cap \mathrm{Sing}(X) = \emptyset$, proving $\Omega_0(X)$ is a closed set and hence the lemma follows. □

Let $X \in G \cap \mathcal{U}$ hence, we have $\Omega_0(X) = \mathrm{cl}(\mathrm{Per}_h(X))$. As a consequence of Theorem 3.2 we know that $\Omega_0(X)$ has a dominated splitting.

Take an admisible neighborhood U of $\Omega_0(X)$, that is, a neighborhood U such that there is $\mathcal{U}(X)$ a neighborhood of X such that for any $Y \in \mathcal{U}(X)$ the maximal invariant set of Y in U also has a dominated splitting. We can assume $\mathcal{U}(X) \subset \mathcal{U}$. Since $X \in G_1$ it is a continuity point of Γ; given a neighborhood U_0 of $\Omega_0(X)$ such that $U_0 \subset \mathrm{cl}(U_0) \subset U$, we can assume that for all $Y \in \mathcal{U}(X)$ verify that $\mathrm{cl}(\mathrm{Per}_h(Y)) \subset U_0$.

Let us prove that for all $Y \in \mathcal{U}(X)$ we have $\Omega_0(Y) \subset U$. Assume this do not happen, so, there is $Y \in \mathcal{U}(X)$ and $x \in \Omega_0(Y) \cap (M \setminus U)$. Notice x is not a periodic point and is non wandering. Moreover, x cannot be accumulated by points in $P_0(Y) \cup F_0(Y)$, otherwise, as in the proof of the Lemma 3.5, we can find an hyperbolic periodic orbit of saddle type outside U_0, which is imposible.

Now the C^1-Closing Lemma [19] allow us to find $\tilde{Y}$ close to Y having a periodic orbit passing through x. Actually, perturbing a bit more if necessary, this orbit is hyperbolic of saddle type. This is a contradiction to the continuity of Γ on X. Hence, for all $Y \in \mathcal{U}(X)$, the set $\Omega_0(Y)$ has a dominated splitting. Finally, the set $\mathcal{U}_1 = \bigcup_{X \in G} \mathcal{U}(X)$ verifies the Proposition. □

Now we can give a proof of Theorem A assuming Theorem B.

Proof of Theorem A. Given $X \in \mathcal{U}$, Proposition 3.4, together with Theorem B imply there is a C^2-Kupka-Smale vector field $Y \in \mathcal{U}_1$ arbitrarily close to X such that $\Omega_0(Y) = \Lambda \cup \mathcal{T}$, where Λ is hyperbolic and $\mathcal{T}$ is a finite union of irrational torus. However, the existence of such torus is not generic, even in the C^2 topology. Hence, making a small perturbation if necessary, we obtain that $\Omega_0(Y)$ is hyperbolic.

Now we claim $\Omega(Y)$ is hyperbolic. For that, we just need to guarantee that $P_0(Y) \cup F_0(Y)$ is a finite set. In fact, for instance, suppose $\#P_0(Y) = \infty$. Let U be a neighborhood of $\Omega_0(Y)$ where the maximal invariant set on U is hyperbolic. Notice we can take U in such a way all periodic orbits in U are of saddle type. On the other hand, all but a finite number of periodic attracting orbits must belong to U. In fact, $\mathrm{cl}(P_0(Y)) \setminus P_0(Y) \subset \Omega_0(Y) \subset U$, by definition. This is a contradiction. Therefore $\#P_0(Y) < \infty$. This argument applies also to $F_0(Y)$. Hence, we conclude $\Omega(Y)$ is hyperbolic.

It is left to prove $\Omega(Y) = \mathrm{cl}(\mathrm{Per}(X))$ and that there are not cycles. Since $L(Y) \subset \Omega(Y)$, the limit set of Y is hyperbolic too and so, the periodic orbits are dense on it. By a result of Newhouse [11], we only need to verify the absence of cycles on $L(Y)$ in order to conclude $L(Y) = \Omega(Y)$. Since there are no singularities of the flow involved, the index of all saddle periodic orbits is one. And since Y is Kupka-Smale

vector field, all the invariant manifolds are in general position, hence there are no cycles.

Now, we have already prove that X can be approximated by Y either: Uniformly hyperbolic with no-cycle condition, exhibiting a homoclinic tangency, or X belongs to the class H. However, if $X \in H$, it is an easy consequence of Hayashi's Connecting Lemma [4] that there is $Y \in \mathcal{X}^1(M)$ arbitrarly close to X showing a singular cycle, concluding the proof of Theorem A. □

4. Central Manifolds

Throughout this section we shall construct central-stable and central-unstable manifolds for invariant sets with dominated splitting. Moreover, we shall see that in our case they are of class C^2. Also we shall prove an analogous version of Pliss' Lemma (see [18]) for continous-time dynamical systems. As a consequence of these, we shall obtain that there can only exist finitely many irrational tori contained in Λ. At the end, we shall reduce the proof of Theorem B to a Main Lemma.

Let $X \in \mathcal{X}^2(M)$ and $\Lambda \subset M$ be a compact invariant set with dominated splitting such that $\mathrm{Sing}(X) \cap \Lambda = \emptyset$ and all periodic points in Λ are hyperbolic saddles. For simplicity on notation we call $x_t = X^t(x)$, for any $x \in M$ and $t \in \mathbb{R}$. We begin establishing the following inequalities.

Lemma 4.1. *If Λ is an invariant set as above, there exist constants $C > 0$ and $\sigma \in (0,1)$ such that for every $x_0 \in \Lambda$ and for all $t \geqslant 0$ the following holds:*

1. $|L^t|_{E(x_0)}||L^{-t}|_{F(x_t)}|^2 \leqslant C\sigma^t$,
2. $|L^t|_{E(x_0)}|^2|L^{-t}|_{F(x_t)}| \leqslant C\sigma^t$,

Proof. We only care about the first inequality since the other is analogous. Take $\alpha \in (0,1)$. We claim there is $T > 0$ such that for any $x \in \Lambda$

$$\frac{|L^T|_{E(x)}|}{|L^T|_{F(x)}|^2} \leqslant \alpha$$

After the claim, it is not difficult to see how the lemma follows; we only have to notice that $|L^t|_E| = \Pi_{j=1}^{\alpha}|L^{jT}|_E||L^r|_E|$ where $t = \alpha T + r$, and the same for the subspace F. Hence, setting

$$C = \max\{|L^r|_{E(x)}|, |L^r|_{F(x)}|; x \in \Lambda,\, r \in [0,T]\},$$

we conclude

$$\frac{|L^t|_{E(x_0)}|}{|L^t|_{F(x_t)}|^2} \leqslant C(\alpha^{1/t_0})^t.$$

Let us assume that the claim do not happens, that is, for all $T > 0$ there is $x^T \in \Lambda$ such that

$$\frac{|L^T|_{E(x^T)}|}{|L^T|_{F(x^T)}|^2} \geqslant \alpha$$

Take the measures $\mu_T = \frac{1}{T}\int_0^T \delta_{x_s^T} ds$, where $x_s^T = X^s(X^T)$ and consider a subsequence $\mu_i := \mu_{T_i}$ converging to a measure μ with $\mathrm{supp}(\mu) \subset \Lambda$. Observe the function $\frac{|L^t|_{E(x)}|}{|L^t|_{F(x_t)}|^2}$ is C^1 with respect to t. Let

$$\varphi(x) = \frac{d}{dt}\log\big(\frac{|L^t|_{E(x)}|}{|L^t|_{F(x_t)}|^2}\big)\big|_{t=0}$$

and notice it is continous on x. It is true that $\int \varphi d\mu \geqslant 0$. In fact,

$$\int \varphi d\mu = \lim_{i\to\infty} \varphi d\mu_i = \lim \frac{1}{T_i}\int_0^{T_i} \varphi(x_s^{T_i}) ds = \lim_{i\to\infty} \frac{1}{T_i}\log\big(\frac{|L^t|_{E(x)}|}{|L^t|_{F(x_t)}|^2}\big)$$

Hence, the Ergodic Decomposition Theorem implies that there is an ergodic measure ν with $\mathrm{supp}(\nu) \subset \mathrm{supp}(\mu)$ and such that $\int \varphi d\nu \geqslant 0$. Denote by $\lambda_E = \lambda_E(\nu) = \int \log |L^1|_E| d\nu$, and $\lambda_F = \lambda_F(\nu) = \int \log |L^1|_F| d\nu$, the Lyapunov Exponents of ν with respect to E and F, respectively. It is true that $\lambda_E < 0$ and $\lambda_F < 0$, since $\int \varphi d\nu = \lambda_E - 2\lambda_F \geqslant 0$ and the condition of domination implies $\lambda_E < \lambda_F$. Hence $0 \leqslant \lambda_E - 2\lambda_F < \lambda_F - 2\lambda_F = -\lambda_F$. Therefore $\lambda_E < 0$ and so $\lambda_F < 0$. Now we claim that the support of ν is contained in the orbit of some periodic point of the flow, that must be a sink. Oseledets' Theorem [13], asserts that ν-almost everywhere there is a unique invariant splitting $E_1 \oplus [X] \oplus E_2$ for X^1. For that and since the splitting $E \oplus F$ is invariant by L^1 we can call E_i that $E \oplus [X] = E_1 \oplus [X]$ and $F \oplus [X] = E_2 \oplus [X]$. Denote λ_1, λ_2, $\lambda_{[X]} = 0$ their respective Lyapunov Exponents. Notice

$$\begin{aligned} \lambda_E &= \int \log(L^t|_E) d\nu = \int \log(\det(DX^1|_{E\oplus[X]})) d\nu \\ &= \int \log(\det(DX^1|_{E_1\oplus[X]})) d\nu = \lambda_1 + \lambda_{[X]} = \lambda_1 \end{aligned}$$

Analogously $\lambda_F = \lambda_2$; therefore $\lambda_1 > 0$ and $\lambda_2 > 0$, and the claim follows. □

Now, our setting fits into the arguments of [6] to prove the existence of central stable and central unstable manifolds tangent to directions E and F, respectively. This is sumarized in the following lemma.

Lemma 4.2. *For each $\varepsilon > 0$ denote the interval $I_\varepsilon = (-\varepsilon, \varepsilon)$ and let $Emb^2(I_1, M)$ be the set of C^2 embeddings of I_1 on M. There exist two continous sections*

$$\Phi^{cs\,(cu)}: \Lambda \longrightarrow Emb^2(I_1, \mathcal{N})$$

such that, if we denote $W_\varepsilon^{cs\,(cu)}(x) = \Phi^{cs\,(cu)}(x)(I_\varepsilon)$, respectively, they verify the following properties:

a) *$T_x W_\varepsilon^{cs}(x) = E(x)$ and $T_x W_\varepsilon^{cu}(x) = F(x)$.*
b) *For all $\varepsilon_1 \in (0,1)$ there exists $\varepsilon_2 > 0$ such that $R^t(W_{\varepsilon_2}^{cs}(x)) \subset W_{\varepsilon_1}^{cs}(x_t)$ and $R^{-t}(W_{\varepsilon_2}^{cu}(x)) \subset W_{\varepsilon_1}^{cu}(x_{-t})$, for all $t \in [0,1]$.*

Notice that $\Phi^{cs}(x)(I_1) \subset \mathcal{N}_x$ since Φ^{cs} is a section on $\mathcal{N}$ and the same for Φ^{cu}. We shall call the manifold $W_{(\varepsilon)}^{cs}$ the (local) central stable manifold and $W_{(\varepsilon)}^{cu}$ the (local) central unstable manifold. Observe that the second part of the lemma implies the next corollary:

Corollary 4.3. *Given $\varepsilon > 0$ there is δ such that:*

a) *If $y \in W_\varepsilon^{cs}(x)$ and $dist(x_t, y_t) \leqslant \delta$ for all $t \in (0, t_0)$, then $R^t(y) \in W_\varepsilon^{cs}(x_t)$ for $t \in (0, t_0)$.*
b) *If $y \in W_\varepsilon^{cu}(x)$ and $dist(x_{-t}, y_{-t}) \leqslant \delta$ for all $t \in (0, t_0)$, then $R^{-t}(y) \in W_\varepsilon^{cs}(x_{-t})$ for $t \in (0, t_0)$.*

It is not difficult to see that in our context it is true the following:

Corollary 4.4. *Let $x \in \Lambda$ such that for some $\gamma \in (0,1)$ we have*

$$|L^t|_{E(x)}| \leqslant \gamma^t, \text{ for all } t \geqslant 0.$$

Then there exists $\varepsilon > 0$ such that $|R^t(W_\varepsilon^{cs}(x))| \to 0$ as $t \to \infty$; that means the central stable manifold is in fact a stable one.

4.1. **Hyperbolic times of a point.** It is possible to prove a version of Pliss' Lemma [18] for flows. In this case, the proof is easier than in the case of diffeomorphisms (see [8]).

Theorem 4.5. *Given any $\varepsilon > 0$, $A \in \mathbb{R}$ and $c > 0$, there exists $L > 0$ such that if $H: [0,T] \longrightarrow \mathbb{R}$ is diferentiable, $H(0) = 0$, $H(T) < cT$ and $\inf(H') > A$, then the set*

$$\mathcal{P}_\varepsilon = \{\tau \in [0,T] | H(s) - H(\tau) \leqslant (c+\varepsilon)(s-\tau) \text{ for all } s \in [\tau, T]\}$$

has Lebesgue measure greater than TL.

Proof. Take $\varepsilon > 0$ and A as in the hypothesis. Define $G(s) = H(s) - (c+\varepsilon)s$. We assume G is of class C^2 and it do not have degenerate critical points; that is, $G'(x) = 0$ if and only if $G''(x) \neq 0$. Actually we can assume all their critical values are different, otherwise, an approximation of G still works. Since $G(0) = 0$ and $G(T) < -\varepsilon T$, it is

possible to define two (perhaps finite) sequences, say $\{a_i\}$ consisting of critical points of G such that $G(x) < G(a_i)$ for every $x > a_i$ (if finite, set the last point $a_{n+1} = T$, if it is not so, then $a_i \to T$) and $\{b_i\}$ as the first number $b > a_i$ such that $G(b_i) = G(a_{i+1})$. Of course $b_i \leqslant a_{i+1}$. Let $B = -\inf G'$, then by the Mean Value Theorem

$$\frac{G(a_i) - G(b_i)}{b_i - a_i} \leqslant B.$$

Hence

$$\frac{G(a_i) - G(b_i)}{B} \leqslant b_i - a_i.$$

On the other hand, $\bigcup_i (a_i, b_i) \subset B_\varepsilon$. In fact,

$$\begin{aligned} H(s) - H(\tau) &= H(s) - (c+\varepsilon)s - (H(\tau) - (c+\varepsilon)\tau) + (c+\varepsilon)(s-\tau) \\ &= G(s) - G(\tau) + (c+\varepsilon)(s-\tau) \end{aligned}$$

Hence, $G(s) - G(\tau) + (c+\varepsilon)(s-\tau) < (c+\varepsilon)(s-\tau)$ if and only if $G(s) < G(\tau)$. Let $t \in (a_i, b_i)$, for some i. If for a given $s > \tau$ we have $G(s) = G(\tau)$ then $\tau = b_i$, because b_i is the first number $b > a_i$ such that $G(b) = G(a_{i+1})$; a contradiction. Therefore,

$$\begin{aligned} |\mathcal{P}_\varepsilon| &\geqslant \textstyle\sum_{i=1}^n (b_i - a_i) \geqslant \frac{1}{B} \sum_{i=1}^n G[(a_i) - G(b_i)] = \\ &= \textstyle\frac{1}{B} \sum_{i=1}^n [G(a_i) - G(a_{i+1})] = \frac{1}{B}(G(a_1) - G(T)) \end{aligned}$$

Notice $G(a_1) \geqslant 0$ since it is a global maximum. Therefore $|\mathcal{P}_\varepsilon| \geqslant \frac{-G(T)}{B} \geqslant \frac{\varepsilon T}{B}$, and then $|\mathcal{P}_\varepsilon| T^{-1} \geqslant \varepsilon B^{-1}$. Finally $G'(t) = H'(t) - (c+\varepsilon)$ and $B = \inf(G') = (c+\varepsilon) - A \geqslant c + \varepsilon - A$, that is

$$\frac{|\mathcal{P}_\varepsilon|}{T} \geqslant \frac{\varepsilon}{c + \varepsilon - A} =: L$$

□

Corollary 4.6. *Let $X \in \mathcal{X}^1(M)$ and let Λ be an invariant compact set with dominated splitting. Given $\varepsilon > 0$, $\gamma < 1$ and $x \in \Lambda$. If there is $T > 0$ such that $|L^T|_{E(x)}| < \gamma^T$, then there is $L > 0$ such that the set*

$$\mathcal{P}_\varepsilon(x) = \{\tau \in [0,T] \,||L^{s-\tau}|_{E(x)}| \leqslant (e^{log(\gamma)+\varepsilon})^{s-\tau} \,\forall s \in [\tau, T]\} \tag{1}$$

is such that leb$(P_\varepsilon(x)) \geqslant LT$, and hence it is non-empty.

Proof. Recall the function $H_E(t) = \log(|L^t|_{E(x)}|)$ is C^1. Moreover $d/dt(H_E) =< D_{x_t}X(E), E >$, and hence $|H'(t)| \leqslant ||X||_{C^1}$. Also remember $DR^t_x(v) = D_xX^t(v) - \frac{<D_xX^t(v), X(x)>}{||X(x)||^2} X(x)$. □

Corollary 4.7. *Given $\varepsilon > 0$, $\gamma < 1$ and $x_0 \in \Lambda$ if there exists $t > 0$ such that $|L^s|_{E(x_t)}| < \gamma^s$, for all $s \geqslant 0$, then for $\tilde{\lambda} = e^{\log(\gamma)+\varepsilon}$, the set*

$$\mathcal{P}(x_0, \tilde{\lambda}) = \{\tau \in \mathbb{R} \,||L^s\,|_{E(x_\tau)}| \leqslant \tilde{\lambda}^s \,\forall s > 0\} \tag{2}$$

is unbounded.

We call the set on (2) the hyperbolic times of x_0 related to $\tilde{\lambda}$.

4.2. **Reducing Theorem B to a Main Lemma.** Now, it is not difficult to see that Theorem B follows from the theorem 4.8 below, once we guarantee there can only be finitely many irrational tori on Λ.

Theorem 4.8. *Let $X \in \mathcal{X}^2(M)$ and let Λ be a compact invariant for X set with dominated splitting; also assume all periodic points in Λ are hyperbolic of saddle type and $Sing(X) \cap \Lambda = \emptyset$. Then one of the following happens:*

1. *Λ is an hyperbolic set.*
2. *There exist an irrational torus $\mathcal{T}$ contained in Λ.*

Proposition 4.9. *There are only finitely many irrational tori in Λ.*

Proof. Assume by contradiction there are infinitely many irrational tori, named $\{\mathcal{T}_k\}_{k\in\mathbb{N}}$. Take numbers $\lambda < \gamma_1 < \gamma_2 < 1$ and $c > 0$ such that $(1+c)\lambda < 1$. On each $\mathcal{T}_k$ there exist a point x^k, and some $t_k > 0$ such that

$$|L^t_{F(x^k)}| \leqslant (1+c)^t, \ \forall t \geqslant t_k$$

and then, by the domination condition we have

$$|L^t_{E(x^k)}| \leqslant \lambda^t(1+c)^t < \gamma_1^t, \ \forall t \geqslant t_k.$$

An application of Lemma 4.5 imply the existence of $T_k > 0$ such that for $y^k := X^{T_k}(x^k)$, $|L^t|_{E(y^k)}| \leqslant \gamma_2^t$, for all $t > 0$. Hence, Corollary 4.4 let us to conclude that each y^k has a stable manifold of size $\varepsilon > 0$.

Let $y \in \Lambda$ be an accumulation point of the sequence $\{y^k\}$ and denote by $\Sigma_y \subset N_y \subset M$ an open ball arround y of radius smaller than $\varepsilon/3$. Still denote y^k the points on Σ_y which are inmediate iterates of the true y^k. Actually, they also have stable manifolds of uniform size ε. Notice that it is not possible $\mathcal{T}_k \cap \Sigma_y$ be a circle for infitely many k, because the splitting on Λ is continous. Moreover, $\mathcal{T}_k \cap \Sigma_y$ is an arc that meets the boundary of the ball in two different connected components. So, it is not difficult to see there are $\mathcal{T}_j$ and $y^k \in \mathcal{T}_k \cap \Sigma_y$, $j \neq k$ that $\mathcal{T}_j \cap \Sigma_y \neq \emptyset$ and $W^s_\varepsilon(y^k) \cap \mathcal{T}_j \neq \emptyset$. This contradiction proves the Proposition. □

In [21], they prove Theorem 4.8 is reduced to the next Main Lemma, and their arguments work for both, diffeomorphisms and vector fields. We only notice that in spite of Lemma 2.2, it is elementary to verify that if Λ is a compact invariant set with dominated splitting such that for every $x \in \Lambda$ we have $|L^t|_{E(x)}| \to 0$ and $|L^{-t}|_{F(x)}| \to 0$ as $t \to \infty$, then Λ is a hyperbolic set.

Main Lemma 1. *Let $\Lambda_0 \subset M$ be a compact transitive invariant set for X^t with a dominated splitting such that $\Lambda_0 \cap Sing(X) = \emptyset$ and it is not an irrational torus neither a periodic orbit. If every properly contained invariant subset of Λ_0 is hyperbolic, then Λ_0 is hyperbolic itself.*

Hence, we have to focus on the proof of the Main Lemma. This will be done in the last section.

5. Asymptotical Behaivor of Dynamical Intervals

In order to obtain hyperbolicity on an invariant set with dominated splitting one must show that the central manifolds on each point are actual stable and unstable manifolds. This is the aim of this section. For that, we shall prove a Denjoy's like property which, roughly speaking, states that smoothness on the vector field X do not allow the existence of some kind of wandering intervals. (The precise statement is Lemma 5.2). Let $X \in \mathcal{X}^2(M)$ and let Λ be compact invariant set with dominated splitting such that $\mathrm{Sing}(X) \cap \Lambda = \emptyset$ and assume that all periodic points in Λ are hyperbolic saddles. Consider the set

$$\Lambda^+ := \cap_{t \geqslant 0} X^{-t}(\mathrm{cl}(V));$$

for some neighborhood $V \subset \mathrm{cl}(V) \subset U_1$. Recall U_1 is a neighborhood of Λ where it is possible define an extension of the invariant 1-cone field on Λ (see Section 2.1). Notice this induces an extension of the dominated splitting of Λ over Λ^+.

We say $I \subset \mathcal{N}_x$ or $I \subset N_x$ are intervals if they are the image of a smooth function $\varphi\colon [0,1] \longrightarrow \mathcal{N}_x$ or N_x, respectively. Denote by $|I|$ the lenght of the interval I. The following definitions collect the insight of central unstable manifolds and the property that will make them actual unstable manifolds.

Definition 2. Let I be an interval contained in $\mathcal{N}_{x_0}$ for some $x_0 \in I$ and let $\delta > 0$.

1. We say I is a-transversal to the E direction for some $a \in (0,1]$ if $T_y I \subset C_a^F(y) \subset \mathcal{N}_{x_0}$ for any $y \in I$. It is E-transversal if it is a-transversal to the direction E for some $0 < a \leqslant 1$.
2. We say I is a (δ, E)-interval if it is E-transversal, $I \subset \cap_{t \geqslant 0} V_t$ and $|R^t(I)| \leqslant \delta$ for all $t \geqslant 0$.

Although these definitions are stated for intervals contained in the normal bundle, we say an interval $I \subset N_x$ on M is E-transversal or (δ, E)-transversal if the interval defined by $\hat{I} := \exp_x^{-1}(I)$ is. Notice if $\hat{I} \subset \mathcal{N}$ is a (δ, E)-interval, then $\exp_x(\hat{I}) \subset \Lambda^+$.

Remark 2. For any $c > 0$ there is $\delta_c > 0$ such that if I is a (δ, E)-interval associated to $x \in I$ with $\delta \leqslant \delta_c$ then, for any $y \in I$:

$$(1-c)^t \leqslant \frac{|D_y R_x^t|_{E(y)}|}{|L^t|_{E(x)}|} \leqslant (1+c)^t$$

And a similar statement with respect to the F-direction.

Any (δ, E)-interval contained on a sets with dominated splitting it has stable manifolds on each of its points, as we shall see on the next Lemma. The key idea behind this is that an interval transversal to direction E which do not grows its lenght on future iterations it must contract exponentially along the E-direction.

Lemma 5.1. *For any $\gamma \in (\lambda, 1)$ there is $\delta_1 > 0$ such that if I is a (δ, E)-interval, $I \subset \mathcal{N}_x$, $x \in I$, with $\delta \leqslant \delta_1$, then there is $t_0 > 0$ such that*

$$|L^s|_{E(x_{t_0})}| < \gamma^s, \forall s \geqslant 0.$$

Moreover, for any $y \in \hat{I}_{t_0} = \exp_{x_{t_0}}(R^{t_0}(I))$ we have

$$|D_y R^s|_{E(x_{t_0})}| < \gamma^s, \forall s \geqslant 0.$$

and hence, on any point of $\hat{I}_{t_0}$ we have stable manifolds of uniform size.

Proof. Fix $\gamma \in (\lambda, 1)$ and $\lambda_2 < 1$ such that $\lambda\lambda_2^{-1} < \gamma$. Let $c > 0$ and $\lambda_3 < 1$ such that $(1-c)\lambda_2 < \lambda_3$. Take $\delta_1 \leqslant \delta_c$ of Remark 2, and let I be a (δ, E)-interval with $\delta \leqslant \delta_1$, associated to $x_0 \in I$

We claim there is $t_0 > 0$ such that $|L^s|_{F(x_{t_0})}| < \lambda_2^{-s}$, for all $s > 0$. To prove the claim suppose there exist a sequence $s_t \to \infty$ as $t \to \infty$ such that for all $t \geqslant 0$

$$|L^{s_t}|_{F(x_t)}| \geqslant \lambda_2^{-s_t}$$

Since $|\hat{I}_t| \leqslant \delta_c$, we have:

$$(1-c)^{-s} \leqslant \frac{|D_y R^s_{x_t}|_{\tilde{F}(y)}|}{|L^s|_{F(x_t)}|} \leqslant (1+c)^s.$$

Then

$$|D_y R^{s_t}_{x_t}|_{\tilde{F}(y)}| \geqslant (1-c)^{-s_t}|L^{s_t}|_{F(x_t)}| \geqslant ((1-c)\lambda_2)^{-s_t} \geqslant \lambda_3^{-s_t}$$

It is a straight forward computation that $|I_{t+s_t}| \geqslant \lambda_3^{-s_t}|I_t|$. If t is big enough, we achieve a contradiction to $|I_{t+s_t}| \leqslant \delta_c$. Hence, the claim follows.

Actually, we have proved that $|L^s|_{E(x_{t_0})}| \leqslant (\lambda\lambda_2^{-1})^s$, for all $s \geqslant 0$. Notice that if $c > 0$ is such that $(1+c)\lambda\lambda_2^{-1} < \gamma$ and $\delta > 0$ is small

enough that for all $y \in \hat{I}_{t_0}$ and for all $s \geqslant 0$

$$(1-c)^{-s} \leqslant \frac{|D_y R^s_{x_t}|_{E(y)}|}{|L^s|_{E(x_t)}|} \leqslant (1+c)^s$$

then $|D_y R^s_{x_t}|_{E(y)}| < \gamma^s$, for all $y \in \hat{I}_{t_0}$ and for all $s \geqslant 0$, as we required. Corollary 4.4 guarantees then stable manifolds on points of $\hat{I}_{t_0}$ of uniform size □

Maximal (δ, E)-intervals. Let $I \subset \mathcal{N}_{x_0}$ be a (δ, E)-interval associated to some $x_0 \in I$. Denote by $I_t := R^t(I)$ for all $t \in \mathbb{R}^+$. We seek for a family of (δ, E)-intervals J_t such that $I_t \subset J_t$ and also $R^{t-s}(J_s) \subset J_t$ for all $0 < s < t$, and any $t \in \mathbb{R}^+$; being maximal for these properties. In order to find it we appeal to Zorn's Lemma in the following way: Consider the set of functions from $\mathbb{R}$ into the compact connected subsets of the normal bundle $\mathcal{N}$, denoted by $P(\mathcal{N})$:

$$\begin{aligned}\mathcal{F} = \quad \{J{:}\mathbb{R} \longrightarrow P(\mathcal{N}) | J(s) \subset \mathcal{N}_{x_s},\ I_s \subset J(s),\ J(s) \text{ is a } (\delta, E)\text{-interval} \\ \text{and } R^t(J(s)) \subset J(s+t),\ \forall s, t \in \mathbb{R}^+\}\end{aligned}$$

with the order relation $J \leqslant K$ if and only if $J(s) \subset K(s)$, $\forall s \in \mathbb{R}^+$. Observe that the function $I(s) = R^s(I)$ belongs to $\mathcal{F}$. Moreover, any totally ordered chain has a supremum in $\mathcal{F}$, say, $\{J^i \in \mathcal{F}; i \in \Gamma\}$, then $J^\infty(s) = \bigcup_{i \in \Gamma} J^i(s)$, defined for each $s \in \mathbb{R}$ belongs to $\mathcal{F}$. Therefore, there is a maximal element on $\mathcal{F}$ denoted by $J_s := J(s)$. If there is an interval $\tilde{K} \subset N_{x_s}$ such that $J_s \subset \tilde{K}$ for some $s \in \mathbb{R}^+$ and $|R^t(\tilde{K})| \leqslant \delta$ for all $t \geqslant 0$, then the family $K(t) := J_t$ for $t < s$ and $K(t) := R^t(\tilde{K})$ for $t \geqslant s$ belongs to $\mathcal{F}$; contradicting the fact $J(\cdot)$ is a maximal element of $\mathcal{F}$.

Notice that Lemma 5.1 implies the existence of $\delta_0 > 0$ such that if $\{J_s\}$ is a maximal family of (δ, E)-intervals related to a (δ, E)-interval I $(\delta \leqslant \delta_0)$, there exist numbers $s > 0$ and $\varepsilon_0 > 0$ such that for every $y \in J_s$, the center-stable manifold $W^{cs}_{\varepsilon_0}(y)$ is in fact a true stable manifold for R^t, that is, $|R^t(W^{cs}_{\varepsilon_0}(y))| \to 0$ as $t \to \infty$. Moreover there exist $C > 0$ and $\gamma \in (0,1)$ such that

$$|R^t_{x_s}(\exp_{x_s}(W^s_{\varepsilon_0}(y)))| < C\gamma^t \text{ for any } t \in \mathbb{R},$$

that is, $\exp_{x_s}(W^s_{\varepsilon_0}(y))$ is a local stable manifold for $R^t_{x_s}$. We will denote also $W^s_{\varepsilon_0}(y) := \exp_{x_s}(\hat{W}^s_{\varepsilon_0}(y))$, except when it is relevant to remark the difference. For any $s \in \mathcal{P}$ denote by $J^\varepsilon_s := \bigcup_{y \in J_s} W^s_\varepsilon(y) \subset M$.

5.1. **Denjoy's Like Property.** Let $\hat{I} \subset \mathcal{N}_{x_0}$ for some $x_0 \in \hat{I}$ be a (δ, E)-interval. Denote by $I_t = \exp_{x_0}(\hat{I}_t) \subset M$, where $\hat{I}_t = R^t(\hat{I})$ and

$x_t := X^t(x)$. Define the ω-limit set of I_0 as $\omega(I_0) = \bigcup_{y\in I_0} \omega(y)$; where $\omega(y)$ is the usual ω-limit set of y according to the flow X^t.

Proposition 5.2. *There is $\delta_0 > 0$ such that if I is a (δ, E)-interval associated to some $x \in I$ for some $\delta < \delta_0$, then there are two posibilities: $\omega(I) \subset \mathrm{Per}(X^t)$ or $\omega(I) = \mathcal{T}$, where $\mathcal{T}$ is an irrational torus.*

Proof. Let $\lambda < 1$ be from the dominated splitting condition. Fix $\gamma \in (\lambda, 1)$ and take $\delta_1 > 0$ from Lemma 5.1. Let $I_0 \subset M$ be a (δ, E)-interval associated to x_0 with $\delta \leqslant \delta_1$. Denote by $I_t = R^t_x(I_0)$. Let $\{\hat{J}_s | s \in \mathbb{R}^+\}$ be a maximal family of (δ, E)-intervals for $\hat{I}_s = \exp^{-1}_{x_s}(I_s)$ constructed above and denote $J_s = \exp_{x_s}(\hat{J}_s) \subset M$. Lemma 5.1 imlplies there exist $t_0 > 0$ for which on any point of J_{t_0} we have defined a stable manifold of size ε_0. This fact, together with Corollary 4.7 imply that the set $\mathcal{P}(x_{t_0})$, the hyperbolic times of x_0 related to γ, is unbounded. Corollary 4.4 assert that for any $s \in \mathcal{P}(x_{t_0})$ and on any point of J_s, we also have stable manifolds of size ε_0. Since we are interested on asymptotical properties of I we shall assume $t_0 = 0$; in fact, $w(I) \subset w(J_{t_0})$. We shall denote, by simplicity $\mathcal{P} = \mathcal{P}(x_0)$. The following situations requires two different arguments:

I. There exist some $t, s \in \mathcal{P}$, $t < s$, for which $J^\varepsilon_s \cap [R^{t-s}(J_s)]^\varepsilon \neq \emptyset$.
II. For any $t, s \in \mathcal{P}$, we have $J^\varepsilon_s \cap [R^{t-s}(J_s)]^\varepsilon = \emptyset$.

Let us first deal with Case I. For that, we shall assume first that $\lim_{s\in\mathcal{P}} |\hat{J}_s| = 0$. If we can guarantee the hypothesis of the following lemma we shall conclude the thesis of Proposition 5.2 in this case.

Lemma 5.3. *If there exist $s \in \mathcal{P}$ and $t > 0$ such that $R^t_{x_s}(J_s) \subset J^\varepsilon_s$, then there is a periodic orbit $p \in \Lambda$ such that $\omega(I) \subset \omega(J_0) = \mathcal{O}(p)$.*

Proof of the Lemma. Define $\pi : J^\varepsilon_s \longrightarrow J_s$, a projection along stable manifolds. Then $\pi \circ R^t_{x_s} : J_s \longrightarrow J_s$ is a continous map of the interval. Since $R^t_{x_s}(J_s) \subset J^\varepsilon_s$, we see $\pi(R^t_{x_s}(J_s)) \subset J_s$. Therefore, there is $z \in J_s$ such that $R^t_{x_s}(z) \in W^s_\varepsilon(z)$; and this can only happen if both $z, R^t_{x_s}(z) \in W^s(p)$ of some periodic point $p \in \Lambda$. Moreover, p must be a sink and $\omega(I) \subset \omega(J_0) = \mathcal{O}(p)$. □

The following two claims let us fit into the hypothesis of the previous Lemma.

Claim 1. There is $\delta' > 0$, $s_0 > 0$ and $c > 0$ such that, if $\tilde{J}$ is a (δ', E)-interval satisfying $J^{c\varepsilon}_s \cap \tilde{J} \neq \emptyset$, with $s_0 < s \in \mathcal{P}$, then $\tilde{J} \subset J^\varepsilon_s$.

Claim 2. There is $s \in \mathcal{P}$, $s > s_0$ and $t > 0$ in such a way $W_{c\varepsilon}(J_s) \cap R^t_{x_s}(J_s) \neq \emptyset$, and $R^t_{x_s}(J_s)$ is a (δ', E)-interval.

Proof of Claim 1. Let $\eta_1 > 0$ such that $(1+\eta_1)\lambda < 1$. As we see in Remark 2, there is $\delta_{\eta_1} > 0$ such that if K is a (δ_{η_1}, E)-interval we have $\forall z, z_0 \in K$

$$\frac{|D_z R^t|_{E(z)}|}{|D_{z_0} R^t|_{E(z_0)}|} < (1+\eta_1)^t, \forall t \geqslant 0. \tag{3}$$

Let us suppose there is $\tilde{J}$ a (δ_{η_1}, E)-interval such that $J_s^{c\varepsilon} \cap \tilde{J} \neq \emptyset$ for some $c > 0$. Observe $\tilde{J} \subset D^t_{x_s}$, for all $t \geqslant 0$. Then there are points $z_0 \in \tilde{J}$ and $y_0 \in J_s$ such that $z_0 \in W^s_{c\varepsilon}(y_0)$. Moreover, for any $c > 0$ there is a positive $\eta_c > 0$ such that

$$\frac{|D_{z_0} R^t_{x_s}|_{E(z_0)}|}{|D_y R^t_{x_s}|_{E(y)}|} < (1+\eta_c)^t, \forall t \geqslant 0 \tag{4}$$

since both points lay on the same stable manifold $W^s_{c\varepsilon}(y)$.

Notice that if $\tilde{\lambda}_c = (1+\eta_1)(1+\eta_c)\lambda < 1$, inequalities (3) and (4) imply that on any point of $\tilde{J}$ there are stable manifolds of size $\varepsilon'(\tilde{\lambda}_c) > 0$. Actually, $\eta_c \to 0$ as $c \to 0$ and $\varepsilon'(\tilde{\lambda}_c)$ does not decrease. So, $|D_z(R^t_{x_s})|_{E(z)}| < \tilde{\lambda}_c^t$ for $t \geqslant 0$ and all $z \in \tilde{J}$.

Fix $c \in (0, \frac{1}{2})$ such that $\tilde{\lambda}_c < 1$ and take $\tilde{\varepsilon} > 0$ such that $c\varepsilon < \tilde{\varepsilon} < (1-c)\varepsilon$ and $\tilde{\varepsilon} \leqslant \varepsilon'(\tilde{\lambda}_c)$. Take $s_0 \in \mathcal{P}$ in such a way that for any $s > s_0$ we have $2|J_s| < \delta$. Then for all $z \in \tilde{J} \cap J_s^{c\varepsilon}$ we have $W^s_{\tilde{\varepsilon}}(z) \subset J_s^{\varepsilon}$ and $W^s_{\tilde{\varepsilon}}(z) \cap J_s \neq \emptyset$.

On the other hand, there is some constant $C > 0$ such that for any E-transversal interval $A \subset \tilde{J}^{\tilde{\varepsilon}}$ we have $|R^t(A)| \leqslant C|R^t(\tilde{J})|$ for any $t \geqslant 0$. Take δ' such that $\delta/2 + C\delta' < \delta$. If $\tilde{J}$ is not properly contained in J_s^{ε}, there is a C^2 extension of the interval J_s, named $J_s \cup A$ with $A \subset \tilde{J}^{\tilde{\varepsilon}}$, which is E-transversal. Moreover,

$$R^t(J_s \cup A) \leqslant |R^t_{x_s}(J_s)| + |R^t_{x_s}(A)| \leqslant \delta/2 + C\delta' < \delta.$$

The last inequality contradicts the fact of J_s being maximal (δ, E)-interval, and give us the claim. □

Proof of Claim 2. Actually, in the previous argument, we could take s_0 big enough in such a way that $|J_s| < \delta'$ and $\lambda^{s_0} < c$ also. The intersection property (hypothesis of case I) implies

$$J_s^{c\varepsilon} \cap R^t_{x_s}(J_s) \neq \emptyset$$

for some positive t. Since $R^t_{x_s}(J_s)$ is a (δ', E)-interval we are done. □

Now we assume $\limsup_{s\in\mathcal{P}} |J_s| > 0$. Considering any convergent subsequence we obtain a limit interval $J_s \to J$ provided of a stable manifolds, say, $J^{\varepsilon} := \cup_{z\in J} W^s_{\varepsilon}(z)$ for any $\varepsilon < \varepsilon_0$. There is no harm if we assume that J^{ε_0} lies in some N_x for some $x \in J \cap \Lambda$.

Lemma 5.4. *If $p \in J^\varepsilon \cap Per(X^t)$ then p is hyperbolic and $\omega(J) \subset Per(X^t)$.*

Proof. Notice that $p \in \Lambda^+$. Let us suppose it is a hyperbolic saddle. Observe that $J \cap W^u_\varepsilon(p)$ is non empty and transversal. Then, the Inclination Lemma [λ-Lemma in [15]] implies that $|W^u_\varepsilon(p)| \leqslant \delta$, since $|R^t_x(J)| \leqslant \delta$ for all $t \geqslant 0$. Hence, there must be two periodical sinks p_1, p_2 bounding a cylinder that contains the orbit of p. Therefore, if $y \in J$ the ω-limit of y is either p_1, p_2 or p. A similar argument implies the thesis in the case of p being a sink or a saddle node, and these are the only posibilities; since $p \in J^\varepsilon$ has a stable manifold. □

Hence, we assume there are no periodic orbits passing through J^ε. Notice that the set ∂J^ε consists of boundary of two kinds: $\partial^s(J^\varepsilon) := \bigcup_{z\in\partial J} W^s_\varepsilon(z)$ and $\partial^u(J^\varepsilon) := \partial(J^\varepsilon) \setminus \partial^s(J^\varepsilon)$. This distinction allows us to make the next definition.

Definition 3. If a set A is such that $A \cap J^\varepsilon \neq \emptyset$ we say A is stably contained on J^ε, $A \subset^s J^\varepsilon$ iff $A \cap \partial^u(J^\varepsilon) = \emptyset$. We also define, for positive ε and $\tilde{\varepsilon}$, the number $\rho(J^\varepsilon, J^{\tilde{\varepsilon}}) = \rho \in \mathbb{R}^+$ as the first time such that $R^\rho_x(J^\varepsilon) \cap J^{\tilde{\varepsilon}} \neq \emptyset$.

Proposition 5.5. *There exists two subintervals $\tilde{L} \subset L \subset J$ such that $w(J) = w(L)$, numbers $0 < t_0 < t_1$ and $\eta > 0$ such that:*

1. *$R^{t_0}_x(L^\eta) \subset^s L^\eta$ for $t_0 = \rho(L^\eta, L^\eta)$, and $R^{t_0}_x(L^\eta) \cap \tilde{L}^\eta = \emptyset$.*
2. *$R^{t_1}_x(\tilde{L}^\eta) \subset^s \tilde{L}^\eta$ for $t_1 = \rho(\tilde{L}^\eta, \tilde{L}^\eta)$.*

Before we prove this Proposition we need a Lemma.

Lemma 5.6. *Let $L \subset J$ be an interval, not containing a periodic orbit:*

1. *If there exists $M > 0$ such that $R^s_x(L) \cap L = \emptyset$ for all $s < M$ then there is $\varepsilon' > 0$ such that $s_0 = \rho(L^{\varepsilon'}, L^{\varepsilon'}) > M$.*
2. *Given any $c \in (0,1)$. If M as above is such that $2\lambda^M < c(1-\lambda^{\eta_*})$, then there is $c' \in (0,c)$ such that $t_0 = \rho(L^{\varepsilon'}, L^{c'\varepsilon'})$ and $R^{t_0}_x(L_{\varepsilon'}) \subset^s L_{c'\varepsilon'}$.*

Proof. Item 1: Fix some $M > 0$ such that $R^s_x(L) \cap L = \emptyset$ for all $s \leqslant M$. Take $\varepsilon \in (0, \varepsilon_0)$, set $s_0 = \rho(L^\varepsilon, L^\varepsilon) > 0$ and assume $s_0 < M$. By hypothesis, $R^{s_0}_x(L) \cap L = \emptyset$; then there is $\varepsilon_1 > 0$ such that $R^{s_0}_x(L^{\varepsilon_1}) \cap L^{\varepsilon_1} = \emptyset$. Notice that $s_1 = \rho(L^{\varepsilon_1}, L^{\varepsilon_1}) > s_0 + \eta_* > s_0$. Repeating this argument a finite number of times we obtain some $\varepsilon' := \varepsilon_k > 0$ and $\rho(L^{\varepsilon'}, L^{\varepsilon'}) > M$ which is what we wanted to prove.

Item 2: Take any $c \in (0,1)$. Since $2\lambda^M < c(1-\lambda^{\eta_*})$ there is $c_1 > 0$ such that $c_1 + \frac{2\lambda^M}{1-\lambda^{\eta_*}} < c$. As in the previous item, let $s_0 = \rho(L^{\varepsilon'}, L^{\varepsilon'}) > M$. We shall construct two sequences of numbers inductively, say

$\{c_i, s_i\}_{i\in\mathbb{N}}$. c_1 is already given. Define for $i \geqslant 1$: $s_i = \rho(L^{\varepsilon'}, L^{c_i\varepsilon'})$ and for $i \geqslant 2$: $c_i = \inf\{b > 0 | R_x^{s_{i-1}}(L^\varepsilon) \subset^s L^{b\varepsilon}\}$.

Let $I = \min\{i \in \mathbb{N} | s_i = s_{i+1}\}$ (or equivalently $c_{i+1} \leqslant c_i$), with the convention $I = \infty$ if the set is empty. Observe that for $i < I$ we have $c_{i+1} > c_i$ and also $s_{i+1} < s_i$; since at the time we consider a wider box arround L it is possible we catch some previous returns to it. Notice $s_1 \geqslant s_0$ since $c_1 < c < 1$. On the other hand, for $i < I$, we have $c_i \leqslant 2\lambda^{s_{i-1}} + c_{i-1}$, that is the maximal stable width of the return plus the width of the $\varepsilon' c_{i-1}$-box arround L. Hence

$$c_i \leqslant 2\varepsilon \sum_{j=1}^{i-1} \lambda^{s_j} + c_1.$$

Inductively, for all $i \in \{j, \dots, I\}$ we have that $s_j \geqslant s_0$ and $c_j < c$. We claim that $c_i < c$ and hence, $s_i \geqslant s_0$ since $L^{c_i\varepsilon'} \subset L^{\varepsilon'}$. Since we have $s_j - s_{j+1} > 0$, then $s_j - s_{j+1} > \eta_*$, and therefore, $s_{i-r} \geqslant s_0 + (r-1)\eta_*$ for $r \in \{1, \dots, i-1\}$. Hence,

$$c_i \leqslant 2\varepsilon \sum_{r=0}^{i-2} \lambda^{s_0 + r\eta_*} + c_1 < \frac{2\varepsilon\lambda^M}{1-\lambda^{\eta_*}} + c_1 < c,$$

that is what we claimed.

Since $s_0 \leqslant s_1$ and $s_i < s_1 - i\eta_*$, there must be some i such that $s_i = s_{i+1}$. Therefore $I < \infty$. Take $c' = c_I$, and $t_0 := s_I = \rho(L^{\varepsilon'}, L^{c'\varepsilon'})$. □

Corollary 5.7. *Denote by* $\hat{L} = \{z \in L | L^{\varepsilon'} \cap R_x^{t_0}(L^{\varepsilon'}) \neq \emptyset\}$, *and set* $\tilde{L} = L \setminus \hat{L}$. *Then* $\tilde{L}$ *is an interval and there is* $c'' > 0$ *such that* $c'' \in (c', c)$ *and* $R_x^{t_1}(\tilde{L}^{\varepsilon'}) \subset^s \tilde{L}^{c''\varepsilon_0}$.

Proof. $\tilde{L}$ is an interval since there are no periodic orbits passing through J^ε. To obtain the second part of the Corollary denote by $d_1 = c'$ (from the previous Lemma) and observe $r_1 = \rho(\tilde{L}^{\varepsilon'}, \tilde{L}_{d_1\varepsilon'}) > t_0 > M$. Then, we can apply the same arguments with the interval $\tilde{L}$ instead of L. □

Lemma 5.8. *If there is no periodic orbit on* J *then for any* $M > 0$ *there is a subinterval* $L \subset J$ *such that* $R_x^s(L) \cap L = \emptyset$ *for* $s \leqslant M$.

Proof. Let us suppose that for $0 < s_0 < M$, the first return of J^ε into itself, happens that $R_x^{s_0}(J) \cap J \neq \emptyset$. Since there is no periodic orbit in J we know one of the borders of J, say J^+, must return inside the box. We call $z \in R_x^{s_0}(J) \cap J$ the farthest point to J^+. Let $w \in J$ such that $R_x^{s_0}(w) = z \in J$.

It is important we verify $R_x^{s_0}([w, J^+]^\varepsilon) \subset J^\varepsilon$. This will guarantee we do not lose any point of the box and so, no point of $\omega(I)$. Let

$L = J \setminus [w, J^+]$ and we seek for the first return of the box on L, that is $s_1 > 0$ such that:

$$R_x^{s_1}(L^\varepsilon) \cap L^\varepsilon \neq \emptyset.$$

Observe now that $R_x^s(L) \cap L = \emptyset$ for $s \leqslant s_0 + \eta_*$.

If now we can say $R_x^s(L) \cap L = \emptyset$ for $s < M$ we are done. However, if it does not happen we repeat the above construction on L, obtaining another subinterval $L' \subset L \subset J$ for which

$$R_x^s(L') \cap L' = \emptyset;\ \forall s \leqslant s_2 + \eta_*.$$

Since $s_2 \geqslant s_1 + \eta_* \geqslant s_0 + 2\eta_*$ we are sure to reach M in finite steps. Observe the way we cut the interval J to obtain L let us guarantee any orbit of J is traced by one of L. □

Now, we can give a proof for Proposition 5.5.

Proof of Proposition 5.5. Fix some $c < 1$ and take $M > 0$ big enough such that $2\lambda^M < (1 - \lambda^{\eta_*})$. Using Lemma 5.8 set $L \subset J$ such that $R_x^s(L) \cap L = \emptyset$ for all $s \leqslant M$ and $\omega(L) = \omega(J)$. Now, Lemma 5.6 says there exists $\varepsilon_0 > 0$ such that $\rho(L^{\varepsilon_0}, L^{\varepsilon_0}) > M$ and there exists $0 < c' < c$ such that $R_x^{t_0}(L^{\varepsilon_0}) \subset^s L^{c'\varepsilon_0}$ where $t_0 = \rho(L^{\varepsilon_0}, L^{c'\varepsilon_0})$.

Moreover, on Corollary 5.7 we defined the set

$$\hat{L} = \{z \in L | L^\varepsilon \cap R_x^{t_0}(L^\varepsilon) \neq \emptyset\}.$$

Denote by $\tilde{L} = L \setminus \hat{L}$; and so, there is $c'' > 0$ such that $c' < c'' < c$ and $R_x^{t_1}(\tilde{L}^{\varepsilon_0}) \subset^s \tilde{L}^{c''\varepsilon_0}$, where $t_1 = \rho(\tilde{L}^{\varepsilon_0}, \tilde{L}^{c''\varepsilon_0})$.

Take $\eta = c''\varepsilon_0$. Since $c' < c'' < c$ we know that $L^{c'\varepsilon_0} \subset L^{c''\varepsilon_0} \subset L^{\varepsilon_0}$ and then $\rho(L^{\varepsilon_0}, L^{c'\varepsilon_0}) = \rho(L^{\varepsilon_0}, L^{c''\varepsilon_0}) = \rho(L^\eta, L^\eta)$. Hence we have found that

1. $R_x^{t_0}(L^\eta) \subset^s L^\eta$ and $R_x^{t_0}(L^\eta) \cap \tilde{L}^\eta = \emptyset$, for $t_0 = \rho(L^\eta, L^\eta)$.
2. $R_x^{t_1}(\tilde{L}^\eta) \subset^s \tilde{L}^\eta$, where $t_1 = \rho(\tilde{L}^\eta, \tilde{L}^\eta)$.

That is what we wanted to prove to conclude Proposition 5.5 □

Now, we are ready to construct an irrational torus and then finish the proof in case I. Let $D_0 := L^\eta$ and $t_0 = \rho(D_0, D_0)$, given on Proposition 5.5. Recall $D_0 \subset N_x \subset M$ for some $x \in \Lambda^+$. Moreover, we can think there is Σ, an extension of N_x where $R_x^{t_0}(D_0) \subset \Sigma$. Define $P_0 : D_0 \longrightarrow \Sigma$ as $P_0(\cdot) = R_x^{t_0}(\cdot)$. Recall that the orbit of no point of D_0 falls into a singularity. Since $R_x^{t_0}(D_0) \cap D_0 \neq \emptyset$, there is a point $z \in D_0$ such that $P_0(z) \in D_0$. Therefore, we can extend the function $\tau_x^{t_0}(\cdot)$ to $D_1 := D_0 \cup P_0(D_0)$. In this way we obtain an extension of P_0 called $P_1 : D_1 \longrightarrow \Sigma$.



Figure I

By induction, we can repeat this procedure until we find some $\tilde{n} \in \mathbb{N}$ such that

$$D_i \cap \bigcup_{j \leqslant \tilde{n}} S_j \neq \emptyset \tag{5}$$

where $S_1 = \{X^s(y) | y \in D_0 \text{ and } s \in [0, \tau_x^s(y))\}$, and S_j is defined inductively. If $D_i \cap S_j \neq \emptyset$ for $1 < j < \tilde{n}$ then $D_{\tilde{n}-j} \cap S_1 \neq \emptyset$. Take $n = \tilde{n} - j$. We can think D_n is glued smoothly to D_0. In fact, this is possible since there is a function $\tau^* : D_n \cap S_1 \longrightarrow \mathbb{R}^- \cup \{0\}$, the time a point on the intesection spent to flow back to D_0, that is: for $y \in D_n \cap S_1$, $X^{\tau^*(y)}(y) \in D_0$. Moreover, we can extend the domain of τ^* to some of the last $D_n \cup D_{n-1} \cup \ldots$, and glueing this function with the zero function defined on the first boxes of the sequence $D_0 \cup D_1 \cup \ldots$. Observe that $\bigcup_{i \leqslant n} \tau^*(D_i)$ is a connected smooth surface emebded in M. Observe the intersection of D_n and D_0 corresponds to the first return of $\tilde{L}_\eta$ into itself, called t_1 in Proposition 5.5. Hence $D_n \subset^s D_0$, and more important, there is such $\tilde{n}$ in (5) and this process of extension stops.

By now, we restrict our attention to D_0: D_0 splits into three rectangles, $A \cup C \cup B$ defined as $A = (R_x^{t_0})^{-1}(D_0 \cap D_1)$, $B = (R_x^{t_1})^{-1}(D_0 \cap D_n)$ and C the region in D_0 in between A and B. Denote by $G : D_0 \longrightarrow D_0$ the first return map of the flow to D_0 and notice that $G(A) = D_0 \cap D_1$ and $G(B) = D_0 \cap D_n$. Moreover, we notice also that $R_x^{t_0}(C) \subset D_1$ and $R_x^{t_1}(C) \subset D_n$. Denote by $\tilde{D}_0 := D_0 \cup R_x^{t_0}(C) \cup R_x^{t_1}(C)$, and consider $\tilde{G} : \tilde{D}_0 \longrightarrow \tilde{D}_0$, the corresponding first return map. Observe that still $\tilde{G}(A) = D_0 \cap D_1$ and $\tilde{G}(B) = D_0 \cap D_n$. Also $\tilde{G}(C) = R_x^{t_0}(C)$ and $\tilde{G}(R_x^{t_0}(C)) = R_x^{t_1}(C)$; the only missing part is the image of $R_x^{t_1}(C)$. It is not difficult to see that the orbit of points in C pass through D_0 near the inner border points of $\tilde{G}(A)$ and $\tilde{G}(B)$ (See Figure I)

Lemma 5.9. *If there there is no periodic orbit passing through D_0 then $\tilde{G}|_{R_x^{t_1}(C)}$ is continous.*

Proof. If it is not the case, the rectangle C splits into at least two subrectangles, call C_A and C_B those shareing part of its boundary with A and B, respectively. Observe $\tilde{G}(R_x^{t_1}(C_A)) = G(C_A)$ and is connected to $\tilde{G}(B)$, and $\tilde{G}(R_x^{t_1}(C_B)) = G(C_B)$ and is connected to $\tilde{G}(A)$. Note that in both cases the other stable boundary must reach the stable boundary of D_0. In fact, if not, we are missing some points of D_0 that actually passes in the formed gap, and this is not possible since $\tilde{G}$ is a true first return map to $\tilde{D}_0$.

It is not difficult to see that in this setting either $\pi \circ G|_{A\cup B}: A\cup B \longrightarrow J$ or $\pi \circ G|_{C_A\cup C_B}: C_A \cup C_B \longrightarrow J$ is onto. Therefore there exists $z \in J$ that $G(z) \in W_\eta^s(z)$, and hence we obtain a periodic orbit in D_0 which is a contradiction. Hence $\tilde{G}|_{R_x^{t_1}(C)}$ is continous and so, $G|_C$ is too. □

Therefore, the domain of P_n, the set $D_0 \cup \ldots \cup D_n$, is an embedded ring. It is not difficult to verify that $P_n^m(D_0 \cup \ldots \cup D_n) \to \tilde{S}$ as $m \to \infty$ where $\tilde{S}$ is an invariant circle. Finally note that the saturated of $\tilde{S}$ by the flow, $\mathcal{T} = \{X^t(y) | y \in \tilde{S} \text{ and } t \in \mathbb{R}\}$ is an embedded torus on M; hence $w(I) \subset w(J) = \mathcal{T}$. Moreover, $P_n|_{\tilde{S}}$ corresponds to the Poincaréś first return map of X^t to $\tilde{S}$. Since P_n is a diffeomorphism of the circle without periodic points, it is conjugated to an irrational rotation (See [5]), Therefore, the restriction of X^t to $\mathcal{T}$ is conjugated to a linear irrational flow on the torus. Here we conclude the proof of case I.

Now we deal with Case II, that is: $J_s^\varepsilon \cap [R_x^{t-s}(J_s)]^\varepsilon = \emptyset$, for all $t, s \in \mathcal{P}$; but first, we shall exploit this disjointness property to measure the area of the strip defined by the family $\{J_s\}$.

Remark 3. For any set $A \subset N_x \subset M$, define

$$S(A) = \{X^s(z) | z \in A,\ s \in \left[-\frac{\eta_*}{2}, \frac{\eta_*}{2}\right]\}.$$

There is a constant $K > 0$ only depending on $||X||_{C^1}$ such that if $A = J^\varepsilon$, for some interval J and $\varepsilon > 0$ then $S(J) := S(J^\varepsilon)$ is such that $K\text{Vol}(S(J)) \geqslant |J|$.

Lemma 5.10. *For any $s \in \mathcal{P}$ we have that:*

$$\int_0^\infty |R_{x_s}^t(J_s)| dt < +\infty,$$

and hence, $|R_{x_s}^t(J_s)| \to 0$ as $t \to \infty$.

Proof. Fix some $s \in \mathcal{P}$ and denote $J := J_s$ and $x := x_s$. Notice that for all $l, t \in \mathcal{P}$ such that $|l - t| \geqslant \eta_*$ we have:

$$S(R_x^l(J)) \cap S(R_x^t(J)) = \emptyset. \tag{6}$$

Let us calculate $\int_0^\infty |R_x^t(J)|dt$. For that, observe $\int_{\mathcal{P}} |R_x^t(J)|dt \leqslant K \operatorname{Vol}(M)$. In fact, let $\mathcal{Q}$ be a partition of $\mathcal{P}$ of elements of diameter less than η_*. Remark 3, together with the disjointness property on (6) imply:

$$\int_{\mathcal{P}} |R_x^t(J)|dt \leqslant \sum_{Q\in\mathcal{Q}} \int_Q |R_x^t(J)|dt \leqslant K \sum_{Q\in\mathcal{Q}} \operatorname{Vol}(S(R_x^{t_Q}(J))) \leqslant K \operatorname{Vol}(M)$$

where $t_Q \in Q \in \mathcal{Q}$.

Now arrange the elements of $\mathcal{Q}$ in an ascending way, say $\mathcal{P} = \{[a_i, b_i]\}$, $i \in \mathbb{N}$, such that $b_{i-1} < a_1$ and $(b_{i-1}, a_i) \cap Q = \emptyset$ for any $Q \in \mathcal{Q}$ and $i \in \mathbb{N}$. Notice that there exists $\tilde{\gamma} > 0$ such that $|L^{a_i - s}|_{E(x_s)}| \geqslant \tilde{\gamma}^{a_i}$ for any $s \in (b_{i-1}, a_i)$; otherwise, there must be a point of $\mathcal{P}$ between b_{i-1} and a_i, which is impossible. Using the fact that

$$|L^{a_i-s}|_{E(x)}||L^{a_i-s}|_{F(x)}|^{-1} \leqslant \lambda^{a_i-s}$$

we obtain that $\forall t \in (0, a_i - b_{i-1})$:

$$|L^{-t}|_{F(x_{a_i})}| \geqslant (\tilde{\gamma}^{-1}\lambda)^t.$$

By a similar argument to the one we use in the proof of Lemma 5.1 there is $\mu \in (0,1)$ such that $|D_z R^{-t}_{x_{a_i}}|_{\tilde{F}(z)}| < \mu^t$, for any $t \in (0, a_i - b_{i-1})$, $z \in R_x^{a_i}(J)$, where $\tilde{F}(z) = T_z(R^{a_i}_{x_{a_i}}(J))$.

Hence,

$$|R^{-t}_{x_{a_i}}(R_x^{a_i}(J))| = \int_{R_x^{a_i}(J)} |D_z(R^{-t}_{x_{a_i}})|_{\tilde{F}}|dz \leqslant \mu^t |R_x^{a_i}(J)|.$$

Therefore,

$$\int_0^{a_i - b_{i-1}} |R_x^{a_i - t}(J)|dt \leqslant |R_x^{a_i}(J)| \int_0^{a_i - b_{i-1}} \mu^t dt \leqslant \frac{|R_x^{a_i}(J)|}{-\log \mu}.$$

A similar argument of disjointness used to bound the integral over the hyperbolic times implies $\sum_i |R_x^{a_{i+1}}(J)| < K \operatorname{Vol}(M)$. Therefore,

$$\int_{\mathbb{R}^+} |R_x^t(J)| \leqslant K \operatorname{Vol}(M)\big(1 - \frac{1}{\log \mu}\big).$$

Finally, the function $\log |R_x^t(J)|$ is uniformly continous on t; hence $|R_x^t(J)| \to 0$ as $t \to \infty$. □

The following lemma measures how much the orbits of the flow X^t differ from the orbits by R^t.

Lemma 5.11. *There exists a constant $C > 0$ such that if an interval $I \subset D^t_x$ for all $t \geqslant 0$ then, for any $y, w \in R^r_x(I)$, and $r, s \geqslant 0$ we have the following inequality:*

$$|\tau^s_{x_r}(y) - \tau^s_{x_r}(w)| \leqslant C \int_r^\infty |R^t_x(I)|dt$$

Proof. Recall that:

$$\tau^s_{x_r}(y) \;=\; \sum_{i=0}^{[s]-1} \tau^1_{x_{i+r}}(R^i_{x_r}(y)) + \tau^{s-[s]}_{x_{[s]+r}}(R^{[s]}_{x_r}(y)),$$

then

$$\begin{aligned} |\tau^s_{x_r}(y) - \tau^s_{x_r}(w)| \;\leqslant\; & \sum_{i=0}^{[s]-1} |\tau^1_{x_{i+r}}(R^i_{x_r}(y)) - \tau^1_{x_{i+r}}(R^i_{x_r}(w))| \\ & + |\tau^{s-[s]}_{x_{[s]+r}}(R^{[s]}_{x_r}(y)) - \tau^{s-[s]}_{x_{[s]+r}}(R^{[s]}_{x_r}(w))| \end{aligned}$$

Now, the function τ^1 is uniformly Lipschitz with some constant $C > 0$. Hence,

$$\leqslant C \sum_{i=0}^{[s]-1} |R^i_{x_r}(I_r)| + |R^{[s]}_{x_r}(I_r)| \leqslant C \int_r^\infty |R^t_{x_0}(I_0)|dt$$

and then the Lemma follows. □

Let us assume there are $t_0 < t_1$, and $t_0, t_1 \in \mathcal{P}$ such that $J^\varepsilon_{t_0} \cap J^\varepsilon_{t_1} \neq \emptyset$. In this case we shall obtain a sequence of times on which this intesection is at a small enough scale that allow us to control the limit. For that, call $Z = J^\varepsilon_{t_0} \cap J^\varepsilon_{t_1}$, and observe that $R^{t_1-t_0}_{x_0}(Z) \subset J^\varepsilon_{t_1}$, since $R^{t_1-t_0}_{x_0}(J^\varepsilon_{t_0}) \subset J^\varepsilon_{t_1}$. This implies the existence of certain $t^* > 0$ such that $R^{t^*}_{x_{t_1}}(J^\varepsilon_{t_1}) \cap J^\varepsilon_{t_1} \neq \emptyset$. Remember it is necessary to fix a destination section any time we define some R^s_y. In this case we select the time t^* in such a way that x_{t^*} lie in some smooth extension of the section over x_{t_1}. Moreover, we can take this extension in such a way the tangent space on x_{t^*} is exactly $\mathcal{N}_{x^*_t}$.

Let $t_2 = t_1 + t^*$ and set $C(t_1) := J^\varepsilon_{t_1}$. Notice that $C(t_1) \subset D^{t^*}_{x_{t_1}}$, then $R^{t_2-t_1}_{x_{t_1}}(C(t_1)) \cap C(t_1) \neq \emptyset$, by construction. Now set $C(t_2) := R^{t_2-t_1}_{x_{t_1}}(C(t_1))$. Inductively, take $t_{i+1} = t_i + \tilde{t^*}$, such that $R^{t_{i+1}-t_i}_{x_{t_i}}(C(t_i)) \cap C(t_i) \neq \emptyset$, and denote $C(t_{i+1}) := R^{t_{i+1}-t_i}_{x_{t_i}}(C(t_i))$ [see Figure II]. Notice that $|t_{i+1} - t_i| > \eta_*$, since any point must flow away certain ammount of time before it returns into a neigbourhood of it. For any



Figure II.

$j, i \in \mathbb{N}$ such that $j < i$, Lemma 5.11 implies that for each $s > 0$ and $y \in R^{t_i-t_j}_{x_{t_j}}(J_{t_j})$ we have that

$$|\tau^s_{x_{t_j}}(y) - s| \leqslant C \int_{t_i-t_j}^{\infty} |R^t_{x_{t_j}}(J_{t_j})| dt.$$

Hence, for j large enough we have that for all $y \in R^{t_i-t_j}_{x_{t_j}}(J_{t_j})$

$$|\tau^s_{x_{t_j}}(y) - s| \leqslant \frac{\eta_*}{3}. \tag{7}$$

Fix $j \in \mathbb{N}$ such that also verifies that $\frac{C\varepsilon_0\lambda^j}{-\log(\lambda)} < \frac{\eta_*}{3}$. We may suppose $x_{t_i} \to z \in M$, as $i \to \infty$. Hence, Lemma 5.10 implies that $R^{t_i-t_j}_{x_{t_j}}(J_{t_j}) = J_{t_i} \to z$, as $i \to \infty$. Moreover, $R^{t_i-t_j}_{x_{t_j}}(C(t_j)) \to z$, as $i \to \infty$. However the inequality on (7) only implies that

$$\lim_{i\to\infty} X^{t_i-t_j}(y) \in \{X^t(z) | t \in [-\frac{\eta_*}{3}, \frac{\eta_*}{3}]\} \tag{8}$$

for any $y \in J_{t_j}$. Moreover (8) is valid also for any $y \in R^{t_j-t_1}_{x_{t_j}}(J^{\varepsilon_0}_{t_1})$ since

$$|\tau^s_{x_j}(y) - s| \leqslant |\tau^s_{x_j}(y) - \tau^s_{x_j}(w)| + |\tau^s_{x_j}(w) - s|$$

for any $y \in R^{t_j-t_1}_{x_{t_1}}(W^s_{\varepsilon_0}(\hat{w}))$, where $\hat{w} \in J_{t_1}$ and $R^{t_j-t_1}_{x_{t_1}}(\hat{w}) = w$.

On the other hand, we know that $C(t_j) \cap C(t_{j+1}) \neq \emptyset$, that is

$$R^{t_j-t_1}_{t_1}(J^{\varepsilon}_{t_1}) \cap R^{t_{j+1}-t_1}_{t_1}(J^{\varepsilon}_{t_1}) \neq \emptyset.$$

This implies that there exists $\hat{w}_1, \hat{w}_2 \in J_{t_1}$ such that

$$R^{t^*}_{x_{t_1}}(W^s_{\varepsilon}(\hat{w}_2)) \subset W^s_{\varepsilon_0}(\hat{w}_1),$$

since $W^{cs}_{\varepsilon_0}(\hat{w}_k) = W^s_{\varepsilon_0}(\hat{w}_k)$, for both, $k = 1, 2$; and hence, they are coherent because the center stable manifolds are dynamically defined.

Let $\hat{w}_3 = R^{t^*}_{x_{t_1}}(\hat{w}_2) \in W^s_{\varepsilon_0}(\hat{w}_1)$. Denote by $w_k = R^{t_j - t_1}_{x_{t_1}}(\hat{w}_k) \in C(t_j)$, for $k = 2, 3$. Notice there is $s > \eta_*$ such that $X^s(w_2) = w_3$. Hence, $\lim_{i\to\infty} X^{t_i - t_j}(w_3) = X^l(z)$, where $|l| < \frac{\eta_*}{3}$. On the other hand

$$\lim_{i\to\infty} X^{t_i - t_j}(X^s(w_2)) = X^{l'}(X^s(z)) \in \{X^t(z) | t \in [-\frac{\eta_*}{3}, \frac{\eta_*}{3}]\},$$

since $s > \frac{\eta_*}{3}$ it implies that z is periodic.

Finaly, if for any $t, s \in \mathcal{P}$ happens that $J^\varepsilon_s \cap J^\varepsilon_t = \emptyset$, by a similar calculation as in the proof the Lemma 5.10, we obtain

$$\int_{\mathcal{P}} |J_s| ds < +\infty \text{ and hence, } |J_s| \to 0 \text{ as } s \to \infty.$$

Therefore, if we prove there exists a sequence of times $\{s_i\}_{i\in\mathbb{N}}$ such that $|J_{s_i}| = \delta$, we get a contradiction. Now, like in Schwartz' proof of Denjoy's Lemma we have that for all J_s maximal there exists $t > 0$ such that $|R^t_{x_s}(J_s)| = \delta$. It is only left to verify that $t + s$ is a hyperbolic time, However, if it is not the case, like in the proof of Lemma 5.10 the interval must increase its length contradicting the fact of the length being δ. Here we finish the proof of Proposition 5.2. □

Once we have proved Proposition 5.2, it is not difficult to verify the following Corollary, regarding that only can exist a finite number of irrational tori on an admissible neighborhood of Λ (see Section 4).

Corollary 5.12. *If we assume that Λ has a dominated splitting and it does not contain an irrational torus, there is $\delta_0 > 0$ and a neighborhood V of Λ where for any (δ, E)-interval, $I \subset V$ with $\delta < \delta_0$, we have that $\omega(I) \subset Per(X^t|_V)$.*

All these properties are still valid for (δ, F)-intervals, considering the flow-time backwards.

5.2. **Dynamically defined local invariant manifolds.** Assume Λ is a compact invariant set of X^t with a dominated splitting, not containing an irrational torus. As we see in Corollary 5.12, there is $\delta_0 > 0$ such that any (δ, E)-interval, $\delta < \delta_0$ has $\omega(I) \subset \mathrm{Per}(X^t|_V)$ for some appropriate admissible neighborhood V.

Lemma 5.13. *There exists $\delta_3 \in (0, \delta_0)$ such that if any periodic point $p \in \Lambda$ has one of its connected components of $W^u(p) - \{p\}$, say W, such that $|R^t_p(W)| \leqslant \delta_3$ for all $t \geqslant 0$, then the other endpoint which is not p is periodic and it is not hyperbolic of saddle type. Moreover, it is a sink or a non hyperbolic periodic point.*

Proof. Let $U_1 \supset \Lambda$ be a neigbourhood where it is defined an extension of the invariant cone field of Λ. Let $\delta_3 < \delta_0$ be a positive number such that any ball of radius δ_3 centered on a point of Λ is contained in U. Take a periodic point p and W as in the hypothesis of the Lemma. There is a number $t_p > 0$ such that $R_p^{t_p}(W) = W$. In fact, t_p is the period of p or twice. Notice that W is a (δ_3, E)-interval. Since $q \in V_1$ it is well defined the direction $F(q) = T_q(W)$ and $|L^{t_q}|_{F(q)}| \leqslant 1$ and hence, by the domination condition we have that $|L^{t_q}|_{E(q)}| \leqslant \lambda < 1$. Hence, if p is hyperbolic it must be a sink. □

The following lemma is a fundamental application of Proposition 5.2.

Lemma 5.14. *For all $\varepsilon \in (0, \delta_0)$ there is $\gamma = \gamma(\varepsilon) > 0$ such that:*

1. *For all $t > 0$ we have that*
   a) $R_x^{-t}(W_\gamma^{cu}(x)) \subset W_\varepsilon^{cu}(x_{-t})$
   b) $R_x^{t}(W_\gamma^{cs}(x)) \subset W_\varepsilon^{cs}(x_{t})$
2. *Given $\tilde{\gamma} \in (0, \gamma(\delta_0))$ we have*
   a) $|R_x^{-t}(W_{\tilde{\gamma}}^{cu}(x))| \to 0$ *as* $t \to \infty$, *or* $x \in W^u(p)$ *for some periodic point p and $p \in W_{\tilde{\gamma}}^{cu}(x)$.*
   b) $|R_x^{t}(W_{\tilde{\gamma}}^{cs}(x))| \to 0$ *as* $t \to \infty$, *or* $x \in W^s(p)$ *for some periodic point p and $p \in W_{\tilde{\gamma}}^{cs}(x)$.*

Proof. We shall only prove the statement for central unstable manifold since the same argument works for the central stable manifolds. For item 1 take δ_0 from Proposition 5.2 and let $\varepsilon < \delta_0$. Recall that Corollary 4.3 assert there is $0 < \delta < \delta_0$ such that if: $|R^{-s}(W_\varepsilon^{cu}(x))| < \delta$ for $s \in [0, t]$ then $R^{-s}(W_\varepsilon^{cu}(x)) \subset W_\varepsilon^{cu}(x_{-s})$ for $s \in [0, t]$. Arguing by contradiction, assume the statement of item 1 does not holds. Then there exist a sequence of numbers $\gamma_n \to 0$, points $x^n \in \Lambda$ and times $m_n \to \infty$ such that $|R^{-m_n}(W_{\gamma_n}^{cu}(x^n))| > \delta$. We should be more precise on how this bound is exeeded; that is, $|R^{-s}(W_{\gamma_n}^{cu}(x^n))| \leqslant \delta$ for $s \in (0, m_n)$, and $|R^{-m_n}(W_{\gamma_n}^{cu}(x^n))| = \delta$. Let us write $I_n = R^{-m_n}(W_{\gamma_n}^{cu}(x^n))$. At least for a subsequence, we can assume that $y^n := X^{-m_n}(x^n) \to z \in \Lambda$ and also I_n converges to some interval I transversal to the E direction. Notice $|I| \leqslant \delta$ and $z \in \mathrm{cl}(I)$. By construction: $I \subset \mathrm{Dom}(R^t) = V_t$ for all $t \geqslant 0$ and $|R^t(I)| \leqslant \delta_0$ for all $t \geqslant 0$. Hence is a (δ, E)-interval. Now, since $z \in \mathrm{cl}(I)$ we know $\omega(z) = p \in \mathrm{Per}(X^t)$ as Proposition 5.2 says. On the other hand, p is hyperbolic since $p \in \Lambda$. Therefore $z \in W^s(p)$. Observe that one connected component of $W^u(p) \smallsetminus \{p\}$ has length less than δ. In fact, $W^u(p)$ is the limit of the intervals $|I_n| \leqslant \delta$ and $z \in \mathrm{cl}(I)$. Hence, the previous lemma implies the other end point must be a periodic point and not an hyperbolic saddle. It may happen that $I_n \cap W^s(p) \neq \emptyset$ for $n >> 1$. In this case we

get a contradiction with the Inclination Lemma, since this intersection is transversal and $R^{m_n}(I_n) = R^{-m_n}(R^{m_n}(W^{cu}_{\gamma_n}(x^n))) \to 0$ as $n \to \infty$. Hence, $I_n \cap W^s(p) = \emptyset$ for n arbitrarily large. In this case, $\omega(y^n) \subset \Lambda$ is the other endpont of the connected component of $W^u(p) \setminus \{p\}$ of length less than δ. Then $\omega(y^n)$ is a periodic sink or a non-hyperblic periodic point. This is a contradiction since all periodic points in Λ are hyperbolic of saddle type, concluding the proof of item 1.

To prove item 2: Take $\delta < \delta(\gamma_0)$, and let $x \in \Lambda$ be a point such that there exists $\eta > 0$ and a sequence of times $t_k \to \infty$ in such a way $|R^{-t_k}(W^{cu}_\gamma(x))| \geqslant \eta$. Last item guarantees that $\eta < \delta_0$, since $|R^{-t}(W^{cu}_\gamma(x))| \leqslant \delta_0$ for any $t \geqslant 0$. As in the previous case, we can obtain a sequence of intervals $I_k := R^{-t_k}(W^{cu}_\gamma(x))$ converging to a (η, E)-interval I, where $z \in \mathrm{cl}(I)$, and $z \in \Lambda$. Hence $z \in W^s(p)$ for some periodic point $p \in \Lambda$ of saddle type. Observe I is transversal to $W^s(p)$, even in the case $z \notin \mathrm{Int}(I)$. Avoiding a contradiction with the Inclination Lemma we have that $|W^u(p)| \leqslant \eta \leqslant \delta_0$. If $z \in \mathrm{Int}(I)$, as $y_k := X^{t_k}(x) \to z$ we conclude $y_k \in W^s(p)$. Otherwise we get a contradiction since both connected components of $W^u(p) \setminus \{p\}$ have length less than δ_0, and hence $\omega(x)$ must be a sink or a non-hyperbolic periodic point. Suppose $z \neq p$. Then for $k >> 1$ all $y_k = X^{-t_k}(x)$ are contained in some fundamental neighborhood of $W^s(p)$. This is a contradiction, since $W^s(p)$ do not have self-intersections. Hence $z = p$. Therefore $y_k \in W^s_{\mathrm{loc}}(p)$. Otherwise $\omega(x)$ must be a sink or a nonhyperbolic periodic point. Hence, $x = p$, proving item 2 in case $z \in \mathrm{int}(I)$.

It is left to verify the case when $z \notin \mathrm{int}(I)$. Observe: $R^{-t_k}(W^{cu}_\gamma(x)) \cap W^s(p) = I_k \cap W^s(p)$ and this intesection is not empty; otherwise, we find a contradiction in a similar way as in the proof of item 1. Combining the following two facts: $|R^t(R^{-t_k}(W^{cu}_\gamma(x)))| \leqslant \delta_0$ for $t \in (0, t_k)$ and $X^{-t_k}(x) \to z$ as $k \to \infty$ we conclude $x \in W^u(p)$. Moreover $x \in W^u_{\delta_0}(p)$. If $p \notin W^{cu}_\gamma(x)$, as $y_k \to p$ when $k \to \infty$ and $|R^{-t_k}(W^{cu}_\gamma(x))|$ do not converges to 0 we have that $R^{-t_k}(W^{cu}_\gamma(x)) \cap W^s(p) \neq \{p\}$ for $k >> 0$, and in particular $I \subset W^s(p)$, a contradiction. Hence $p \in W^{cu}_\gamma(x)$ and we are done. □

6. Proof of Main Lemma

We shall devote this last section to prove the Main Lemma stated on Section 4. For that, we shall first provide non-periodic points of Λ (Λ as in the hypothesis of Main Lemma) with certain two-dimensional boxes in which we have some nice properties of recurrence; this will spent most of our effort. Another important notion are the returns associated to such boxes, and here we prove they behave in the same

way as their conterpart for surface's diffeomorphisms ([21], pag 1007). On the other hand, under the same hypothesis of Main Lemma, we shall obtain more dynamical properties for center-stable and center-unstable manifolds. All these shall allow us to fit into the last calculations of [21] and therefore finish the proof.

Let us begin defining the standard box arround a point in Λ, and the notion of distortion along the cs-direction on such boxes:

Let $x \in \Lambda$ be given. For any two intervals $J \subset W^{cu}_{\gamma}(x)$ and $T \subset W^{cs}_{\gamma}(x)$, both containing the point $(x,0) \in \mathcal{N}_x$, we call the box, or (J,T)-box, arround x the set $B_T(J) = \exp_x(T \times J)$, with boundaries: $\partial^{cs}(B_T(J)) = \exp_x(\partial T \times J)$ and $\partial^{cu}(B_T(J)) = \exp_x(T \times \partial J)$. Observe that if the size of the box, say $|T|+|J|$ is small enough, then $\partial^{cs}(B_T(J))$ is transversal to the E-direction and also $\partial^{cu}(B_T(J))$ is transversal to the F-direction. Also we can assume that for any $y \in \Lambda$, $\partial^{\alpha}(B_T(J)) \cap W^{\alpha}_{\gamma}(y)$ are relative open sets for $\alpha = cs, cu$. If $B_T(J)$ is a box and $x \in T' \subset T$, we say $B_{T'}(J)$ is a subbox of $B_T(J)$ if it is a box and $\partial^{cs}B_{T'}(J) \subset \partial^{cs}B_T(J)$. For simplicity, we shall denote only $B_{\varepsilon}(J)$, when $T \subset W^{cs}_{\varepsilon}(x)$.

Definition 4. A box $B_{\varepsilon}(J)$ has distortion $C > 0$, or precisely distortion along the cs-direction, if for any two intervals $J_1, J_2 \subset B_{\varepsilon}(J)$, both transversal to the E direction and whose end points lie in $\partial^{cs}B_{\varepsilon}(J)$ happens that:

$$C^{-1} \leqslant \frac{l(J_1)}{l(J_2)} \leqslant C.$$

An equivalent way to obtain such distortion is by means of a projection along some foliation close to the E-direction defined on the box. Precisely, let $\mathcal{F}^{cs}$ be the foliation obtained by integrating some C^1 vector field Y defined on $B_{\varepsilon}(J)$ such that $Y(p) \in C^E_a(p)$, for all $p \in \mathrm{int}(B_{\varepsilon}(J))$ and for $p \in \partial^{cu}(B_{\varepsilon}(J))$, $Y(p) \in T_p\partial^{cu}(B_{\varepsilon}(J))$. If we denote by $\Pi = \Pi(J_1, J_2)$ the projection along the leaves of $\mathcal{F}^{cs}$ betweeen the two intervals J_1, J_2, then the box $B_{\varepsilon}(J)$ has distortion C if there exists $C > 0$ such that $C^{-1} \leqslant ||\Pi'|| \leqslant C$.

In [21] they guarantee distortion on a dynamical box if it is provided of some bound on the sum of the lenght of the preimages of the leaves of the foliation on the box. Under the same hypothesis, this is still valid in our context, since the generalized Poincaré transformation, behaves like a diffeomorphism (See Corolary 3.5 of [21]). Let us state the following Corolary to be precise:

Corollary 6.1. *There exists a positive number $\tau(< \delta_2)$ such that if for some $z \in B_\varepsilon(J) \cap \Lambda$ and $n > 0$ we have*

$$||L^t_{E(z_{-t})}(z)|| < \lambda_1^t, \text{ for } 0 \leqslant t \leqslant n$$

and the box $B(n) := B(R_z^{-n}(J(z)))$ satisfies that:

1. *$R^n_{z_{-n}}(B(n)) \subset B_\varepsilon(J)$ and $R^n_z(\partial^{cs} B(n)) \subset \partial^{cs} B_\varepsilon(J)$,*
2. *$R^t_{z_{-n}}(R_z^{-n}(J(z)))$ has diameter less than τ, for $0 \leqslant t \leqslant n$,*

then there exists $C_1 > 0$ such that $B(R_z^{-n}(J(z)))$ has distortion C_1.

Using this Corollary, we can prove more dynamical properties on the central manifolds.

Lemma 6.2. *Assume Λ is transitive and every proper compact invariant subset of Λ is hyperbolic, then either $|R_x^{-t}(W^{cu}_\gamma(x))| \to 0$ as $t \to \infty$ for any $x \in \Lambda$ or the direction F is expanding (that is, $|L^{-t}|F(x)| \to 0$ as $t \to \infty$).*

Proof. Observe Λ is not a periodic orbit of the flow. Suppose there is $x \in \Lambda$ such that $\limsup |R_x^{-t}(W^{cu}_\gamma(x))| > 0$. Lemma 5.14 implies there is a periodic orbit $p \in \Lambda$ such that one of the components of $W^u(p) - \{p\}$ has lenght less than δ_0. Also Lemma 5.13 implies the endpoint of this component $q \neq p$, is a sink or a non-hyperbolic periodic point, and so, $q \notin \Lambda$.

Since Λ is transitive and non-trivial there is $x_0 \in \Lambda \smallsetminus \mathrm{Per}(X^t)$ such that $x_0 \in W^s(p)$. Moreover, there is a small neighborhood of x_0, call it $U \subset \mathcal{N}_p$ divided into two components by $W^{cs}(x_0) \subset W^s(p)$, where of one side, say the "upper" one, have the same ω-limit as q.

Take $\lambda < \lambda_1 < \lambda_2 < \lambda_3 < 1$, and $c > 0$ such that $\lambda \lambda_1^{-1} < \lambda_2$ and $(1+c)\lambda_2 < \lambda_3$. Take $\tilde{\delta} > 0$ such that if I is a (δ, E)-interval on z for $\delta < \tilde{\delta}$, then for all $y \in I$ we have:

$$(1-c)^t \leqslant \frac{|L^t|_{F(y)}|}{|DR^t_y|_{\tilde{F}(y)}|} \leqslant (1+c)^t$$

where $\tilde{F}(y) = T_y I$.

Let $B_\varepsilon(J)$ be a box for x_0 contained in U and even so small such that

$$R^t_{x_0}(B_\varepsilon(J^+)) \cap B_\varepsilon(J^+) = \emptyset, \ \forall t \neq 0$$

and $|R^{-t}_{x_0}(J(y))| \leqslant \delta$, for any $y \in B_\varepsilon(J) \cap \Lambda$ and $t \geqslant 0$; where $\delta + \varepsilon < \eta < \tau$ (τ from Corollary 6.1 and η the size of the box).

For any given $z \in B_\varepsilon(J) \cap \Lambda$ denote by $J^+(z) = J(z) \cap B_\varepsilon(J^+)$ and for it we have:

$$R_z^{-t}(J^+(z)) \cap B_\varepsilon(J^+) = \emptyset, \ \forall t > 0.$$

Let be $B_{\varepsilon_1}(J)$ a subbox for a positive $\varepsilon_1 < \varepsilon/4$. We claim there is $K > 0$ such that for any $z \in B_{\varepsilon_1}(J) \cap \Lambda$:

$$\int_{t \geqslant 0} |R^{-t}(J^+(z))| dt \leqslant K$$

for any $z \in B_{\varepsilon_1}(J) \cap \Lambda$.

Once established this claim, the argument on the proof of Lemma 3.5.2 of [21] concludes the thesis of the Lemma.

Let us prove the claim. Fix an arbitrary point $z \in B_{\varepsilon_1}(J) \cap \Lambda$. And consider the set

$$M_z = \{m \in \mathbb{R}^+ \,||L^t|_{E(z_{-m})}| \leqslant \lambda_2^t, \, \forall t \in [0, m]\}$$

Corollary 6.1 implies that for any $m \in M_z$ the box

$$B(m) = B_{\varepsilon_1}(R_z^{-m}(J^+(z)))$$

has distortion C. In fact, it is true that $R^{m_i}_{z_{-m}}(B(m)) \subset B_\varepsilon(J)$ and $R^{m_i}_{z_{-m}}(\partial^{cu} B(m_i)) \subset \partial^{cu} B_\varepsilon(J)$. Notice also that for $m, n \in M_z$, $m \neq n$, then $B(m) \cap B(n) = \emptyset$. Take $\varepsilon_2 = \varepsilon_1/4$. For each $y \in \Lambda$ consider the three dimensional cube (defined on section 3):

$$S(y) := S(B_{\varepsilon_2}(W^{cu}_{\gamma/2}(y))).$$

Since Λ is compact, it can be covered by a finite number of these cubes:

$$\Lambda \subset \bigcup_{k=0}^{k_0} S(y_k).$$

For each $m \in M_z$, the point z_{-m} belongs to some $S(y_k)$. Denote by $J(m) := B(m) \cap S(y_k)$. If the size of the box η is small enough, $J(m)$ is a non empty interval contained in $B(m)$. Notice there are constants $K_1, K_2 > 0$ only depending on η_* such that

$$|J(m)| \leqslant K_1 |W^{cu}_{\gamma/2}(y_k)| \leqslant K_2 \mathrm{Vol}(S(y_k)).$$

Since we have distortion C on $B(m)$, we have

$$\frac{1}{C} \leqslant \frac{|J(m)|}{|R_z^{-m}(J^+(z))|} \leqslant C$$

and hence,

$$|R_z^{-m}(J^+(z))| \leqslant C|J(m)|.$$

Therefore,

$$\int_{M_z} |J(t)| dt \leqslant K \sum_{k=0}^{k_0} \int_{-\eta_*/2}^{\eta_*/2} |X^t(W^{cu}_{\gamma/2}(y_k))| dt \leqslant K \mathrm{Vol}(M)$$

Now we have to bound the integral in the gaps of M_z. Let $M, m \in M_z$ such that $M > m$ and $M_z \cap (m, M) = \emptyset$ (perhaps $M = \infty$). For

$t \in (m, M)$, we have that $|L^t|_{E(z_{-m-t})}| \geqslant \lambda_1^t$. Then, the dominated splitting condition implies:

$$|L^{-t}|_{F(z_{-m})}| \leqslant (\lambda\lambda_1^{-1})^t \leqslant \lambda_2^t$$

and hence,

$$|DR_y^{-t}|_{\tilde{F}(y)}| \leqslant \lambda_3^t$$

where $\tilde{F}(y) = T_y R_z^{-m}(J^+(z))$. Denote $J_m = R_z^{-m}(J^+(z))$, then

$$\int_m^M |R^t(J_m)|dt \leqslant \frac{1}{1-\lambda_3}|J_m|.$$

Hence,

$$\begin{aligned}\int_{t\geqslant 0} |R^t(J^+(z))|dt &\leqslant \textstyle\int_{M_z} |R^t(J^+(z))|dt + \sum_j \int_{m_j}^{M_j} |R^t(J_{m_j})|dt \\ &\leqslant K\mathrm{Vol}(M) + \tfrac{1}{1-\lambda_3}\textstyle\sum_j |J_{m_j}|\end{aligned}$$

where $\{(m_j, M_j)\}$ denotes all the gaps of M_z. However, $m_j \in M_z$ and hence, the box $B(m_j)$ has distortion C. Hence, we can bound the last sum above $\sum_j |J_{m_j}| \leqslant C\mathrm{Vol}(M)$, proving the claim. □

An analogous result holds also for center stable manifolds, allowing us to conclude that for all $x \in \Lambda$ we have both, as $t \to \infty$:

$$|R_x^{-t}(W_\gamma^{cu}(x))| \to 0 \text{ and } |R_x^t(W_\gamma^{cs}(x))| \to 0.$$

In the sequel, we shall take $\gamma \leqslant \gamma_1$.

Corollary 6.3. *Given $\varepsilon > 0$ there exists some $t_0 \geqslant 0$ such that for any $x \in \Lambda$ and $t \geqslant t_0$ we have:*

$$|R_x^{-t}(W_\gamma^{cu}(x))| \leqslant \varepsilon \text{ and } |R_x^t(W_\gamma^{cs}(x))| \leqslant \varepsilon.$$

6.1. **Returns.** Let $B_\varepsilon(J)$ be a box and suppose $x \in J \cap \Lambda$. Recall that for any $y \in B_\varepsilon(J) \cap \Lambda$ we have defined $J(y) = W_\gamma^{cu}(y) \cap B_\varepsilon(J)$. Given a neighborhood V, we consider the set of points whose entire orbit stays on V and they are asymptotic to Λ

$$\tilde{\Lambda} = \{y \in V | \mathrm{dist}(\Phi^t(x), \Lambda) \to 0 \text{ as } t \to \pm\infty\}$$

Notice that $\tilde{\Lambda}$ also has dominated splitting since it is contained in V. Although it is not compact, the arguments in Lemmas 5.14 and 6.2 apply, and so, there are central stable and central unstable manifolds on points of $\tilde{\Lambda}$ with the corresponding dynamical properties.

Definition 5. For $\delta > 0$, we say the box $B_\varepsilon(J)$ is δ-adapted if for every $y \in B_\varepsilon(J) \cap \Lambda$ the following conditions are satisfied:

1. $|R_x^{-t}(J(y))| \leqslant \delta$ for all $t \geqslant 0$
2. $R_x^{-t}(J(y)) \cap B_\varepsilon(J) = \emptyset$ or $R_x^{-t}(J(y)) \subset B_\varepsilon(J)$, for all $t \geqslant 0$.

Before we prove the existence of adapted boxes for certain points on Λ, we need to recognize some configuration that implies the presence of periodic orbits in a box, as the following lemma says.

Lemma 6.4. *Let $J \subset W^{cu}_\gamma(x)$ be an interval, for some $x \in \Lambda$ and assume $W^{cu}_\gamma(x)$ is dynamically defined. Suppose there is $t > 0$ such that $R^{-t}_x(J) \cap J \neq \emptyset$. Then, the α-limit is a periodic orbit. Hence x is periodic.*

This lemma is also true replacing *cu* by *cs*, α-limit by ω-limit and takeing positive iterations.

Proof. Suppose there is $t > 0$ such that $R^t_x(J) \cap J \neq \emptyset$, then $x_t \in W^{cu}_\gamma(x)$. Therefore there is a sequence $t_k \to \infty$ such that $x_{t_k} \in W^{cu}_\gamma(x)$ and that $x_{t_k} \to x$ as $k \to \infty$. Since $W^{cu}_\gamma(x)$ is dynamically defined, for k big enough: $R^{-t_k}_x(W^{cu}_\gamma(x)) \subset W^{cu}_\gamma(x)$, and hence $\alpha(x)$ is a periodic orbit. Moreover x is periodic. □

Lemma 6.5. *For any δ small and any point $x \in \Lambda \setminus Per(X^t)$ there exists δ-adapted boxes associated to x.*

Proof. Let $x \in \Lambda$ be a non-periodic point. Notice that condition 1 of the definition is easely satisfied if the height of the box is small enough. So, we are only interested on condition 2. In spite of the previous Lemma, there is an interval $J \subset W^{cu}_\gamma(x)$ for which $\Phi^{-t}(J) \cap J = \emptyset$, for any $t > 0$. Given some order relation on J we can split J into two intervals acording to x, say $J^+ = \{y \in J | y > x\}$ and $J^- = \{y \in J | y < x\}$. Now we define

$$A^+ = \{y \in J^+ | \exists z \in \tilde{\Lambda}, W^{cs}_\gamma(z) \cap J^+ = \{y\}\}$$

and the corresponding set for J^-. Let $\{\varepsilon_k > 0\}_{k\in\mathbb{N}}$ be a decreasing sequence of numbers converging to 0 and let B_{ε_k} be a sequence of boxes arround J such that $B_{\varepsilon_{k+1}} \subset B_{\varepsilon_k}$ and $\bigcap_k B_{\varepsilon_k} = J$. For any $y \in A^+$ and any positive integer k we denote by $U(y, \varepsilon_k)$ the connected component of $W^{cu}_{2\gamma}(z) \cap B_{\varepsilon_k}(J)$ which contains y. Consider also

$$I(y, \varepsilon_k) = \{(z, n) | z \in B_{\varepsilon_k}(J) \cap \Lambda, n > 0 \text{ and such that } R^{-n}_x(J(z)) \cap U(y, \varepsilon_k) \neq \emptyset\}$$

If there exist points $y^+ \in J^+$ and $y^- \in J^-$ and there is an integer k with $\#I(y^\pm, \varepsilon_k) = 0$ then the box $B_{\varepsilon_k}(J')$ arround $J' = (y^-, y^+)$ is δ-adapted. Notice $\partial^{cs} B_{\varepsilon_k}(J') = U(y^-, \varepsilon_k) \cup U(y^+, \varepsilon_k)$. Actually, we only need $\#I(y, \varepsilon_k) < \infty$ for some $y \in J$ since then, for some $k' > k$ we obtain $\#I(y, \varepsilon'_k) = 0$. So, we shall assume $\#I(y, \varepsilon_k) = \infty$ for any k and $y \in A^+ \cup A^-$, for instance take $y \in A^+$ and we seek for a contradiction. If the set $K := \{n | \exists (z, n) \in I(y, \varepsilon_k)\}$ is bounded for some k then it is

bounded for the same y and any $k' > k$. In fact $I(y, \varepsilon_{k'}) \subset I(y, \varepsilon_k)$. Hence, for some k' big enough one must have either $I(y, \varepsilon_k) = \emptyset$ or $R_x^{-t}(J) \cap J \neq \emptyset$ for some $t > 0$, which is a contradiction. Therefore, the set K is unbounded for any (y, ε_k). As a consequence, for any $y \in A^+$ there are sequences $z_n \in B_{\varepsilon_n}(J)$ and $m_n \to \infty$ such that

$$R_x^{-m_n}(J(z_n)) \cap U(y, \varepsilon_n) \neq \emptyset.$$

Observe $|R_x^{-m_n}(J(z_n))| \to 0$ as $n \to \infty$; otherwise there is a limit point z of the sequence $\{z_n\}$ whose backward orbit shadows any z_n until time $-m_n$ and shadows x for the entire past too. Therefore, $\limsup_{t\to\infty} |R_x^{-t}(W_\gamma^{cu}(z))| > 0$, which is a contradiction. Hence, the sequence $R_x^{-m_n}(z_n)$ converges to $y \in \Lambda$. Actually, $A^+ \subset \Lambda$. Notice that if there is some $y \in A^+ \cap W^s(p)$ for some $p \in \mathrm{Per}(X^t)$ we achieve a contradiction again. In fact, in such case, for k big enough, the set $U(y, \varepsilon_k)$ must be contained in some fundamental domain of $W^s(p)$. On the other hand, we can take ε_{n_0} small enough that $\mathcal{O}(p) \cap B_{\varepsilon_{n_0}}(J) = \emptyset$. Nevertheless, for n big, $y_n = R_x^{-m_n}(J(z_n)) \cap W^s(p)$ and then $y_n \to \mathcal{O}(p)$ as $n \to \infty$. However,

$$R_{x_{m_n}}^{m_n}(y_n) \in R_{x_{-m_n}}^{m_n}(R_x^{-m_n}(J(z_n))) = J(z_n) \subset B_{\varepsilon_n}(J) \subset B_{\varepsilon_{n_0}}(J)$$

which is a contradiction. Therefore, no point on A^+ belongs to the stable manifold of a periodic point in Λ.

Take some $y \in A^+$, $(z_n, n) \in I(y, \varepsilon_k)$ and $m_n > 0$, such that for any $w \in J(z_n) \cap \tilde{\Lambda}$ we have

$$W_\gamma^{cs}(R_x^{-m_n}(w)) \cap J \subset J^+.$$

Consider a map $\pi \colon A^+ \longrightarrow A^+$ defined as follows: For each $z \in A^+$, let $w = W_\gamma^{cs}(z) \cap J(z_n)$. Then define $\pi(z) := W_\gamma^{cs}(R_x^{-m_n})(w) \cap J$. By construction, $\mathrm{dist}(R_x^t(z), R_x^t(w)) \to 0$ and $\mathrm{dist}(R_x^{-t}(z_n), R_x^{-t}(w)) \to 0$ as $t \to \infty$. Then, $w \in \tilde{\Lambda}$ and hence this map is well defined. Moreover, it is continous and monotone. This implies that there exists $w_0 \in A^+ \subset J$ such that $\pi(w_0) = w_0$, that is, $R_x^{-m_n}(w_0)$ belongs to $W_\gamma^{cs}(w_0)$. As in the proof of Lemma 6.4 this situation implies that the α-limit of w_0 is a periodic orbit as well for the α limit of y. Hence x must be periodic. This contradiction implies there is $y^+ \in A^+$ such that $I(y^+, \varepsilon_k) = \emptyset$. Hence, if x is accumulated on both sides of $W_\gamma^{cs}(x)$ we are done. If x is accumulated only on one side, the arguments of the proof of Lemma 3.6.1 in [21] apply directly. □

We shall reffer to those points accumulated only on one side as boundary points of $\tilde{\Lambda}$. Notice that if $B_\varepsilon(J)$ is an adapted box, then any subbox is adapted too. As a consequence of the previous argument, for any non periodic point $x \in \Lambda$ small adapted boxes can be taken in such

a way $\partial^{cu}(B_\varepsilon(J))$ is contained in some central stable manifolds, unless the point x is a boundary point of $\tilde{\Lambda}$.

Consider an adapted box $B_\varepsilon(J)$ associated to some $x \in \Lambda$. If there is $z \in B_\varepsilon(J) \cap \Lambda$ such that $z_{-t} \in B_\varepsilon(J)$ $(t > 0)$ for the first time, we say (ψ_z, D_ψ) is a *return* of $B_\varepsilon(J)$ if D_ψ is the connected component which contain z of $D_z^{-t} \cap B_\varepsilon(J)$ and $\psi_z = R_z^{-t}|_{D_\psi}$. Notice the holonomy map is defined on some normal section that contains the box. Denote by $\mathcal{R} = \mathcal{R}(B_\varepsilon(J), \Lambda)$ the set of all returns associated to $B_\varepsilon(J)$. For $\psi \in \mathcal{R}$ we say $|\psi'| < \xi < 1$ if and only if for any $y \in J(z)$, $z \in D_\psi \cap \Lambda$ we have that $|DR_z^{-t}|_{\tilde{F}(y)}| < \xi$, where $\tilde{F}(y) = T_y W_\gamma^{cu}(z)$.

Let $\psi \in \mathcal{R}$. If $y \in D_\psi \cap \Lambda$ then $J(y) \subset D_\psi$; so, it is saturated by central unstable manifolds and hence, is a vertical strip; since any point $\tilde{y} \in J(y)$ shadows y until time $\tau_z^{-t}(\tilde{y})$. On the other hand, if $D_\psi \subset^s B_\varepsilon(J)$, then the image $B_\psi := \mathrm{img}(\psi)$ is an horizontal strip, *i.e.* intersects both components of $\partial^s B_\varepsilon(J)$, by continuity

We also say $\psi > \varphi$ for $\psi, \varphi \in \mathcal{R}$ if $\psi \circ \varphi^{-1}$ is obtained by flowing the orbits a negative amount of time. Remember we are considering first returns for the past.

Lemma 6.6. *For $\psi_1, \psi_2 \in \mathcal{R}$, if $\psi_1 \neq \psi_2$ then $D_{\psi_1} \cap D_{\psi_2}$ is not a domain of a return if the intersection does not contains $\partial^u(B_\varepsilon(J))$.*

Proof. Take $\varphi \in \mathcal{R}(B_\varepsilon(J), \Lambda)$ such that $D_\varphi = D_{\psi_1} \cap D_{\psi_2}$. Since $D_\varphi \cap \partial^u B_\varepsilon(J) = \emptyset$ then B_φ is an horizontal strip. Now $\psi_1(D_\varphi) \subset B_{\psi_1} \subset B_\varepsilon(J)$ since $D_\varphi \subset D_{\psi_1}$. Since φ is a return, there is $z \in D_\varphi$ such that $\varphi(z)$ is the first return of z to $B_\varepsilon(J)$. Hence $\psi_1 > \varphi$. Observe $\psi_1 \circ \varphi^{-1}(B_\varepsilon(J)) \subset B_\varepsilon(J)$, because it is an adapted box. This is a contradiction to the fact of Λ being transitive and not a single periodic orbit. □

Definition 6. An adapted box $B_\varepsilon(J)$ is well adapted if there is a subbox $B_{\varepsilon'}(J)$ and two disjoint vertical strips S_1, S_2 such that

$$B_\varepsilon(J) \setminus B_{\varepsilon'}(J) = S_1 \cup S_2$$

each one satisfies either: $S_i \cap \Lambda = \emptyset$ or S_i is a domain of some $\psi_i \in \mathcal{R}(B_\varepsilon(J), \Lambda)$, for which $\psi_i(S_i)$ is an horizontal strip.

Lemma 6.7. *Given any $x \in \Lambda \setminus Per(X^t)$, there exists well adapted boxes associated to x of arbitrarily small size.*

Proof. Let $B_\varepsilon(J)$ be a small adapted box associated to $x \in \Lambda \setminus \mathrm{Per}(X^t)$, for some $J \subset W_\gamma^{cu}(x)$. We shall prove there is a subbox $B_{\varepsilon'}(J)$ which is well adapted $(\varepsilon' \leqslant \varepsilon)$. It is important to remark that subboxes are not necessarily symetric with respect to J, that is, we shall only need

to find an interval $T \subset W^{cs}_{\varepsilon}(x)$, $x \in T$, such that $B_T(J)$ have the desidered properties. We shall divide the proof into two parts:

First we shall assume that for all $\varepsilon' \leqslant \varepsilon$:

$$B_{\varepsilon'}(J) \neq \{y \in J(z) | z \in \mathrm{cl}(B_{\varepsilon'}(J)) \cap \Lambda\}.$$

For instance, this happens on the right side of the box $B^+_{\varepsilon'}(J)$ and latter we shall see it implies the same happens on the left one. That is, for any $\varepsilon' \leqslant \varepsilon$:

$$B^+_{\varepsilon'}(J) \neq \{y \in J(z) | z \in \mathrm{cl}(B^+_{\varepsilon'}(J)) \cap \Lambda\}. \tag{9}$$

Then, there exists some $w \in W^{cs}_{\varepsilon}(x)$, on the right hand of x such that $w \notin J(z)$, for any $z \in \mathrm{cl}(B^+_{\varepsilon'}(J)) \cap \Lambda$. According to this order we have $x < w$ on $W^{cs}_{\varepsilon}(x)$. Define $w_z = J(z) \cap W^{cs}_{\varepsilon}(x)$ for any such z. If $w > w_z$ for all z then $S_2 = B^+_{\varepsilon'}(J) \setminus B_{(x,\tilde{w_z})}(J)$ is a vertical strip contained $B^+_{\varepsilon'}(J)$, where $\tilde{w_z} = \sup\{w_z;\, z\}$ and such that $S_2 \cap \Lambda = \emptyset$ as the definition requires. Otherwise, there is $z \in \mathrm{cl}(B^+_{\varepsilon'}(J)) \cap \Lambda$ such that $w_z > w$. Let $w_0 = \inf\{w_z | w_z > w\}$. Hence, reducing the right side of the box $B_{\varepsilon'}(J)$ to $B_{(x,w_0)}(J)$ we have that $S_2 = B^+_{\varepsilon'}(J) \setminus B_{(x,\tilde{w_z})}(J)$ and $S_2 \cap \Lambda = \emptyset$. Now we shall see that the left side of the box also verificates (9). In fact, assume by contradiction that for some $\varepsilon' > 0$

$$B^-_{\varepsilon'}(J) = \{y \in J(z) | z \in \mathrm{cl}(B^-_{\varepsilon'}(J)) \cap \Lambda\}. \tag{10}$$

We claim there is $\psi \in \mathcal{R}$ such that

$$D_\psi \cap B^-_{\varepsilon'}(J) \neq \emptyset \text{ and } D_\psi \cap (\partial^u B^-_{\varepsilon'}(J) \setminus J) = \emptyset.$$

If the claim holds, then B_ψ is an horizontal strip that crosses all $B_\varepsilon(J)$ and in particular $B_\psi \cap B^+_\varepsilon(J) \neq \emptyset$. This is a contradiction to (9). Therefore, (9) holds also for the left side of the box. Repeating the argument we have done for the right side on the left side, we obtain the desidered subbox of $B_\varepsilon(J)$ which is well adapted.

To prove the claim, first notice there are $\psi \in \mathcal{R}$ such that $D_\psi \cap B^-_{\varepsilon'}(J) \neq \emptyset$, since there are points of Λ on the left side of the box and Λ is transitive. If there is only one return, then $B^-_{\varepsilon'}(J) \subset D_\psi$ and also $B_\psi \subset (B^-_{\varepsilon'}(J) \setminus J)$, since $X^t(J) \cap J = \emptyset$, $\forall t \neq 0$ and also B_ψ satisfies (10). This is a contradiction to the transitivity of the set Λ. Hence, there are at least two returns on the left side of the box.

Suppose there are two different $\psi_1, \psi_2 \in \mathcal{R}$ such that

$$(\partial^u(B^-_{\varepsilon'}(J))) \setminus J) \subset D_{\psi_1} \cap D_{\psi_1},$$

$\psi_1 < \psi_2$ and there is no other return preceding ψ_2 except ψ_1. Notice $B_{\psi_1} \not\subseteq D_{\psi_1}$. Moreover, $B_{\psi_1} \subset (B^-_{\varepsilon'}(J) \setminus J)$, since $B^+_\varepsilon(J) \cap \Lambda$ have holes as we assumed in (9) and $X^t(J) \cap J = \emptyset$ for any $t \neq 0$. Also $D_{\psi_1} \subset B^-_{\varepsilon'}(J)$ since, in the other case $B^-_{\varepsilon'}(J) \subset D_{\psi_1}$ contradicting

again the transitivity of Λ. Notice $D_{\psi_1} \subset D_{\psi_2}$ since $\psi_1 < \psi_2$ and they are different returns. Therefore there exisits $z \in B^-_{\varepsilon'}(J) \setminus D_{\psi_1}$ that returns into $B^-_{\varepsilon'}(J)$ before ψ_2 by a continous extension of ψ_1. However, $z \in D_{\psi_2} \setminus D_{\psi_1}$; this contradiction proves our claim.

Now we assume that

$$B_\varepsilon(J) = \{y \in J(z) | z \in \mathrm{cl}(B_\varepsilon(J)) \cap \Lambda\}.$$

Take ψ_+ and $\psi_- \in \mathcal{R}$ such that $D_+ \subset B^+_\varepsilon(J)$, $D_- \subset B^-_\varepsilon(J)$ and that B_{ψ_+}, B_{ψ_-} are horizontal strips. It is not difficult to see that we can find two periodic intervals $J(w_+) \subset D_+$ and $J(w_-) \subset D_-$, that is

$$\psi_\pm(J(w_\pm)) \subset J(w_\pm).$$

Cutting the box along both intervals we obtain the desidered well adapted box, proving the Lemma. □

6.2. **Proof of Main Lemma.** Let $\Lambda_0 \subset M$ be a non trivial compact invariant set for X^t provided with a dominated splitting which is not an irrational torus. Assume that any compact invariant subset of Λ_0 is hyperbolic. We shall prove Λ_0 is hyperbolic. In order to do that, we need to prove both: $|L^{-t}|_{F(z_0)}| \to 0$ and $|L^t|_{E(z_0)}| \to 0$ as $t \to \infty$ for any $z_0 \in \Lambda_0$. We shall prove only the former limit. The latter is analogous.

First assume Λ_0 is not a minimal set. Recall a compact invariant set is minimal if and only if any orbit is dense. Now we are ready to apply the following lemmas. For their proof, see Lemma 3.7.4 and Lemma 3.7.2 in [21].

Lemma 6.8. *Let $B_\varepsilon(J)$ be a well adapted box such that $\#R(B_\varepsilon(J), \Lambda)$ is not bounded. Then, there exists a return $\psi_0 \in R(B_\varepsilon(J), \Lambda)$ such that the adapted box $B_{\psi_0} = img(\psi_0)$ satisfies that for every $\psi \in R(B_{\psi_0}, \Lambda)$ we have $|\psi'| < \frac{1}{2}$.*

Lemma 6.9. *Let $B_\varepsilon(J)$ be an adapted box such that for any $\psi \in \mathcal{R}(B_\varepsilon(J), \Lambda)$ we have $|\psi'| \leqslant \xi < 1$. Then, for every $y \in B_\varepsilon(J) \cap \Lambda_0$ we have*

$$\int_0^\infty |R^{-t}(J(y))| dt < \infty$$

and therefore $|D_y R^{-t}|_{F(y)}| \to 0$ as $t \to \infty$

Notice that $|D_y R^{-t}|_{F(y)}| \to 0$ as $t \to \infty$ implies $|L^{-t}|_{F(y)}| \to 0$ as $t \to \infty$.

By hypothesis there is $x \in \Lambda_0$ such that $x \notin \omega(x)$. Lemma 6.7 implies the existence of a small well adapted box $B_\varepsilon(J)$ associated to x such that $B_\varepsilon(J) \cap \{X^{-t}(x) | t > 0\} = \emptyset$. Notice $\#R(B_\varepsilon(J), \Lambda_0) = \infty$,

since Λ_0 is transitive. Therefore combining Lemma 6.8 and Lemma 6.9 we conclude that there exists a subbox B_0 ($= \mathrm{Img}(\psi_0)$ of the Lemma 6.8) where, for all $y \in B_0 \cap \Lambda_0$ we have $|D_y R^{-t}|_{F(y)}| \to 0$ as $t \to \infty$, and hence $|L^{-t}|_{F(y)}| \to 0$ as $t \to \infty$, as we have remarked above.

Now take $z_0 \in \Lambda_0$, if $\alpha(z_0) \neq \Lambda_0$, by hypothesis is a hyperbolic set. Therefore $|L^{-t}|_{F(z_0)}| \to 0$ as $t \to \infty$. On the other hand, if $\alpha(z_0) = \Lambda_0$, then there is $m > 0$ such that $z_{-m} \in B_0$. Then $|L^{-t}|_{F(z_{-m})}| \to 0$ as $t \to \infty$, and this is what we wanted to prove.

For the case Λ_0 is minimal, we have to use a diferent argument. Again, we state first two lemmas which correspond to Lemma 3.7.5 and Lemma 3.7.6 in [21].

Lemma 6.10. *There is $x_0 \in \Lambda_0$ and an adapted box of arbitrarily small size associated to x_0 such that $B_\varepsilon(J^+) \cap \Lambda_0 = \emptyset$ or $B_\varepsilon(J^-) \cap \Lambda_0 = \emptyset$.*

Lemma 6.11. *Let $B_\varepsilon(J)$ be an adapted box such that $B_\varepsilon(J^+) \cap \Lambda_0 = \emptyset$. Then there exists $K > 0$ such that $\forall y \in B_\varepsilon(J) \cap \Lambda_0$*

$$\int_0^\infty |R^{-t}(J^+(y))| dt < K$$

and moreover, there is $J_1(y)$, $J^+(y) \subset J_1(y) \subset J(y)$ where both components of $J_1(y) - J^+(y)$ have length bounded away from zero indepentently of y and some $\tilde{K} > 0$ where

$$\int_0^\infty |R^{-t}(J_1(y))| dt < \tilde{K}$$

In spite of these, notice that $B = \bigcup_{y \in B_\varepsilon(J) \cap \Lambda_0} J_1(y)$ is a two-dimensional open set contained in some transversal section to the flow where for any $y \in B \cap \Lambda_0$ we have:

$$\int_0^\infty |R^{-t}(J_1(y))| dt < \tilde{K}$$

and hence $|L^{-t}|_{F(y)}| \to 0$ as $t \to \infty$.

Let $z_0 \in \Lambda_0$. There is some $m > 0$ such that $z_{-m} \in B$. Then $|L^{-t}|_{F(z_{-m})}| \to 0$ as $t \to \infty$. This finishes the proof of the Main Lemma and concludes this work.

Instituto de Matemática Pura e Aplicada (IMPA), Brazil.
E-mail address: aubin@impa.br

References

[1] Bonatti, C., Viana, M. *SRB measures for partially hyperbolic dynamical systems whose central direction is mostly contracting*, Israel J. Math. 115 (2000), 157-193

[2] Doering, C.I. *Persistently transitive vector fields on three manifolds*, Dynam. Syst. And Biff. Theory, Pitman Res # 160 pp 59-89.
[3] Guckenheimer, J., Williams, R.F. *Structural stability of Lorenz attractors.*, Inst. Hautes Études Sci. Publ. Math. No. 50 (1979), 59–72.
[4] Hayashi, S. *Connecting invariant manifolds and the solution of the C^1 stability and Ω-stability conjectures for flows.*, Ann. of Math. (2) 145 (1997), no. 1, 81-137.
[5] Herman, M-R. *Sur la conjugaison différentiable des difféomorphismes du cercle à des rotations.* Inst. Hautes Études Sci. Publ. Math. No. 49 (1979), 5–233.
[6] Hirsch, M. W., Pugh, C. C., Shub, M. *Invariant manifolds.* Lecture Notes in Mathematics, Vol. 583. Springer-Verlag, Berlin-New York, 1977. 149 pp.
[7] Labarca, R., Pacífico M.J. *Stability of singularity horseshoes.* Topology 25 (1986), no. 3, 337–352
[8] Mañé, R. *Ergodic Theory and Differential Dynamics* Springer Verlag, New York, 1987.
[9] Morales, C.A., Pacífico, M. J.; Pujals, E. *On C^1 robust singular transitive sets for three-dimensional flows.* C. R. Acad. Sci. Paris Sér. I Math. 326 (1998), no. 1, 81–86.
[10] Newhouse, S. *Diffeomorphisms with infinitely many sinks.* Topology 13 (1974), 9–18.
[11] Newhouse, S. *Hyperbolic Limit Sets*, Trans. A.M.S. 167 (1972), 125 - 150.
[12] Newhouse, S. *Lectures on dynamical systems. Dynamical systems* (C.I.M.E. Summer School, Bressanone, 1978), Progr. Math., 8, Birkhuser, Boston, Mass., 1980, pp. 1–114.
[13] Oseledets, V.I. *A multiplicative ergodic theorem: Lyapunov characteristic numbers for dynamical systems.* Trans. Moscow. Math. Soc. 19:197-231, 1968.
[14] Palis, J. *A Global View of Dynamics and a Conjecture on the denseness of finitude of attractors* , Asterisque 261 (2000) 339-351.
[15] Palis, J. *On Morse-Smale Dynamical Systems*, Topology 8 (1968) 385-404.
[16] Palis, J., Smale, S. *Structural stability theorems*, Proc. A.M.S. Symp. Pure Math. 14 (1970), 223-232.
[17] Palis, J., Takens, F. *Hyperbolicity and sensitive chaotic dynamics of homoclinic bifurcations*, Cambridge Univ. Press. Cambridge, 1993.
[18] Pliss, V.A. *On a Conjecture of Smale*, Diff. Uravneija, 8, (1972), 268-282.
[19] Pugh, C. *The closing lemma*, Amer. J. Math., 89 (1967) 956-1009.
[20] Pugh, C. *An improved closing lemma and a general density theorem*, Amer. J. Math., 89 (1967) 1010-1021.
[21] Pujals, E., Sambarino, M. *Homoclinic tangencies and hyperbolicity for surface diffeomorphisms.*, Ann. of Math. (2) 151 (2000), no. 3, 961–1023.
[22] Pujals, E., Sambarino, M. *On the dynamics of Dominated Splitting*, to appear.
[23] Schwartz, A. J. *A generalization of a Poincaré-Bendixon Theorem to closed two dimensional Manifolds* Amer. J. Math. 85 (1963), 453-458, Errata, *ibid* 85 (1963), 753.
[24] Smale, S. *Differentiable dynamical systems.* Bull. A.M.S, 73, (1967), 747-817.
[25] Tahzibi, A. *Stably ergodic systems which are not partially hyperbolic*, to appear.