On the topological invariance of the algebraic multiplicity of a holomorphic vector field.

Rudy Rosas Bazán

September 2, 2005

Resumo

Nesta tese estudamos a equivalência topológica entre germes de campos de vetores holomorfos com uma singularidade isolada na origem de $\mathbb{C}^n$ $(n \geq 2)$. Assim, se a equivalência topológica é de classe C^1, provamos que a multiplicidade algébrica do campo (na origem) é invariante. Alem disso, se considerarmos somente campos de vetores em $\mathbb{C}^2$, é suficiente supor que a equivalência topológica é diferenciável na origem de $\mathbb{C}^2$.

Palavras chaves: vector field, holomorphic, multiplicity.

Preface

Given a curve $f : (\mathbb{C}^2, 0) \to (\mathbb{C}, 0)$, singular at $0 \in \mathbb{C}^2$, we define its *algebraic multiplicity* as the degree of the first nonzero jet of f, that is, $\nu(f) = \nu$ where

$$f = f_\nu + f_{\nu+1} + \cdots$$

is the Taylor development of f and $f_\nu \neq 0$. A well known result by Burau [5] and Zariski [6] states that ν is a *topological invariant*, that is, given $\widetilde{f} : (\mathbb{C}^2, 0) \to (\mathbb{C}, 2)$ and a homeomorphism $\mathrm{h} : U \to \widetilde{U}$ between neighborhoods of $0 \in \mathbb{C}^2$ such that $\mathrm{h}(f^{-1}(0) \cap U) = \widetilde{f}^{-1}(0) \cap V$ then $\nu(f) = \nu(\widetilde{f})$. Consider now a holomorphic vector field Z in $\mathbb{C}^2$ with a singularity at $0 \in \mathbb{C}^2$. If

$$Z = Z_\nu + Z_{\nu+1} + \cdots$$

we define $\nu = \nu(Z)$ as the *algebraic multiplicity* of Z. A natural question, posed by J.F.Mattei is: is $\nu(Z)$ a topological invariant of $\mathcal{F}_Z$?. In [2], the authors give a positive answer if $\mathcal{F}_Z$ is a *generalized curve*, that is, if the desingularization of Z does not contain complex saddle-nodes. In this work, we impose conditions on the topological equivalence $\mathrm{h} : U \to \widetilde{U}$. Thus, in Part II we prove the following:

Theorem. *Let* $\mathrm{h} : U \to \widetilde{U}$ *be a topological equivalence between* $\mathcal{F}_Z$ *and* $\mathcal{F}_{\widetilde{Z}}$ *and assume that* h *preserves the orientation of* $\mathbb{C}^2$*. Suppose that* h *is differentiable at* $0 \in \mathbb{C}^2$ *and such that* $d\,\mathrm{h}(0) : \mathbb{R}^4 \to \mathbb{R}^4$ *is a real isomorphism. Then the algebraic multiplicities of* Z *and* $\widetilde{Z}$ *are the same.*

In Part I, we consider holomorphic vector fields Z and $\widetilde{Z}$ in $\mathbb{C}^n \, (n \geq 2)$ with isolated singularity at $0 \in \mathbb{C}^n$ and prove the following:

Theorem. *Let* U *and* $\widetilde{U}$ *be neighborhoods of* $0 \in \mathbb{C}^n$ *and let* $\mathrm{h} : U \to \widetilde{U}$ *be a* C^1 *equivalence between* $\mathcal{F}_Z$ *and* $\mathcal{F}_{\widetilde{Z}}$*, that is, a* C^1 *diffeomorphism taking leaves of* $\mathcal{F}_Z$ *to leaves of* $\mathcal{F}_{\widetilde{Z}}$*. Then the algebraic multiplicities of* Z *and* $\widetilde{Z}$ *are equal.*

The two parts of this thesis are independent and can be read in any order.

Contents

Part I

The C^1 case

1 Introduction

Let $\mathcal{F}$ be a holomorphic foliation by curves of a neighborhood U of $0 \in \mathbb{C}^n$ with a unique singularity at $0 \in \mathbb{C}^n\,(n \geq 2)$. We assume that $\mathcal{F}$ is generated by the holomorphic vector field

$$V = \sum_{i=1}^{n} a_i \frac{\partial}{\partial z_i},\ a_i \in \mathcal{O}_U,\ g.c.d.(a_1, a_2, \ldots, a_n) = 1.$$

The algebraic multiplicity of $\mathcal{F}$ (at $0 \in \mathbb{C}^n$) is the minimum vanishing order at $0 \in \mathbb{C}^n$ of the functions a_i. Let $\widetilde{\mathcal{F}}$ be another holomorphic foliation by curves of a neighborhood $\widetilde{U}$ of $0 \in \mathbb{C}^n$ and let $\mathrm{h} : U \to \widetilde{U}$ be a topological equivalence between $\mathcal{F}$ and $\widetilde{\mathcal{F}}$, that is, a homeomorphism taking leaves of $\mathcal{F}$ to leaves of $\widetilde{\mathcal{F}}$. A natural question, posed by J.F.Mattei is: are the algebraic multiplicities of $\mathcal{F}$ and $\widetilde{\mathcal{F}}$ the same?. In [2], the authors give a positive answer if $n = 2$ and $\mathcal{F}$ is a *generalized curve*, that is, if its desingularization does not contain complex saddle-nodes. In this work we give a sufficient condition on the topological equivalence $\mathrm{h} : U \to \widetilde{U}$ for the algebraic multiplicity to be invariant. Let $\pi : \widehat{\mathbb{C}^n} \to \mathbb{C}^n$ be the quadratic blow up with center at $0 \in \mathbb{C}^n$. Clearly the map $h := \pi^{-1}\,\mathrm{h}\,\pi$ is a homeomorphism between $\pi^{-1}(U\backslash\{0\})$ and $\pi^{-1}(\widetilde{U}\backslash\{0\})$. Then we prove the following:

Theorem 1.1. *Suppose that h extends to the divisor $\pi^{-1}(0)$ as a homeomorphism between $\pi^{-1}(U)$ and $\pi^{-1}(\widetilde{U})$. Then the algebraic multiplicities of $\mathcal{F}$ and $\widetilde{\mathcal{F}}$ are the same.*

If h is a C^1 diffeomorphism, we prove that h extends to the divisor. Thus, we obtain that the algebraic multiplicity is invariant by C^1 equivalences:

Theorem 1.2. *Let $\mathcal{F}$ and $\widetilde{\mathcal{F}}$ be two foliations by curves of neighborhoods U and $\widetilde{U}$ of $0 \in \mathbb{C}^n$, $n \geq 2$. Let $\mathrm{h} : U \to \widetilde{U}$ be a C^1 equivalence between $\mathcal{F}$ and $\widetilde{\mathcal{F}}$, that is, a C^1 diffeomorphism taking leaves of $\mathcal{F}$ to leaves of $\widetilde{\mathcal{F}}$. Then the algebraic multiplicities of $\mathcal{F}$ and $\widetilde{\mathcal{F}}$ are equal.*

It is known that there exists a unique way of extending the pull back foliation $\pi^*(\mathcal{F}|_{U\backslash\{0\}})$ to a singular analytic foliation $\mathcal{F}_0$ on $\pi^{-1}(U)$ with singular set of codimension ≥ 2. We say that $\mathcal{F}_0$ is the strict transform of $\mathcal{F}$ by π. Let $\widetilde{\mathcal{F}}_0$ be the strict transform of $\widetilde{\mathcal{F}}$ by π. In order to prove Theorem 1.1 we show that the algebraic multiplicity of $\mathcal{F}$ depends on the Chern class of the tangent bundle of $\mathcal{F}_0$. To relate the Chern classes of the tangent bundles of $\mathcal{F}_0$ and $\widetilde{\mathcal{F}}_0$ we use the following theorem (see [1]).

Theorem 1.3. *Let $\mathcal{F}$ and $\widetilde{\mathcal{F}}$ be foliations by curves on the complex manifolds M and $\widetilde{M}$ respectively. Let $c(T\mathcal{F})$ denote the Chern class of the tangent bundle $T\mathcal{F}$ of $\mathcal{F}$. Let $h : M \to \widetilde{M}$ be a topological equivalence between $\mathcal{F}$ and $\widetilde{\mathcal{F}}$ and consider the map $h^* : H^2(M, \mathbb{Z}) \to H^2(\widetilde{M}, \mathbb{Z})$ induced in the cohomology. Then $h^*(c(T\mathcal{F})) = c(T\widetilde{\mathcal{F}})$.*

Clearly the homeomorphism $h : \pi^{-1}(U\backslash\{0\}) \to \pi^{-1}(\widetilde{U}\backslash\{0\})$ is a topological equivalence between $\mathcal{F}_0|_{\pi^{-1}(U\backslash\{0\})}$ and $\widetilde{\mathcal{F}}_0|_{\pi^{-1}(\widetilde{U}\backslash\{0\})}$. To be able to apply Theorem 1.3 we show that h extends as a topological equivalence between $\mathcal{F}_0$ and $\widetilde{\mathcal{F}}_0$. This is the non trivial part of the proof. Thus, we prove the following.

Theorem 1.4. *Let V and $\widetilde{V}$ be complex manifolds, let $Y \subset V$ and $\widetilde{Y} \subset \widetilde{V}$ be analytic subvarieties of codimension ≥ 1 and, let $\mathcal{F}$ and $\widetilde{\mathcal{F}}$ be holomorphic foliations by curves on V and $\widetilde{V}$ respectively. Suppose there is a homeomorphism h between V and $\widetilde{V}$ with $h(Y) = \widetilde{Y}$ and such that $h|_{V\backslash Y}$ is a topological equivalence between $\mathcal{F}|_{V\backslash Y}$ and $\widetilde{\mathcal{F}}|_{\widetilde{V}\backslash\widetilde{Y}}$. Then h is a topological equivalence between $\mathcal{F}$ and $\widetilde{\mathcal{F}}$.*

This part is organized as follows. In section 2, we state and prove a lemma used in section 3 to prove Theorem 1.4. In section 4 we prove Theorem 1.1. Finally, section 5 discusses the case C^1.

2 A fundamental lemma

Let $\mathbb{D} = \{z \in \mathbb{C} : ||z|| < 1\}$ and $\mathbb{B} = \{z \in \mathbb{C}^{n-1} : ||z|| < 1\}$ where $n \geq 2$. Let M be a complex manifold of complex dimension n and let D be a subset of M homeomorphic to a disc. We say that D is a *singular disc* if for all $x \in D$ there exist a neighborhood $\mathcal{D}$ of x in D, and an injective holomorphic function $f : \mathbb{D} \to M$ such that $f(\mathbb{D}) = \mathcal{D}$. If $f'(0) = 0$ we say that x is a *singularity* of D, otherwise x is a *regular point* of D (this does not depend on f). The set S of singularities of D is discrete and closed in D and we have that $D\backslash S$ is a complex submanifold of M. Thus, if x is a regular point of D, there is a neighborhood U of x in M and holomorphic coordinates (w, z), $w \in \mathbb{B}$, $z \in \mathbb{D}$ on U such that $D \cap U$ is represented by $(w = 0)$. If D does not have singularities we say that it is a *regular disc*. In this case, by uniformization, there is a holomorphic map $f : E \to M$, where $E = \mathbb{D}$ or $\mathbb{C}$, such that f is a biholomorphism between E and D.
Example. Let $\mathcal{F}$ be a holomorphic foliation by curves on the complex manifold M and let $D \subset M$ be homeomorphic to a disc. If D is contained in a leaf of $\mathcal{F}$ then it is a regular disc.

Lemma 2.1. *Let $F : \mathbb{D}\times[0, 1] \to \mathbb{C}^n$ be a continuous map such that for all $t \in [0, 1]$, the map $F(*, t) : \mathbb{D} \to \mathbb{C}^n$ is a homeomorphism onto its image. Thus, we have a continuous family of discs $D_t := F(\mathbb{D} \times \{t\})$. Suppose D_t is a regular disc for each $t > 0$. Then D_0 is a singular disc.*

Remark. Actually, we may only assume that $\mathcal{D}_t$ is a singular disc for all $t > 0$.
We now state the lemmas used in the proof of Lemma 2.1.

Lemma 2.2. *Let U be a simply connected domain in the complex plane such that $\partial\,\mathrm{U}$ is a Jordan curve. Then any uniformization $f : \mathbb{D} \to \mathrm{U}$ extends as a homeomorphism between $\overline{\mathbb{D}}$ and $\overline{\mathrm{U}}$.*

Lemma 2.3. *For each $k \in \mathbb{N}$, let $\phi_k : \mathbb{S}^1 \to \mathbb{S}^1$ be a homeomorphism. Then $\{\phi_k\}$ has a subsequence which converges almost everywhere with respect to the Lebesgue measure in the circle.*

Lemma 2.4. *Let $\phi : \mathbb{S}^1 \to \mathbb{C}$ be a bounded measurable function. Suppose that $\int_{\mathbb{S}^1} z^n\phi(z)dz = 0$ for all $n \in \mathbb{Z}$. Then, with respect to the Lebesgue measure in the circle, ϕ vanishes almost everywhere.*

Lemma 2.5. *Let $\mathcal{D} \subset \mathbb{C}^n$ be a bounded set homeomorphic to a disc. Let $p \in \mathcal{D}$ and suppose that $\mathcal{D}\backslash\{p\}$ has a complex structure such that the inclusion $\mathcal{D}\backslash\{p\} \to \mathbb{C}^n$ is holomorphic. Then there is a holomorphic injective map $g : \mathbb{D} \to \mathbb{C}^n$ with $g(\mathbb{D}) = \mathcal{D}$, $g(0) = p$.*

Proof of Lemma 2.1. Let $p = F(x_0, 0)$ be any point in D_0. Let $\mathrm{U} \subset \mathbb{D}$ be a disc centered at x_0 and such that $\overline{\mathrm{U}} \subset \mathbb{D}$. Let $t_k > 0$ be such that $t_k \to 0$ as $k \to \infty$ an define $\mathcal{D}_k = F(\mathrm{U} \times \{t_k\})$. Clearly $\overline{\mathcal{D}}_k \subset \overline{D}_{t_k}$. By uniformization, D_{t_k} is equivalent to a subset of $\mathbb{C}$ and, since $\mathcal{D}_k$ is a proper subset of D_{t_k}, we have (again by uniformization) that $\mathcal{D}_k$ is holomorphically equivalent to the unitary disc $\mathbb{D}$. Then there is a holomorphic map $f_k : \mathbb{D} \to \mathbb{C}^n$ which is a biholomorphism between $\mathbb{D}$ and $\mathcal{D}_k$. If we think that D_{t_k} is a subset of $\mathbb{C}$, by applying Lemma 2.2, we conclude that f_k extends as a homeomorphism $f_k : \overline{\mathbb{D}} \to \overline{\mathcal{D}}_k$. We may assume that $f_k(0) = F(x_0, t_k)$ for all k; otherwise we compose f_k with a suitable Moebius transformation. Observe that $f_k(\mathbb{D})$ is contained in the compact set $F(\overline{\mathrm{U}} \times [0,1])$, hence $\{f_k\}$ is uniformly bounded and, by Montel's theorem, we can assume that f_k converges uniformly on compact sets to a holomorphic function $f : \mathbb{D} \to \mathbb{C}^n$. Note that $f(0) = p$, since

$$f(0) = \lim_{k\to\infty} f_k(0) = \lim_{k\to\infty} F(x_0, t_k) = F(x_0, 0) = p.$$

Let $\mathbb{S}^1 = \partial\mathbb{D}$ and consider for each k the homeomorphism

$$\varphi_k := f_k|_{\mathbb{S}^1} : \mathbb{S}^1 \to \partial\mathcal{D}_k.$$

Define

$$\pi : F(\partial\,\mathrm{U} \times [0,1]) \to \mathbb{S}^1$$
$$\pi(F(\zeta, t)) = \zeta.$$

Clearly, π maps $\partial\mathcal{D}_k = F(\partial\,\mathrm{U} \times \{t_k\})$ homeomorphically onto $\mathbb{S}^1$. Therefore for each k, the map

$$\phi_k := \pi \circ \varphi_k : \mathbb{S}^1 \to \mathbb{S}^1$$

is a homeomorphism. By taking a subsequence, we may assume that ϕ_k converges a.e. to a function $\phi : \mathbb{S}^1 \to \mathbb{S}^1$ (Lemma 2.3). Therefore φ_k converges a.e. to

$$\varphi := \pi^{-1} \circ \phi : \mathbb{S}^1 \to \partial\mathcal{D}_0.$$

Fix $x \in \mathbb{D}$. Since $\{\varphi_k\}$ is uniformly bounded, by the dominated convergence theorem we have that

$$\frac{1}{2\pi i}\int_{\mathbb{S}^1} \frac{\varphi_k(w)}{w - x}dw \to \frac{1}{2\pi i}\int_{\mathbb{S}^1} \frac{\varphi(w)}{w - x}dw \tag{1}$$

as $k \to \infty$. By the Cauchy's integral formula, the left part of (1) is equal to $f_k(x)$ and, since $f_k(x) \to f(x)$, we conclude that

$$f(x) = \frac{1}{2\pi i}\int_{\mathbb{S}^1} \frac{\varphi(w)}{w - x}dw. \tag{2}$$

Assertion 1. *$f : \mathbb{D} \to \mathbb{C}^n$ is not constant.*

Proof. Assume by contradiction that f is a constant function. Then $f^{(n)}(0) = 0$ for all $n \in \mathbb{N}$, where $f^{(n)}$ is the nth derivative of f. From (2), by induction on n, it is not difficult to prove that

$$f^{(n)}(0) = \frac{1}{2\pi i}\int_{\mathbb{S}^1} \frac{\varphi(w)}{w^{n+1}}dw = 0$$

for all $n \geq 1$. Hence

$$\int_{\mathbb{S}^1} w^n \varphi(w)dw = 0 \quad \text{for all} \quad n \leq -2. \tag{3}$$

On the other hand, for each k and any $n \geq 0$ we have that $\int_{\mathbb{S}^1} w^n \varphi_k(w)dw = 0$ because $w^n\varphi_k(w)$ extends holomorphically to $\mathbb{D}$ as $w^n f_k(w)$. Then, by the dominated convergence theorem we have that

$$\int_{\mathbb{S}^1} w^n \varphi(w)dw = \lim_{k\to\infty}\int_{\mathbb{S}^1} w^n \varphi_k(w)dw = 0 \quad \text{for all} \quad n \geq 0. \tag{4}$$

Thus, from (3) and (4):

$$\int_{\mathbb{S}^1} w^n \varphi(w)dw = 0 \quad \text{for all} \quad n \in \mathbb{Z}\backslash\{-1\}.$$

And easy computation shows that each coordinate of the function $\phi := \varphi - p$ satisfies the hypothesis of Lemma 2.4. Then $\phi = 0$ a.e. and therefore $\varphi = p$ a.e., which is a contradiction because $\varphi(\mathbb{S}^1) \subset \partial\mathcal{D}_0$ and $p \notin \partial\mathcal{D}_0$.

Assertion 2. $f(\mathbb{D}) \subset D_0$.

Proof. Let $z \in \mathbb{C}^n$ be such that $z = f(x)$. Then $z = \lim f_k(x)$. Since $f_k(x)$ is contained in $\mathcal{D}_k = F(\mathrm{U}\times\{t_k\})$, we have that $f_k(x) = F(x_k, t_k)$ with $x_k \in \mathrm{U}$. By taking a subsequence, we may assume that $x_k \to \bar{x} \in \overline{\mathrm{U}}$. Then

$$z = \lim_{k\to\infty} f_k(x) = \lim_{k\to\infty} F(x_k, t_k) = F(\lim_{k\to\infty} x_k, 0) = F(\bar{x}, 0) \in D_0.$$

Therefore $f(\mathbb{D}) \subset D_0$.

It follows from Assertion 1 that f' does not vanish. Then, we know that the zero set of f' is discrete and closed in $\mathbb{D}$. Hence, there exists a disc $\Omega \subset \mathbb{D}$ centered at 0 such that $f' \neq 0$ on $\Omega\backslash\{0\}$. Since f is not constant, 0 is an isolated point in $f^{-1}(p)$. Thus, we assume Ω to be small enough such that $\overline{\Omega} \cap f^{-1}(p) = \{0\}$. In particular, $f(\partial\Omega)$ does not pass through p.

Assertion 3. There is a disc $\mathcal{D} \subset D_0$ with $p \in \mathcal{D}$ and such that $\mathcal{D} \subset f(\Omega)$.

Proof. Let $x \in \Omega\backslash\{0\}$. Then $f'(x) \neq 0$ and there exists a disc $\Delta \subset \Omega$ with $x \in \Delta$ and such that $f|_{\overline{\Delta}} : \overline{\Delta} \to \mathbb{C}^n$ is injective, hence a homeomorphism onto its image, since $\overline{\Delta}$ is compact. Then $f(\Delta)$ is homeomorphic to a disc and, since $f(\Delta) \subset D_0$, we have that $f(\Delta)$ is open in D_0. Then $f(x)$ is an interior point of $f(\Omega)$ as a subset of D_0. It follows that every point $z \in f(\Omega\backslash\{0\})$ is an interior point of $f(\Omega) \subset D_0$. Thus if

z is a point in the boundary of $f(\Omega)$, since z is not an interior point, we have that $z \notin f(\partial\Omega \cup \{0\}) = f(\partial\Omega) \cup \{p\}$. Therefore:

$$\partial f(\Omega) \subset f(\partial\Omega) \cup \{p\}.$$

Since $f(\partial\Omega)$ does not pass through p, we may take a disc $\mathcal{D} \subset D_0$ containing p and such that $\mathcal{D}$ is disjoint of $f(\partial\Omega)$. Finally, we claim that $\mathcal{D} \subset f(\Omega)$. Let $z \in \mathcal{D}$ and suppose that $z \notin f(\Omega)$. Since $\mathcal{D}$ contains p, we may take $x \neq 0$, close enough to 0, such that $z' := f(x) \in \mathcal{D}$. We have $z' \neq p$ because $x \neq 0$, hence we may take a path γ in $\mathcal{D}\backslash\{p\}$ connecting z and z'. Since $z \notin f(\Omega)$ and $z' \in f(\Omega)$, there exists $z'' \in \gamma$ such that $z'' \in \partial f(\Omega)$. Then, since $\partial f(\Omega) \subset f(\partial\Omega) \cup \{p\}$, we have $z'' \in f(\partial\Omega) \cup \{p\}$. But this is a contradiction because $z'' \in \gamma$ is contained in $\mathcal{D}\backslash\{p\}$, which is disjoint of $f(\partial\Omega) \cup \{p\}$.

Let $z \in \mathcal{D}\backslash\{p\}$. By Assertion 3, $z = f(x)$ with $x \in \Omega$. Since $z \neq p$ we have $x \neq 0$, hence $f'(x) \neq 0$. Then, from the proof of Assertion 3, there exists a disc $\Delta \subset \Omega$, $x \in \Delta$, such that $f|_\Delta : \Delta \to \mathbb{C}^n$ is injective and $f(\Delta)$ is a neighborhood of z in $\mathcal{D}\backslash\{p\}$. Since f is holomorphic it follows that $\mathcal{D}\backslash\{p\}$ is a Riemann surface and the inclusion $\mathcal{D}\backslash\{p\} \to \mathbb{C}^n$ is a holomorphic map. Then, by Lemma 2.5, there is a holomorphic injective map $g : \mathbb{D} \to \mathbb{C}^n$ with $g(\mathbb{D}) = \mathcal{D} \subset D_0$. Since p was arbitrary, it follows that D_0 is a singular disc, which finishes the proof of Lemma 2.1. □

Proof of Lemma 2.2. See [7] p.310. □

Proof of Lemma 2.3. We give a sketch of the proof. By taking a subsequence we may assume that ϕ_k converges on a dense subset of $\mathbb{S}^1$. Let $\pi : \mathbb{R} \to \mathbb{S}^1$ be a covering. For each k, we may choose a lifting $f_k : \mathbb{R} \to \mathbb{R}$ of ϕ_k by π. Since ϕ_k is a homeomorphism, f_k is monotone and we may assume that f_k is increasing for all k. We may also assume that f_k converges on a dense subset R of $\mathbb{R}$. For all $y \in R$, we define $f(y) = \lim f_k(y)$. Observe that f is increasing, since so is f_k for all k. We extend f to $\mathbb{R}$ as

$$f(x) = \limsup_{y \in R, y < x} f(y).$$

It is not difficult to see that $f : \mathbb{R} \to \mathbb{R}$ is increasing. Then f is continuous on a set A of total measure. Now, it is not difficult to prove that for all $x \in A$, the sequence $f_k(x)$ converges to $f(x)$ and the lemma follows. □

Proof of Lemma 2.4. We claim that $\int_{\mathbb{S}^1} f(z)\phi(z)dz = 0$ for all continuous function $f : \mathbb{S}^1 \to \mathbb{C}$. By the Stone -Weierstrass approximation theorem (see [9]), f can be uniformly approximated by a sequence of functions $P_k = A_k + iB_k$, $k \in \mathbb{N}$, where $A_k, B_k : \mathbb{R}^2 \to \mathbb{R}$ are real polynomials. Let $z = x + iy$ and observe that $\bar{z} = 1/z$ if $z \in \mathbb{S}^1$. Then

$$\begin{aligned} P_k(z) &= A_k(x,y) + iB_k(x,y) = A_k(\frac{z+\bar{z}}{2}, \frac{z-\bar{z}}{2}) + iB_k(\frac{z+\bar{z}}{2}, \frac{z-\bar{z}}{2}) \\ &= A_k(\frac{z+1/z}{2}, \frac{z-1/z}{2}) + iB_k(\frac{z+1/z}{2}, \frac{z-1/z}{2}). \end{aligned}$$

Hence $P_k(z) = \sum a_j z^j$ where j runs on the integers, the sum is finite and the coefficients a_j depend only on the coefficients of A_k and B_k. Then

$$\int_{\mathbb{S}^1} P_k(z)\phi(z)dz = \int_{\mathbb{S}^1} (\sum_j a_j z^j)\phi(z)dz = \sum_j a_j \int_{\mathbb{S}^1} z^j \phi(z)dz = 0,$$

by hypothesis. Since P_k converges uniformly to f and ϕ is bounded we have that

$$\int_{\mathbb{S}^1} P_k(z)\phi(z)dz \to \int_{\mathbb{S}^1} f(z)\phi(z)dz$$

as $k \to \infty$ and therefore $\int_{\mathbb{S}^1} f(z)\phi(z)dz = 0$.

Now, we take an uniformly bounded sequence of continuous functions $f_k : \mathbb{S}^1 \to \mathbb{C}$ which converges a.e. to $\bar{\phi}$. Then $\{f_k\phi\}$ is uniformly bounded and converges a.e. to $\bar{\phi}\phi = |\phi|^2$. Thus by the dominated convergence theorem we have that

$$\int_{\mathbb{S}^1} f_k(z)\phi(z)dz \to \int_{\mathbb{S}^1} |\phi(z)|^2 dz$$

as $k \to \infty$. Therefore $\int_{\mathbb{S}^1} |\phi(z)|^2 dz = 0$ and it follows that $\phi = 0$ almost everywhere. $\square$

Proof of Lemma 2.5. Let A_r denote the annulus $\{z \in \mathbb{C},\ r < |z| < 1\}$ where $r \geq 0$. Since $\mathcal{D} \subset \mathbb{C}^n$ is bounded and $\mathcal{D}\backslash\{p\}$ is homeomorphic to an annulus we have (see [10]) that there exist a biholomorphism

$$g : A_r \to \mathcal{D}\backslash\{p\},$$

such that $g(z) \to p$ as $|z| \to r$. Take R with $r < R < 1$ and let Γ_r and Γ_R be denote the circles $|z| = r$ and $|z| = R$ respectively. For $r < |z| < R$ we have the formula

$$g(z) = \frac{1}{2\pi i}\int_{\Gamma_R} \frac{g(w)}{w - z}dw - \frac{1}{2\pi i}\int_{\Gamma_r} \frac{p}{w - z}dw,$$

since g extends continuously to Γ_r as $g|_{\Gamma_r} = p$. But the second integral is equal to zero because $p/(w - z)$ is holomorphic on the disc $|w| < r$, then

$$g(z) = \frac{1}{2\pi i}\int_{\Gamma_R} \frac{f(w)}{w - z}dw.$$

Therefore g extends to the disc $|z| < R$. Since g is not constant and $g|_{\Gamma_r} = p$ we necessarily have $r = 0$ and the lemma follows. $\square$

3 An extension theorem

This section is devoted to prove Theorem 1.4. We show first that Theorem 1.4 is a consequence of the following theorem.

Theorem 3.1. *Let $\mathcal{F}$ be a foliation by curves on the complex manifold M. Let $X \subset M$ be an analytic subvariety of codimension ≥ 1. Suppose that:*

(i) $\mathcal{F}$ is generated by a holomorphic vector field.

(ii) There exists a homeomorphism $h : \Sigma \times D \to M$, where Σ is a ball in $\mathbb{C}^{n-1}$ and D is a disc in $\mathbb{C}$.

(iii) If $D_z := h(\{z\} \times D)$ then for all z: either D_z is contained in X, or $D_z \cap X$ is discrete and $D_z\backslash X$ is contained in a leaf of $\mathcal{F}$.

Then $\mathcal{F}$ is regular and the sets D_z are the leaves of $\mathcal{F}$.

Proof of Theorem 1.4. Let p be a point in Y which is regular for $\mathcal{F}$. Let Σ denote a ball in $\mathbb{C}^{n-1}$ and D a disc in $\mathbb{C}$. Consider a neighborhood W of p on which $\mathcal{F}$ is a product foliation, that is, $W \simeq \Sigma \times D$ and the sets $\{z\} \times D$ are the leaves of $\mathcal{F}|_W$. We take W small enough such that $\widetilde{\mathcal{F}}$ restricted to $M := h(W)$ is generated by a holomorphic vector field. Let X be the intersection between M and $\widetilde{Y}$. We will show that the hypothesis of Theorem 3.1 hold for $\widetilde{\mathcal{F}}$ restricted to M. Hypothesis (i) and (ii) of 3.1 evidently hold. Let $D_z = h(\{z\} \times D)$.

Assertion 1. For all $z \in \Sigma$, either $\{z\} \times D$ is contained in Y, or $S'_z := (\{z\} \times D) \cap Y$ is discrete and closed in $\{z\} \times D$.

Proof. Since Y is closed, we have that $A = \{z\} \times D \cap Y$ is closed in $\{z\} \times D$. Suppose that A is not discrete. Let $x \in A$ be an accumulation point of A and let f be a holomorphic function which defines Y on a neighborhood of x. Let $\mathcal{D}$ be a disc in $\{z\} \times D$, with $x \in \mathcal{D}$, and such that f is defined on $\mathcal{D}$. Since $A \subset Y$ we have that f vanishes on $A \cap \mathcal{D}$. Then f vanishes on $\mathcal{D}$ because $A \cap \mathcal{D}$ has $x \in \mathcal{D}$ as an acummulation point. Thus, $\mathcal{D} \subset Y$ and we have therefore $\mathcal{D} \subset A$. It follows that A is open in $\{z\} \times D$ and, by connectedness, $A = \{z\} \times D \cap Y$. Thus $\{z\} \times D$ is contained in Y.

Suppose that D_z is not contained in X. Let $S_z = h(S'_z)$, where S'_z is given by Assertion 1. Then S_z is discrete in D_z. Observe that $(\{z\} \times D)\backslash S'_z$ is contained in a leaf of $\mathcal{F}|_{M\backslash Y}$. Then, since $h|_{M\backslash Y}$ is a topological equivalence between $\mathcal{F}|_{V\backslash Y}$ and $\widetilde{\mathcal{F}}|_{\widetilde{V}\backslash\widetilde{Y}}$, it follows that

$$D_z\backslash S_z = h((\{z\} \times D)\backslash S'_z)$$

is contained in a leaf of $\widetilde{\mathcal{F}}$. Thus, hypothesis (iii) of 3.1 holds. Then $\widetilde{\mathcal{F}}$ is regular on $M = h(W)$ and every D_z is contained in a leaf of $\widetilde{\mathcal{F}}$. Therefore we conclude:

Assertion 2. If p is a point in Y which is regular for $\mathcal{F}$, then p is mapped by h to a regular point of $\widetilde{\mathcal{F}}$. Moreover, there exists a neighborhood Ω of p in its leaf which is mapped by h onto a neighborhood of $h(p)$ in its leaf.

Now, by using Assertion 2 for h and h^{-1}, we deduce that p is regular for $\mathcal{F}$ if and only if $h(p)$ is regular for $\widetilde{\mathcal{F}}$. Hence

$$h(\mathrm{Sing}(\mathcal{F})) = \mathrm{Sing}(\widetilde{\mathcal{F}}).$$

It remains to prove that h maps any leaf of $\mathcal{F}$ onto a leaf of $\widetilde{\mathcal{F}}$. Let p be a regular point of $\mathcal{F}$. Let L be the leaf of $\mathcal{F}$ passing through p and let $\widetilde{L}$ be the leaf of $\widetilde{\mathcal{F}}$ passing through $h(p)$. Let A be the set of points in L which are mapped by h into $\widetilde{L}$. By Assertion 2, if $x \in A$ there exists a neighborhood of x in L_p contained in A. Therefore A is open. Now, let $x \notin A$. Then $h(x) \notin \widetilde{L}$. Thus, if $L' \neq L$ is the leaf of $\widetilde{\mathcal{F}}$ passing through $h(x)$ it follows by Assertion 2 that there exists a neighborhood Ω of x in L which is mapped by h into $L' \neq \widetilde{L}$, hence Ω is contained in $L\backslash A$. Then A is also closed and it follows by conectedness that $A = L$, that is, $h(L) \subset \widetilde{L}$. Analogously, we prove that $h^{-1}(\widetilde{L}) \subset L$. Therefore $h(L) = \widetilde{L}$. $\square$

We proceed now to prove Theorem 3.1.

Proposition 3.2. *Let $\mathcal{F}$ be a foliation by curves on the complex manifold M. Let $X \subset M$ be an analytic subvariety of codimension ≥ 1. Suppose that:*

(i) There exists a homeomorphism $h : \Sigma \times D \to M$, where Σ is a ball in $\mathbb{C}^{n-1}$ and D is a disc in $\mathbb{C}$.

(ii) If $D_z := h(\{z\} \times D)$ then for all z: either D_z is contained in X, or D_z is contained in a leaf of $\mathcal{F}$.

Consider $z' \in \Sigma$ and suppose that $D_{z'}$ is a singular disc. Let $S_{z'}$ the set of singularities of $D_{z'}$. Then $D_{z'} \backslash S_{z'}$ is contained in a leaf of $\mathcal{F}$.

Proof. It is sufficient to prove the following.

Assertion. If $p \in D_{z'} \backslash S_{z'}$ then p has a neighborhood in $D_{z'} \backslash S_{z'}$ contained in a leaf of $\mathcal{F}$.

Suppose Assertion holds. let L be a leaf of $\mathcal{F}$ and let $x \in (D_{z'} \backslash S_{z'}) \cap L$. By Assertion, there is a neighborhood Δ of x in $D_{z'} \backslash S_{z'}$ such that $\Delta \subset L$. Then $\Delta \subset (D_{z'} \backslash S_{z'}) \cap L$ and it follows that the intersection of $D_{z'} \backslash S_{z'}$ with any leaf is open in $D_{z'} \backslash S_{z'}$. Then, since $D_{z'} \backslash S_{z'}$ is connected, we have that it is contained in a unique leaf.

Proof of Assertion. Let p in $D_{z'} \backslash S_{z'}$. Since p is a regular point of the singular disc $D_{z'}$, on a neighborhood $U \subset M$ of p we may consider coordinates (w, y), $w \in \mathbb{B}$, $y \in \mathbb{D}$ with $p = (0,0)$ and such that $D_{z'} \cap U$ is represented by $(w = 0)$. Suppose that $p = h(z', t')$. Let Σ' be a ball in Σ containing z' and let D' be a disc in D containing t'. Then $W = \Sigma' \times D'$ is a neighborhood of (z', t') and, by taking W small enough, we assume $h(\overline{W}) \subset U$. Let $D'_z = h(\{z\} \times D')$. Note that $D'_{z'} \subset D_{z'} \cap U$, hence $D'_{z'}$ is contained in $(w = 0)$. Let $g : U \to \mathbb{D}$ be the projection $g(w, y) = y$. Consider $z \in \Sigma'$ and suppose $D_z \backslash X \neq \emptyset$. By hypothesis (ii), D_z is contained in a leaf of $\mathcal{F}$. Therefore D'_z is contained in leaf of $\mathcal{F}$ and we have that $g|_{D'_z} : D'_z \to \mathbb{D}$ is a holomorphic map. Remember that $D'_{z'} \subset (w = 0)$. Then $g|_{D'_{z'}} : D'_{z'} \to \mathbb{D}$ is given by $(0, y) \to y$ and is therefore a one to one map. Then $g(D'_{z'})$ is a disc in $\mathbb{D}$ with $g(\partial D'_{z'})$ as boundary. Note that $p = (0,0) \in D'_{z'}$, hence 0 is contained in the disc $g(D'_{z'})$. Therefore the curve $g(\partial D'_{z'})$ winds once around 0. By the continuity of h we assume Σ' small enough such that $g(\partial D'_z)$ is homotopic to $g(\partial D'_{z'})$ in $\mathbb{D} \backslash \{0\}$ for all $z \in \Sigma'$. Then $g(\partial D'_z)$ winds once around 0 and $g|_{D'_z}$ has therefore a unique zero. In other words, the plaque D'_z intersects $Y = \mathbb{B} \times \{0\} \subset U$ at a unique point. Thus, we can define the map $f : h(W) \backslash X \to Y$ by $f(D'_z \backslash X) = D'_z \cap Y$ whenever $D'_z \backslash X \neq \emptyset$. We have that f is holomorphic because it is constant along the leaves and, restricted to any transversal, is a holonomy map. Since f is bounded and X has codimension ≥ 1, by the generalized Riemann's extension theorem, f extends to a holomorphic function on $h(W)$. Observe that f restricted to Y is the identity map, then f is a submersion in a neighborhood V of Y. Hence f defines a regular foliation $\mathcal{N}$ on V. It is easy to see that $\mathcal{N}$ coincides with $\mathcal{F}$ on $V \backslash X$, thus $\mathcal{N} = \mathcal{F}$. Therefore $p \in Y$ is a regular point of $\mathcal{F}$.

Now, by reducing the neighborhood $W = \Sigma' \times D'$ of (z', t'), we may assume that $h(W)$ is contained in a neighborhood of p where $\mathcal{F}$ is given by a submersion f. Obviously

$D'_{z'}$ is a neighborhood of p in D_z. We shall prove that $D'_{z'}$ is contained in a leaf of $\mathcal{F}$ (the leaf passing through p). If $D'_{z'}$ is not contained in X, so is $D_{z'}$ and, by hypothesis (ii), we have that $D'_{z'}$ is contained in a leaf of $\mathcal{F}$. On the other hand, suppose that $D'_{z'}$ is contained in X. Then there exists a sequence of points $z_k \to z'$ such that $h(\{z_k\} \times D)$ is not contained in X, otherwise $h(\Sigma'' \times D) \subset X$ for some neighborhood $\Sigma'' \subset \Sigma$ of z', which is a contradiction because X has codimension ≥ 1. Thus, by (ii), we have that D'_{z_k} is contained in a leaf of $\mathcal{F}$ for all k. Recall $D'_{z_k} \subset h(W)$ is contained in a domain where $\mathcal{F}$ is given by the submersion f. Then f is constant over $D'_{z_k} = h(\{z_k\} \times D')$ and in particular, for all $t \in D'$ we have $f(h(z_k,t)) = f(h(z_k,t'))$. Then:

$$\begin{aligned} f(h(z',t)) &= f(h(\lim_{k\to\infty} z_k,t)) = \lim_{k\to\infty} f(h(z_k,t)) \\ &= \lim_{k\to\infty} f(h(z_k,t')) = f(h(\lim_{k\to\infty} z_k,t')) \\ &= f(h(z',t')). \end{aligned}$$

Therefore, for all $t \in D'$ we have that $h(z',t)$ and $h(z',t')$ are contained in the same Leaf. It follows that $D'_{z'}$ is contained in the leaf passing trough $h(z',t')$. Thus, Assertion is proved. $\square$

Proposition 3.3. *Let $\mathcal{F}$ be a foliation by curves on the complex manifold M such that:*

(i) *$\mathcal{F}$ is generated by a holomorphic vector field.*

(ii) *There exists a homeomorphism $h : \Sigma \times D \to M$, where Σ is a ball in $\mathbb{C}^{n-1}$ and D is a disc in $\mathbb{C}$.*

(iii) *For all z, there is a discrete closed set $S_z \subset D_z := h(\{z\} \times D)$ such that $D_z \backslash S_z$ is contained in a leaf of $\mathcal{F}$.*

Then $\mathcal{F}$ is regular and the sets D_z are the leaves of $\mathcal{F}$.

We need the following Lemmas.

Lemma 3.4. *Let $f : \overline{\mathbb{D}} \to \mathbb{C}$ be smooth, and holomorphic on $\mathbb{D}$. Suppose that f is regular on $\mathbb{S}^1$. Then f is a regular map if and only if the curve $f|_{\mathbb{S}^1} : \mathbb{S}^1 \to \mathbb{C}$ has degree 1*[1]*.*

Proof. If ζ is a point in $\mathbb{S}^1$, then $i\zeta$ is the unitary vector tangent to $\mathbb{S}^1$ at ζ. Thus the velocity vector of the curve $f|_{\mathbb{S}^1}$ in $f(\zeta)$ is the image by $f'(\zeta)$ of $i\zeta$, that is, $f'(\zeta)i\zeta$. The winding number of the curve $f'(\zeta)i\zeta$ is the number of zeros on $\mathbb{D}$ of the function $izf'(z)$, and this number is equal to 1 if and only if $f' \neq 0$ on $\mathbb{D}$. $\square$

Lemma 3.5. *Let M be a complex manifold and $D \subset M$ a singular disc. Then there exists a holomorphic injective map $g : E \to M$, where $E = \mathbb{D}$ or $\mathbb{C}$, such that $g(E) = D$.*

[1]The degree of a parameterized regular curve in the plane is defined as the winding number around 0 of its velocity vector.

Proof. Let S be the set of singularities of D. Observe that $D\backslash S$ is homeomorphic to a subset of the plane. Then, since $D\backslash S$ is a Riemann surface, we have that $D\backslash S$ is actually holomorphically equivalent to a proper subset of the Riemann sphere (see [10]). Thus, there is a biholomorphism

$$g : U \to D\backslash S$$

where U is a domain in $\mathbb{C}P^1$. Consider $p \in S$. Since D is a singular disc, there is a holomorphic injective map $h : \mathbb{D} \to \mathcal{D}$, $h(0) = p$, where $\mathcal{D}$ is disc in D. We may assume that h is regular on $\overline{\mathbb{D}}\backslash\{0\}$. Then p is the unique singularity in $\overline{\mathcal{D}} \subset D$ and therefore the holomorphic function

$$g^{-1} \circ h : \overline{\mathbb{D}}\backslash\{0\} \to U$$

is well defined. Since $g^{-1} \circ h$ is injective, we have that $g^{-1} \circ h(\mathbb{S}^1)$ is a Jordan curve, hence it divides the sphere $\mathbb{C}P^1$ in two domains both homeomorphic to a disc. Then the image of $g^{-1} \circ h$ is contained in one of these domains and it follows by Riemann's extension theorem that $g^{-1} \circ h$ extends to $\mathbb{D}$ as a one to one map. Then, since $h : \mathbb{D} \to \mathcal{D}$ is a homeomorphism, we have that g^{-1} extends to $\mathcal{D}$ as a homeomorphism. Therefore, g^{-1} extends to D as a homeomorphism onto a simply connected domain $\Omega \subset \mathbb{C}$. Then $U = \Omega\backslash S'$, where $S' = g^{-1}(S)$ is a discrete closed set and we have therefore that $g : \Omega\backslash S' \to M$ extends to Ω as a holomorphic injective function $g : \Omega \to M$. Since $\Omega \subset \mathbb{C}P^1$ is simply connected and clearly $\Omega \neq \mathbb{C}P^1$, the lemma follows from Riemann's Uniformization Theorem. ☐

Proof of Proposition 3.3.

Assertion 1. For all z, we have that D_z is a singular disc and the sets $D_z\backslash \operatorname{Sing}(\mathcal{F})$ are the nonsingular leaves of $\mathcal{F}$.

Proof. Let $x \in D_z$. Since S_z is a discrete closed subset of D_z, there is a disc $\mathcal{D} \subset D_z$ with $x \in \mathcal{D}$ such that $\mathcal{D}\backslash\{x\} \subset D_z\backslash S_z$. Then, from hypothesis (iii), $\mathcal{D}\backslash\{x\}$ is contained in a leaf of $\mathcal{F}$. If $\mathcal{D}$ is small enough, we may think that $\mathcal{D}$ is contained in $\mathbb{C}^n$. Hence, by applying Lemma 3.5, there exists a holomorphic injective map $g : \mathbb{D} \to M$ with $g(\mathbb{D}) = \mathcal{D}$. Since that $x \in D_z$ was arbitrary, it follows that D_z is a singular disc.

Let L be a leaf of $\mathcal{F}$ and suppose that $x \in L \cap (D_z\backslash \operatorname{Sing}(\mathcal{F}))$ for some z. Take $\mathcal{D} \subset D_z$ as above. We assume $\mathcal{D}$ small enough such that it is contained in a neighborhood U of x where $\mathcal{F}$ is trivial and given by the submersion f. Then $\mathcal{D}\backslash\{x\}$ is contained in a leaf of $\mathcal{F}|_U$ and f is therefore constant over $\mathcal{D}\backslash\{x\}$. Hence, by continuity, f is constant over $\mathcal{D}$. Then $\mathcal{D}$ is contained in a leaf of $\mathcal{F}|_U$ and we have therefore $\mathcal{D} \subset L$. Thus we have $\mathcal{D} \subset L \cap (D_z\backslash \operatorname{Sing}(\mathcal{F}))$. It follows that $L \cap (D_z\backslash \operatorname{Sing}(\mathcal{F}))$ is an open subset of both L and $D_z\backslash \operatorname{Sing}(\mathcal{F})$ for all L and z. Now, fix a leaf L. Since the intersection of L with any $D_z\backslash \operatorname{Sing}(\mathcal{F})$ is open in L, it follows by connectedness that L is contained in a unique $D_z\backslash \operatorname{Sing}(\mathcal{F})$. For this $D_z\backslash \operatorname{Sing}(\mathcal{F})$, we also have that its intersection with any leaf is open in $D_z\backslash \operatorname{Sing}(\mathcal{F})$. Again by connectedness $D_z\backslash \operatorname{Sing}(\mathcal{F})$ is contained in a unique leaf, thus we necessarily have $D_z\backslash \operatorname{Sing}(\mathcal{F}) \subset L$ and it follows that $D_z\backslash \operatorname{Sing}(\mathcal{F}) = L$. Therefore Assertion 1 is proved.

Fix $p \in M$. We have $p \in D_{z'}$ for some $z' \in \Sigma$. Take p' in $D_{z'} \backslash S_{z'}$. From hypothesis (iii), p' is a regular point of $\mathcal{F}$. We have $p' = h(z', t')$ with $t' \in D$. If $B \subset \Sigma$ is a ball containing z', then $\Sigma_0 := B \times \{t'\}$ is a $(n-1)$ ball passing through (z', t'). We assume B small enough such that $\overline{\Sigma}_0$ is mapped by h into a neighborhood W of p' where $\mathcal{F}$ is equivalent to a product foliation. Let $\widetilde{\Sigma}$ (submanifold of W) be a global transversal to $\mathcal{F}|_W$. If w is a point contained in $\overline{h(\Sigma_0)}$, the leaf of $\mathcal{F}|_W$ passing through it intersects $\widetilde{\Sigma}$ in a unique point $\psi(w)$. We claim that ψ is a homeomorphism of $\overline{h(\Sigma_0)}$ onto its image. Since $\overline{h(\Sigma_0)}$ is compact, it suffices to prove that ψ is injective on $\overline{h(\Sigma_0)}$. Suppose that w_1 and w_2 are two points in $\overline{h(\Sigma_0)}$ contained in the same leaf L of $\mathcal{F}|_W$. From Assertion 1, we have that $L \subset D_z$ for some z. Then $h^{-1}(L) \subset \{z\} \times D$, hence $h^{-1}(w_1)$ and $h^{-1}(w_2)$ are two different points in the intersection of $(z \times D)$ with $\overline{\Sigma}_0$, which is a contradiction because $\overline{\Sigma}_0 \subset \Sigma \times \{t'\}$ intersects $(z \times D)$ only at (z, t').

We redefine $\widetilde{\Sigma}$ as $\widetilde{\Sigma} = \psi(h(\Sigma_0))$. Then for all $z \in B$ we have that D_z intersects $\widetilde{\Sigma}$ at the unique point $\psi(h(z, t_0))$; otherwise, if D_z intersects $\widetilde{\Sigma}$ in two different points x_1 and x_2, then $h^{-1}\psi^{-1}(x_1)$ and $h^{-1}\psi^{-1}(x_2)$ would be two different points in the intersection of $\{z\} \times \mathbb{D}$ with Σ_0.

We may define the map

$$g : V = h(B \times \mathbb{D}) \to \widetilde{\Sigma},$$

$$g(D_z) = D_z \cap \widetilde{\Sigma}.$$

By Assertion 1, each leaf of $\mathcal{F}$ is contained in some D_z. Then g is constant along the leaves. Therefore, since the restriction of g to any transversal is a holonomy map, we have that g is holomorphic on $V \backslash \operatorname{Sing}(\mathcal{F})$. Actually, since $\operatorname{Sing}(\mathcal{F})$ has codimension ≥ 2, g is holomorphic on V.

Consider $x \in \widetilde{\Sigma} \backslash g(\operatorname{Sing}(\mathcal{F}))$. Then $D = g^{-1}(x)$ does not intersect $\operatorname{Sing}(\mathcal{F})$. Clearly D is equal to some D_z. Then, by Assertion 1, $D \backslash \operatorname{Sing}(\mathcal{F}) = D$ is a leaf of $\mathcal{F}$. Thus, we conclude that for all $x \in \widetilde{\Sigma} \backslash g(\operatorname{Sing}(\mathcal{F}))$, the leaf passing through x is simply connected. Moreover, since $\operatorname{Sing}(\mathcal{F})$ has codimension ≥ 2, we have that $g(\operatorname{Sing}(\mathcal{F}))$ has codimension ≥ 1 in $\widetilde{\Sigma}$ and we have therefore that:

Assertion 2. For all x in a dense subset of $\widetilde{\Sigma}$, the leaf passing through x is simply connected.

Let Z be a holomorphic vector field which generates $\mathcal{F}$ on V and φ the local complex flow of Z. Let L be a leaf of $\mathcal{F}|_V$ and let x_L be its intersection with $\widetilde{\Sigma}$ ($g(L) = \{x_L\}$). There exists $\varepsilon_L > 0$ such that $\varphi(x_L, *)$ maps the disc $|t| < \varepsilon_L$ biholomorphically onto a neighborhood D_L of x_L in L. Thus, given any x in D_L there exists a unique $\tau_L(x)$ with $|\tau_L(x)| < \varepsilon_L$ such that $\varphi(x_L, \tau_L(x)) = x$. The function $\tau_L : D_L \to \mathbb{C}$ is the complex time between x_L and x. Clearly τ_L is holomorphic on D_L.

Assertion 3. The function τ_L can be analytically continued on L along any path $\gamma : [0,1] \to L$ with $\gamma(0) = x_L$.

Proof. Since γ does not intersect $\operatorname{Sing}(\mathcal{F})$ there exists $\delta > 0$ such that for all x in $\gamma([0,1])$, the map $\varphi(x, *)$ is a biholomorphism between $\mathbb{D}_{2\delta}$ and its image. Denote x_L

by x_0 and let $0 = s_0 < s_1 < \cdots < s_r = 1$ and $x_1 = \gamma(s_1), \ldots, x_r = \gamma(s_r)$ be such that:

(i) The open sets $\varphi(x_i, \mathbb{D}_\delta)$ for $i = 0, \ldots, r$ cover $\gamma([0,1])$.

(ii) x_i is contained in $\varphi(x_{i-1}, \mathbb{D}_\delta)$ for $i = 1, \ldots, r$.

For each $i = 0, \ldots, r$ let $\tau_i' : \varphi(x_i, \mathbb{D}_{2\delta}) \to \mathbb{D}_{2\delta}$ be defined by $\varphi(x_i, \tau_i'(x)) = x$. Let $x \in \varphi(x_{i-1}, \mathbb{D}_\delta) \cap \varphi(x_i, \mathbb{D}_\delta)$. Let $t_i = \tau_{i-1}'(x_i)$ for $i = 1, \ldots, r$ and define $t_0 = 0$. Clearly, $|t_i|$ and $|\tau_i'(x)|$ are less than δ, hence $|t_i + \tau_i'(x)| < 2\delta$ and we have that

$$\begin{aligned}
\varphi(x_{i-1}, t_i + \tau_i'(x)) &= \varphi(\varphi(x_{i-1}, t_i), \tau_i'(x)) \\
&= \varphi(\varphi(x_{i-1}, \tau_{i-1}'(x_i)), \tau_i'(x)) \\
&= \varphi(x_i, \tau_i'(x)) \\
&= x.
\end{aligned}$$

Then, by definition of τ_{i-1}' we obtain:

$$t_i + \tau_i'(x) = \tau_{i-1}'(x). \tag{5}$$

For each $i = 1, \ldots, r$ let τ_i be the holomorphic function on $\varphi(x_i, \mathbb{D}_\delta)$ defined by

$$\tau_i = \tau_i' + t_0 + \cdots + t_i.$$

By using (5) we deduce that $\tau_{i-1} = \tau_i$ on $\varphi(x_{i-1}, \mathbb{D}_\delta) \cap \varphi(x_i, \mathbb{D}_\delta)$. Moreover, it follows from the definition that τ_0 is equal to τ_L in a neighborhood of $x_0 = x_L$. Therefore, τ_0,...,τ_r give an analytic continuation of τ_L along γ.

Assertion 4. Let L be any leaf of $\mathcal{F}|_V$ and let $\gamma', \gamma'' : [0,1] \to L$ be paths such that $\gamma'(0) = \gamma''(0) = x_L$ and $\gamma'(1) = \gamma''(1) = x \in L$. Let τ_L' be the analytic continuation of τ_L along γ' and let τ_L'' be the analytic continuation of τ_L along γ''. Then $\tau_L'(x) = \tau_L''(x)$. Thus, τ_L extends as a holomorphic function on L. Therefore we may define $\tau : V \setminus \operatorname{Sing}(\mathcal{F}) \to \mathbb{C}$ by $\tau = \tau_L$ on L. Then τ is holomorphic on $U \setminus \operatorname{Sing}(\mathcal{F})$ and extends to U because $\operatorname{Sing}(\mathcal{F})$ has codimension ≥ 2. Moreover, if restricted to a leaf, τ is a regular map. In particular, τ is a submersion on $U \setminus \operatorname{Sing}(\mathcal{F})$.

Proof. Fix L and denote x_L by x_0. Let $0 = s_0 < \cdots < s_r = 1$, let $\Sigma_0, \ldots, \Sigma_r$ be transversals to the foliation at the points $x_0, x_1 = \gamma(s_1), \ldots, x_r = \gamma(s_r)$ respectively, and let $\delta > 0$ with the following properties:

(i) $\Sigma_0 \subset \widetilde{\Sigma}$.

(ii) The flow φ maps $\Sigma_i \times \mathbb{D}_{2\delta}$ biholomorphically onto its image, for all $i = 0, \ldots, r$.

(iii) The transversal Σ_i is contained in $\varphi(\Sigma_{i-1} \times \mathbb{D}_\delta)$, for all $i = 1, \ldots, r$.

(iv) For all $i = 1, \ldots, r$ we have that $\Sigma_i = h_i(\Sigma_0)$, where h_i is the holonomy map along γ.

Denote by V' the union of the sets $\varphi(\Sigma_i \times \mathbb{D}_\delta)$ for $i = 0, \ldots, r$. Consider $x \in V'$ and let L_x be the leaf passing through x. Let $k \in \{0, \ldots, r\}$ be such that $x \in \varphi(\Sigma_k \times \mathbb{D}_\delta)$. Then L_x intersects Σ_k and it follows from hypothesis (iv) that L_x intersects each Σ_i. Since $\Sigma_0 \subset \widetilde{\Sigma}$ we have that L_x intersects Σ_0 in a unique point and, by (iv), the same holds for each Σ_i. Then we may define $\rho_i : V' \to \Sigma_i$ such that $\rho_i(x)$ is the point of intersection between L_x and Σ_i. Let $\tau_i'(x) \in \mathbb{D}_\delta$ be defined by $\varphi(\rho_i(x), \tau_i'(x)) = x$. Since $\rho_i(x) \in \Sigma_i$, by hypothesis (iii) we have that $\rho_i(x) \in \varphi(\Sigma_{i-1} \times \mathbb{D}_\delta)$ for $i = 1, \ldots, r$. Then for $i = 1, \ldots, r$ we may define $t_i : V' \to \mathbb{D}_\delta$ as $t_i = \tau_{i-1}' \circ \rho_i$. Define $t_0 : V' \to \mathbb{D}_\delta$ as the zero function. Clearly, ρ_i, τ_i and t_i are holomorphic functions. We proceed as in the proof of Assertion 3. Let $x \in \varphi(\Sigma_i \times \mathbb{D}_\delta) \cap \varphi(\Sigma_{i-1} \times \mathbb{D}_\delta)$. Since $|t_i(x)|$ and $|\tau_i'(x)|$ are less than δ, then $|t_i(x) + \tau_i'(x)| < 2\delta$. Thus, by hypothesis (ii), $\varphi(\rho_{i-1}(x), t_i(x) + \tau_i'(x))$ is well defined and:

$$\begin{aligned}
\varphi(\rho_{i-1}(x), t_i(x) + \tau_i'(x)) &= \varphi(\varphi(\rho_{i-1}(x), t_i(x)), \tau_i'(x)) \\
&= \varphi(\varphi(\rho_{i-1}(x), \tau_{i-1}' \circ \rho_i(x)), \tau_i'(x)) \\
&= \varphi(\rho_i(x), \tau_i'(x)) \\
&= x.
\end{aligned}$$

Then by definition of τ_{i-1}' we deduce that

$$t_i(x) + \tau_i'(x) = \tau_{i-1}'(x).$$

Thus, the holomorphic functions on $\varphi(\Sigma_i \times \mathbb{D}_\delta)$ defined as

$$\tau_i(x) = \tau_i'(x) + t_0(x) + \cdots + t_i(x) \tag{6}$$

for each $i = 0, \ldots, r$ are such that

$$\tau_i = \tau_{i-1}$$

on $\varphi(\Sigma_i \times \mathbb{D}_\delta) \cap \varphi(\Sigma_{i-1} \times \mathbb{D}_\delta)$. Observe that for any leaf L', the restriction $\tau_0|_{L'}$ coincides with $\tau_{L'}$ on a neighborhood of $x_{L'}$. Then $\tau_0|_{L'}, \cdots, \tau_r|_{L'}$ give an analytic continuation of $\tau_{L'}$. Thus, $\tau_r|_L$ is the analytic continuation of τ_L along γ', hence $\tau_r(x) = \tau_L'(x)$. We denote τ_r by τ'. Analogously we construct τ'' for γ''. Then we have that $\tau''|_{L'}$ is an analytic continuation of $\tau_{L'}$ and, $\tau''|_L$ is the analytic continuation of τ_L along γ'', hence $\tau''(x) = \tau_L(x)$. By Assertion 2, we may take a sequence $\{x_k\}$ of points in Σ_0 with $x_k \to x$ as $k \to \infty$ and such that the leaf L_k passing through x_k is simply connected for all k. From above $\tau'|_{L_k}$ and $\tau''|_{L_k}$ are analytic continuations of τ_{L_k}. Since L_k is simply connected and, by Assertion 2, τ_{L_k} has an analytic continuation along any path, then $\tau'|_{L_k}$ and $\tau''|_{L_k}$ coincide on a neighborhood of x_k. In particular, $\tau'(x_k) = \tau''(x_k)$. Making $k \to \infty$ it follows by continuity that $\tau'(x) = \tau''(x)$, that is, $\tau_L'(x) = \tau_L''(x)$. Therefore, τ_L extends to L.

We define $\tau : V \setminus \operatorname{Sing}(\mathcal{F}) \to \mathbb{C}$ by $\tau|_L = \tau_L$. From above, τ coincides with the holomorphic function τ' on a neighborhood of the point x (arbitrary point). Then τ is holomorphic. Finally, remember (equation 6) that on a neighborhood of any non singular point, τ is expressed as

$$\tau_r(x) = \tau_r'(x) + t_0(x) + \cdots + t_r(x).$$

If we restrict x to a leaf, the first term of the sum above is a regular map and the other terms are constants. Hence τ is a regular map of any leaf. This finishes the proof of Assertion 4.

Given $x \in \widetilde{\Sigma}$, we know that $g^{-1}(x)$ is equal to D_z for some z. We denote $g^{-1}(x)$ by D_x. Thus, we have $p \in D_x$ for $x = g(p)$. It follows from hypothesis (iii) that there is a disc $\mathcal{D}' \subset D_x$ containing p such that $\mathcal{D}' \backslash \{p\}$ is contained in a leaf. Lemma 2.5 implies that there is a holomorphic bijective map $f : \Omega \to \mathcal{D}'$, $f(0) = p$, where $\Omega \subset \mathbb{C}$ is a disc containing $\mathbb{D}$. Thus if $\mathcal{D} = f(\mathbb{D})$, we have that $f : \overline{\mathbb{D}} \to \overline{\mathcal{D}}$ is holomorphic and regular on $\overline{\mathbb{D}} \backslash \{0\}$. Since $\overline{\mathcal{D}} \backslash \{p\}$ is contained in a leaf and by Assertion 3 we have that τ is a submersion on $U \backslash \operatorname{Sing}(\mathcal{F})$, then there exists a neighborhood V of $\partial\Delta$ on which τ defines a foliation by transversal balls along $\partial\Delta$. If we denote by Σ_ζ the transversal passing trough $\zeta \in \partial\Delta$ we have that τ is constant along Σ_ζ. Recall that $y \in \widetilde{\Sigma}$ is the unique point in the intersection of D_y and $\widetilde{\Sigma}$. It follows from the transversal uniformity of the foliation that if $y \in \widetilde{\Sigma}$ is close to x then D_y intersects only one time each transversal Σ_ζ. Let $\theta_y(\zeta)$ be the intersection of D_y with Σ_ζ. Since $\theta_y(\zeta)$ and ζ are both contained in Σ_ζ, we have that $\tau(\theta_y(\zeta)) = \tau(\zeta)$ for all $\zeta \in \partial\Delta$. Note that $\theta_y := \theta_y(\partial\Delta)$ is a smooth Jordan curve in D_y. By Assertion 2, we may choose y such that D_y is a leaf. We consider $\mathcal{D}_y \subset D_y$, the regular disc bounded by θ_y.



Let $f_y : \mathbb{D} \to \mathcal{D}_y$ be a uniformization map. Since θ_y is a smooth Jordan curve, f_y extends as a diffeomorphism $f_y : \overline{\mathbb{D}} \to \overline{\mathcal{D}}_y$ (see [8], p.323). By Assertion 3, we have that τ is regular on $\overline{\mathcal{D}}_y$. It follows that $\tau \circ f_y : \overline{\mathbb{D}} \to \mathbb{C}$ is a regular map. Therefore, by Lemma 3.4, the curve $\tau \circ f_y : \mathbb{S}^1 \to \mathbb{C}$ has degree 1. Remember that $\tau(\theta_y(\zeta)) = \tau(\zeta)$ for all $\zeta \in \partial\Delta$, thus $\tau(\partial\mathcal{D}_y) = \tau(\partial\mathcal{D})$. Then

$$\tau \circ f_y(\mathbb{S}^1) = \tau(\partial\mathcal{D}_y) = \tau(\partial\mathcal{D}) = \tau \circ f(\mathbb{S}^1).$$

Therefore $\tau \circ f : \mathbb{S}^1 \to \mathbb{C}$ is only a reparametrization of $\tau \circ f_y : \mathbb{S}^1 \to \mathbb{C}$, hence $\tau \circ f : \mathbb{S}^1 \to \mathbb{C}$ is regular and has degree 1. Again by Lemma 3.4, $\tau \circ f : \overline{\mathbb{D}} \to \mathbb{C}$ is also a regular map and in particular, $\tau \circ f$ is locally injective. Therefore there exists a disc $\mathrm{U} \subset \mathbb{D}$, centered at 0, such that $\tau \circ f$ is injective on $\overline{\mathrm{U}}$. Then

$$\tau \circ f(\partial \mathrm{U})$$

is a Jordan curve in $\mathbb{C}$. We also denote $f(\mathrm{U})$ by $\mathcal{D}$. Again, let Σ_ζ be the transversal ball through $\zeta \in \partial\mathcal{D}$. Proceeding as above, if Σ' is a small enough ball in $\widetilde{\Sigma}$ containing $x = g(p)$, we obtain that for all $y \in \Sigma'$ the set D_y intersects each Σ_ζ at the unique point $\theta_y(\zeta)$. Thus we have the Jordan curve θ_y in D_y such that $\tau(\theta_y) = \tau(\partial D)$. Remember that $\tau(\partial D) = \tau \circ f(\partial \mathrm{U})$ is a Jordan curve in $\mathbb{C}$. It follows that $\tau(\theta_y)$ is Jordan curve in $\mathbb{C}$ for all y. Let $\mathcal{D}_y \subset D_y$ be the disc bounded by θ_y. Since D_y is a singular disc, by Lemma 3.5, there is an injective holomorphic map $f_y : E \to M$, where $E = \mathbb{D}$ or $\mathbb{C}$, such that $f_y(E) = D_y$. Let $\Omega_y \subset E$ be such that $f_y(\Omega_y) = \mathcal{D}_y$. Clearly Ω_y is a disc and $f_y(\partial\Omega_y) = \partial\mathcal{D}_y$. Then

$$\tau \circ f_y(\partial\Omega_y) = \tau(\partial\mathcal{D}_y)$$

is, from above, a Jordan curve in $\mathbb{C}$. Hence we deduce that the holomorphic function $\tau \circ f_y : \Omega_y \to \mathbb{C}$ is injective on $\overline{\Omega}_y$. Thus, since f_y is injective, we conclude that

$$\tau : \overline{\mathcal{D}}_y \to \mathbb{C}$$

is injective for all $y \in \Sigma'$.

Denote by W the union of the discs $\mathcal{D}_y$ for all $y \in \Sigma'$. It is easy to see that W is a neighborhood of p. Define

$$F : \overline{W} \to \widetilde{\Sigma} \times \mathbb{C}$$

$$F(w) = (g(w), \tau(w))$$

Assertion 5. F is a biholomorphism between W and its image.

Proof. Clearly F is holomorphic on W. We shall prove that F injective on $\overline{W}$. Suppose $F(w) = F(w')$. Then $g(w) = g(w') = y$, hence $w, w' \in D_y$ and, since $\overline{W} \cap D_y = \overline{\mathcal{D}}_y$, we have $w, w' \in \overline{\mathcal{D}}_y$. On the other hand $\tau(w) = \tau(w')$ and since τ is injective on $\overline{\mathcal{D}}_y$ we conclude that $w = w'$. Now, since $\overline{W}$ is compact, F is a homeomorphism onto its image and it follows that F is a biholomorphism.

Now, we will prove that $p \in W$ is regular for $\mathcal{F}$. Let $\mathcal{N}$ be the regular foliation on $\widetilde{\Sigma} \times \mathbb{C}$ whose leaves are the sets $\{*\} \times \mathbb{C}$. Let $\mathcal{F}'$ be the pull-back foliation of $\mathcal{N}$ by the biholomorphism F. Then $\mathcal{F}'$ is regular and it is easy to see that $\mathcal{F}'$ coincides with $\mathcal{F}$ out on a open set of W (out of $\mathrm{Sing}(\mathcal{F})$). Then $\mathcal{F}' = \mathcal{F}$ on W and $\mathcal{F}$ is therefore regular at p. Since $p \in U$ was arbitrary, we have proved that $\mathrm{Sing}(\mathcal{F})$ is empty. Then, from Assertion 1, the sets D_z are the leaves of $\mathcal{F}$. The proof of Proposition 3.3 is complete. ☐

Lemma 3.6. *Let M be a complex manifold and let $X \subset M$ be a subvariety of codimension $k \geq 1$. Let p be a point in X. Then there exist a path x_t, $t \in [0, 1)$, with $x_0 = p$ and such that x_t is out of X for all $t > 0$.*

Proof. Let U be a neighborhood of p in M and $f : U \to \mathbb{C}^r$ be such that $f^{-1}(0) = X \cap U$. There is a complex disc D passing through p such that $f|_D$ does not vanish, otherwise

$f^{-1}(0) \subset X$ contains a neighborhood of p in M. By reducing D, we may assume that $p \in D$ is the unique zero of $f|_D$. Then $D \backslash \{p\}$ is disjoint of X and it is sufficient to take any path x_t in D with $x_0 = p$ and $x_t \neq p$ for all $t > 0$. $\square$

Proof of Theorem 3.1.

Assertion 1. Let $z \in \Sigma$ such that D_z is not contained in X. Then D_z is contained in a leaf of $\mathcal{F}$.

Proof. Take $t_0 \in D_z$ such that $h(z, t_0) \notin X$. Since X is closed in M, if Σ' is a small enough neighborhood (ball) of z in Σ, we have that $h(z', t_0) \notin X$ for all $z' \in \Sigma'$. Hence, for all $z' \in \Sigma'$ we have that $D_{z'}$ is not contained in X. Then, by hypothesis (ii), $S_{z'} := D_{z'} \cap X$ is discrete and $D_{z'} \backslash S_{z'}$ is contained in a nonsingular leaf of $\mathcal{F}$. Therefore, $\mathcal{F}$ restricted to $M' := h(\Sigma' \times D)$ satisfies the hypothesis of Proposition 3.3 and we have therefore that D_z is contained in a leaf of $\mathcal{F}$.

Assertion 2. Let $z \in \Sigma$ such that D_z is contained in X. Then D_z is a singular disc.

Proof. Let $x \in D_z$, $x = h(z, t)$. Let $\Sigma' \subset \Sigma$ be a neighborhood (a ball) of z and $D' \subset D$ be a neighborhood (a disc) of t. If Σ' and D' are small enough, we may assume that $M' := h(\Sigma' \times D')$ is a domain in $\mathbb{C}^n$. Since X has codimension ≥ 1, by Lemma 3.6, there is a path $x_s = h(z_t, t_s)$ in M' such that $x_0 = x$ and $x_s \notin X$ for all $s > 0$. Then $D_s := D_{z_s}$ is not contained in X for all $s > 0$ and it follows by Assertion 1 that D_s is contained in a leaf. Hence D_s is a regular disc for all $s > 0$. Then, we may apply Lemma 2.1 to the family of discs D_s and conclude that $D_z = D_0$ is a singular disc.

Assertion 3. Let z be such that $D_z \subset X$. Let S_z be the set of singularities of the singular disc D_z. Then $D_z \backslash S_z$ is contained in a leaf of $\mathcal{F}$.

Proof. By Assertion 2, if D_z is not contained in X we have that D_z is contained in a leaf of $\mathcal{F}$. Therefore, the hypothesis of Proposition 3.2 holds for $\mathcal{F}$ and Assertion 3 follows.

Let z be such that D_z is not contained in X. By hypothesis (iii) of 3.1, we have that $S_z := D_z \cap X$ is discrete and $D_z \backslash S_z$ is contained in a leaf of $\mathcal{F}$. From this and Assertion 3 we conclude: for all z there is a discrete set S_z such that $D_z \backslash S_z$ is contained in a leaf of $\mathcal{F}$. Therefore the hypothesis of Proposition 3.3 holds and Theorem 3.1 follows. $\square$

4 The algebraic multiplicity and the Chern class of the tangent bundle of the strict transform

Let $\mathcal{F}_0$, $\widetilde{\mathcal{F}}_0$ and h as in §1.

Proposition 4.1. *If h extends to the divisor as a homeomorphism between $\pi^{-1}(U)$ and $\pi^{-1}(\widetilde{U})$, then the extension also denoted by h is a topological equivalence between $\mathcal{F}_0$ and $\widetilde{\mathcal{F}}_0$.*

Proof. Is a direct application of Theorem 1.4. $\square$

Proof of Theorem 1.1. Suppose that $\mathcal{F}$ is generated on U by the holomorphic vector field

$$V=\sum_{i=1}^{n} a_i\frac{\partial}{\partial z_i},\ a_i\in\mathcal{O}_U,\ g.c.d.(a_1,a_2,\ldots,a_n)=1.$$

For each $j=1,2,\ldots,n$, let $U_j=(x_j\neq 0)$ and $U'_j=\pi^{-1}(U_j)$. Let $V_j=\pi^*(V|_{U_j})$. If $(x^j_1,\ldots,x^j_n)$ are coordinates on U'_j such that

$$\pi(x^j_1,\ldots,x^j_n)=(x^j_jx^j_1,\ldots,x^j_j,\ldots,x^j_jx^j_n),$$

then

$$V_j=a_j\frac{\partial}{\partial x^j_j}+\sum_{i=1,i\neq j}^{n}\frac{a_i-x^j_ia_j}{x^j_j}\frac{\partial}{\partial x^j_i},$$

where $a_i=a_i\circ\pi$ for $i=1,\ldots,n$. On U'_j, $\mathcal{F}_0$ is defined by the vector field

$$W_j=\frac{1}{(x^j_j)^{r-\xi}}V_j,$$

where r is the algebraic multiplicity of V at $0\in\mathbb{C}^n$ and $\xi=1$ or 0 depending on the divisor being invariant or not by $\mathcal{F}_0$. Evidently $V_i=V_j$ on $U'_i\cap U'_j$. Then

$$W_i=\left(x^j_j/x^i_i\right)^{r-\xi}W_j\quad\text{on}\quad U'_i\cap U'_j.$$

It follows from this equation that the tangent bundle $T\mathcal{F}_0$ of $\mathcal{F}_0$ is isomorphic to $L^{\xi-r}$, where L is the line bundle associated to the divisor $E=\pi^{-1}(0)$. Then the Chern class $c(T\mathcal{F}_0)$ of $T\mathcal{F}_0$ is equal to $(\xi-r)c(L)$. It is natural consider E as an element in $H_{n-2}(U',\mathbb{Z})$, where $U'=\pi^{-1}(U)$. We know that $c(L)$ is equal to $d(E)\in H^2(U',\mathbb{Z})$, the dual of E. Therefore

$$c(T\mathcal{F}_0)=(\xi-r)d(E).$$

On the other hand, make $\widetilde{U}'=\pi^{-1}(\widetilde{U})$ and observe that the divisor E is invariant by $\mathcal{F}_0$ if and only if it is by $\widetilde{\mathcal{F}}_0$. Then analogously we have

$$c(T\widetilde{\mathcal{F}}_0)=(\xi-\widetilde{r})\widetilde{d}(E),$$

where $\widetilde{r}$ is the algebraic multiplicity of $\widetilde{\mathcal{F}}$ and $\widetilde{d}(E)\in H^2(\widetilde{U}',\mathbb{Z})$ is the dual of E. By Proposition 4.1 we have that $h:U'\to\widetilde{U}'$ is a topological equivalence between $\mathcal{F}_0$ and $\widetilde{\mathcal{F}}_0$. Then Theorem 1.3 implies that

$$(\xi-r)h^*(d(E))=(\xi-\widetilde{r})\widetilde{d}(E).\tag{7}$$

We may assume that U is a ball in $\mathbb{C}^n$. Thus, we have that U' is a tubular neighborhood of E and therefore $H^2(U',\mathbb{Z})\simeq\mathbb{Z}$. Since the cohomology is invariant by homeomorphisms, we also have $H^2(\widetilde{U}',\mathbb{Z})\simeq\mathbb{Z}$. Can be proved that $d(E)$ and $\widetilde{d}(E)$ are generators of $H^2(U',\mathbb{Z})$ and $H^2(\widetilde{U}',\mathbb{Z})$ respectively. Then we have that $h^*(d(E))=\widetilde{d}(E)$ or $h^*(d(E))=-\widetilde{d}(E)$. By using this in (7) we obtain $r=\widetilde{r}$ or $r+\widetilde{r}=2\xi$. The second possibility implies $r = \widetilde{r} = \xi$, since $r \geq 1$, $\widetilde{r} \geq 1$ and $\xi = 1$ or 0. Therefore we conclude that $r = \widetilde{r}$. $\square$

Remark. Under the hypothesis of Theorem 1.1, we have another invariants. The restriction of $\mathcal{F}_0$ to the divisor is a foliation with $\mathrm{Sing}(\mathcal{F}_0)$ as singular set. It is well known that this foliation coincides out of the singular set with a unique foliation $\mathcal{N}$ of codimension ≥ 2 in the divisor (the saturated foliation). We will say that $\mathcal{N}$ is the foliation induced by $\mathcal{F}_0$ in the divisor. Let $\widetilde{\mathcal{N}}$ be the foliation induced by $\widetilde{\mathcal{F}}_0$ in the divisor. It follows from Theorem 1.4 that $\mathcal{N}$ and $\widetilde{\mathcal{N}}$ are topologically equivalent. Thus, since the divisor is isomorphic to $\mathbb{P}^{n-1}$, Theorem 1.3 implies that $d(\mathcal{N}) = d(\widetilde{\mathcal{N}})$. In other words, the degree of the foliation induced in the divisor is invariant.

From above, $\mathcal{F}_0$ is generated by the holomorphic vector fields W_i and

$$W_i = \left(x_j^j / x_i^i\right)^{r-\xi} W_j \quad \text{on} \quad U_i' \cap U_j',$$

where $\xi = 1$ or 0. Let $x \in U_i' \cap U_j'$. Let $x^i = (x_1^i, \ldots, x_n^i)$ be the coordinates of x in U_i' and let $x^j = (x_1^j, \ldots, x_n^j)$ be the coordinates of x in U_j'. Since $\pi(x^i) = \pi(x^j)$, we have that

$$(x_i^i x_1^i, \ldots, x_i^i, \ldots, x_i^i x_n^i) = (x_j^j x_1^j, \ldots, x_j^j, \ldots, x_j^j x_n^j),$$

hence $x_j^j / x_i^i = x_j^i$. Replacing in last equation we obtain:

$$W_i = \left(x_j^i\right)^{r-\xi} W_j \quad \text{on} \quad U_i' \cap U_j'. \tag{8}$$

Observe that $\pi^{-1}(0) \cap U_i'$ is represented by $(x_i^i = 0)$. Recall that $\pi^{-1}(0)$ is canonically isomorphic to $\mathbb{P}^{n-1}$. A point p in $\pi^{-1}(0) \cap U_i'$ given by

$$(x_1^i(p), \ldots, 0_i, \ldots, x_n^i(p))$$

is represented in homogeneous coordinates by

$$[z_1 : \cdots : z_n](p) = [x_1^i(p) : \cdots : 1_i : \cdots : x_n^i(p)],$$

hence $x_j^i(p) = (z_j/z_i)(p)$. Thus, if $\mathcal{U}_i = U_i' \cap \pi^{-1}(0)$ and $J_i = W_i|_{\mathcal{U}_i}$, it follows from (8) that

$$J_i = (z_j/z_i)^{r-\xi} J_j \quad \text{on} \quad \mathcal{U}_i \cap \mathcal{U}_j. \tag{9}$$

Let S be the union of the components of codimension 2 of $\mathrm{Sing}\,(\mathcal{F}_0)$. Then S is the codimension 1 part (respect to the divisor) of the zero set of $\{J_i\}$. Each J_i may be expressed as $J_i = f_i Z_i$, where f_i is a holomorphic function on $\mathcal{U}_i$ and the vector field Z_i has singular set of codimension ≥ 2. It follows from (9) that

$$Z_i = (f_j/f_i)\,(z_j/z_i)^{r-\xi} Z_j \;\text{ on }\; \mathcal{U}_i \cap \mathcal{U}_j.$$

From this equation, it is not difficult to conclude that

$$r = d(\mathcal{N}) - \deg(S) - 1 + \xi,$$

where $\deg(S)$ is the degree of S as a divisor of $\pi^{-1}(0)$. Then, since the algebraic multiplicity y and the degree of the foliation induced in the divisor are invariants, we deduce that the degree of the codimension 1 part of the singular set of the strict transform is also an invariant. Moreover it is not difficult to see that $h(S) = \widetilde{S}$, where $\widetilde{S}$ is the union of the components of codimension 2 of $\mathrm{Sing}(\widetilde{\mathcal{F}}_0)$.

5 The case C^1

In this section we prove Theorem 1.2. In view of Theorem 1.1, it is sufficient to show the following.

Proposition 5.1. *Let $\mathcal{F}$ and $\widetilde{\mathcal{F}}$ be two foliations by curves of neighborhoods U and $\widetilde{U}$ of $0 \in \mathbb{C}^n$. Let $\mathrm{h} : U \to \widetilde{U}$ be a C^1 equivalence. Let $h : \pi^{-1}(U \backslash \{0\}) \to \pi^{-1}(\widetilde{U} \backslash \{0\})$ be as before. Then h can be extended to the divisor as a homeomorphism between $\pi^{-1}(U)$ and $\pi^{-1}(\widetilde{U})$.*

We start the proof.

Proposition 5.2. *Under the conditions of Proposition 5.1, we have that $d\,\mathrm{h}(0) : \mathbb{C}^n \to \mathbb{C}^n$ maps complex lines onto complex lines. Furthermore, if $J : \mathbb{C}^n \to \mathbb{C}^n$ is the conjugation $J(z) = \bar{z}$, then either $d\,\mathrm{h}(0) : \mathbb{C}^n \to \mathbb{C}^n$ is a $\mathbb{C}$-linear isomorphism, or $d\,\mathrm{h}(0) = Q \circ J$, where $Q : \mathbb{C}^n \to \mathbb{C}^n$ is a $\mathbb{C}$-linear isomorphism. Thus, $d\,\mathrm{h}(0)$ induces a diffeomorphism of $\mathbb{P}^{n-1}$ onto itself.*

Proof. let L be a complex line, $0 \in L \subset \mathbb{C}^n$. There exists $\mathbb{C}$-linear functions $A_i : \mathbb{C}^n \to \mathbb{C}$ for $i = 1, \ldots, (n-1)$, such that

$$L = \{z \in \mathbb{C}^n \,:\, A_i(z) = 0, \;\text{ for all }\; i = 1, 2, \ldots, (n-1)\}.$$

Let $V : U \to \mathbb{C}^n$ be a holomorphic vector field which generates $\mathcal{F}$. The set:

$$B = \{z \in \mathbb{C}^n \,:\, A_i \circ V(z) = 0, \;\text{for all }\; i = 1, 2, \ldots, (n-1)\}$$

is an analytic variety and it is easy to see that $0 \in B$. Then, there exists a complex curve contained in B and passing through 0. In particular there exists a sequence of points $z_k \in \mathbb{C}^n \backslash \{0\}$, $z_k \to 0$, such that $A_i \circ V(z_k) = 0$ for all $k \in \mathbb{N}$ and all $i = 1, 2, \ldots, (n-1)$. In other words, $T_{z_k}\mathcal{F} = L$ for all $k \in \mathbb{N}$. Now, since h is a C^1 equivalence, $d\,\mathrm{h}_{z_k}(T_{z_k}\mathcal{F}) = T_{\mathrm{h}(z_k)}\widetilde{\mathcal{F}}$, that is, $d\,\mathrm{h}_{z_k}(L) = T_{\mathrm{h}(z_k)}\widetilde{\mathcal{F}}$ is a complex line for all $k \in \mathbb{N}$. Making $k \to \infty$, since $\mathrm{h} \in C^1$ and the space of complex lines of $\mathbb{C}^n$ is compact, we obtain that $d\;h_0(L)$ is also a complex line. The second part of the proposition is an immediate consequence of the following lemma. □

Lemma 5.3. *Let $A : \mathbb{R}^{2n} \to \mathbb{R}^{2n}$ be a $\mathbb{R}$-linear isomorphism. Identify $\mathbb{R}^{2n}$ with $\mathbb{C}^n$ and assume that A maps complex lines onto complex lines. Then, either A is a $\mathbb{C}$-linear isomorphism, or $A = Q \circ J$ with $Q : \mathbb{C}^n \to \mathbb{C}^n$ a $\mathbb{C}$-linear isomorphism.*

Proof. Since A maps any complex line onto a complex line, for all $v \in \mathbb{C}^n \backslash \{0\}$ there exists $\theta(v) \in \mathbb{C} \backslash \{0\}$ such that $A(iv) = \theta(v) A(v)$. Let v_1 and v_2 be two $\mathbb{C}$-linearly independent vectors. Then

$$A(iv_1 + iv_2) = A(iv_1) + A(iv_2) = \theta(v_1)A(v_1) + \theta(v_2)A(v_2).$$

Moreover:

$$\begin{aligned} A(iv_1 + iv_2) &= A(i(v_1 + v_2)) = \theta(v_1 + v_2)A(v_1 + v_2) \\ &= \theta(v_1 + v_2)A(v_1) + \theta(v_1 + v_2)A(v_2). \end{aligned}$$

From the equations above, we obtain:

$$(\theta(v_1) - \theta(v_1 + v_2))A(v_1) + ((\theta(v_2) - \theta(V_1 + v_2))A(v_2) = 0. \tag{10}$$

Let L_1 and L_2 be the complex lines generated by v_1 and v_2 respectively. Since v_1 and v_2 are $\mathbb{C}$-linearly independent, we have that L_1 and L_2 are different. This implies, since A is an isomorphism, that $A(L_1)$ and $A(L_2)$ are different complex lines. Then, since $A(L_1)$ and $A(L_2)$ are generated by $A(v_1)$ and $A(v_2)$ respectively, we have that $A(v_1)$ and $A(v_2)$ are $\mathbb{C}$-linearly independent. Thus, it follows from equation (10) that

$$\theta(v_1) = \theta(v_1 + v_2) = \theta(v_2).$$

It is now easy to see that $\theta(v) = \theta_0,\ \forall v \in \mathbb{C}^n \backslash \{0\}$. We know that there exists two $\mathbb{C}$-linear transformations $P : \mathbb{C}^n \to \mathbb{C}^n$ and $Q : \mathbb{C}^n \to \mathbb{C}^n$ such that

$$A(z) = P(z) + Q(\bar{z}), \text{ for all } z \in \mathbb{C}^n.$$

Then

$$A(iz) = iP(z) - iQ(\bar{z}).$$

On the other hand

$$A(iz) = \theta_0 A(z) = \theta_0 P(z) + \theta_0 Q(\bar{z}), \text{ for all } z \in \mathbb{C}^n.$$

consequently

$$(\theta_0 - i)P(z) + (\theta_0 + i)Q(\bar{z}) = 0.$$

Since, as functions of z, $(\theta_0 - i)P$ and $(\theta_0 + i)Q \circ J$ are holomorphic and anti-holomorphic respectively, we have that

$$(\theta_0 - i)P \equiv 0, \ (\theta_0 + i)Q \circ J \equiv 0.$$

From this it is easy to see that either $P = 0$, or $Q = 0$. This proves the lemma. □

Definition 5.4. *Let $\{z_k\}$ be a sequence of points in $\mathbb{C}^n \backslash \{0\}$. Let L be a complex line in $\mathbb{C}^n$. We say that $\{z_k\}$ is tangent to L at 0 if $z_k \to 0$ and every accumulation point of $\{z_k/||z_k||\}$ is contained in L.*

Let $\pi : \widehat{\mathbb{C}^n} \to \mathbb{C}^n$ be the blow up at $0 \in \mathbb{C}^n$. We know that $\pi^{-1}(0)$ is naturally isomorphic to $\mathbb{P}^{n-1}$. Thus, for each $p \in \pi^{-1}(0)$ we denote by L_p the respective complex line in $\mathbb{C}^n$.

Proposition 5.5. *Let $\{p_k\}$ be a sequence of points in $\widehat{\mathbb{C}^n} \backslash \pi^{-1}(0)$. Then $p_k \to p \in \pi^{-1}(0)$ if and only if $\{\pi(p_k)\}$ is tangent to L_p at 0.*

Proof. Let

$$U_1 = \{(z_1, z_2, \ldots, z_n) \in \mathbb{C}^n : z_1 \neq 0\}.$$

There exist coordinates $(x_1, x_2, \ldots, x_n)$ on $\pi^{-1}(U_1)$ such that

$$\pi(x_1, x_2, \ldots, x_n) = (x_1, x_1 x_2, \ldots, x_1 x_n)$$

and $\pi^{-1}(0)$ is represented by $x_1 = 0$. Without lost of generality we may assume that $p = (0, 0, \ldots, 0)$, $L_p = \{z_2 = 0, ..., z_n = 0\}$, and $p_k = (x_1{}^k, x_2{}^k, \ldots, x_n{}^k) \in \pi^{-1}(U_1)$ for all $k \in \mathbb{N}$. Assume first that $p_k \to p$. Evidently, $\pi(p_k) \to 0$. On the other hand

$$\frac{\pi(p_k)}{||\pi(z_k)||} = \frac{x_1{}^k}{|x_1{}^k|} \frac{(1, x_2{}^k, \ldots, x_n{}^k)}{||(1, x_2{}^k, \ldots, x_n{}^k)||}$$

and since

$$\frac{(1, x_2{}^k, \ldots, x_n{}^k)}{||(1, x_2{}^k, \ldots, x_n{}^k)||} \to (1, 0, 0, \ldots, 0),$$

any accumulation point of $\{\pi(p_k)/||\pi(p_k)||\}$ have the form $\lambda(1, 0, \ldots, 0)$ with $\lambda \in \mathbb{C}\backslash\{0\}$. Therefore $\{\pi(p_k)\}$ is tangent to L_p in 0. Conversely, suppose that $\{\pi(p_k)\}$ is tangent to L_p in 0. Let $\pi(p_k) = z^k = (z_1^k, ..., z_n^k)$. Since any accumulation point of

$$\frac{z^k}{||z^k||} = (\frac{z_1^k}{||z^k||}, ..., \frac{z_n^k}{||z^k||})$$

is contained in $L_p = \{z_2 = 0, ..., z_n = 0\}$, Then for k big enough and all $j = 2, ..., n$ we have that $\frac{z_j^k}{||z^k||} < \epsilon$, with $\epsilon > 0$ arbitrary. Then

$$||z^k|| \leq |z_1^k| + \ldots + |z_n^k| \leq |z_1^k| + (n-1)\epsilon||z^k||,$$

hence $|z_1^k| \geq (1 - (n-1)\epsilon)||z^k||$ and therefore $z_1^k \neq 0$ if ϵ is small enough. Thus, we may assume that $p_k = (x_1^k, ..., x_n^k) \in \pi^{-1}(U_1)$. Then $(x_1^k, x_1^k x_2^k, \ldots, x_1^k x_n^k) \to 0$ and therefore $x_1{}^k \to 0$. On the other hand, since any accumulation point of

$$\frac{\pi(p_k)}{||\pi(p_k)||} = \frac{x_1{}^k}{|x_1{}^k|} \frac{(1, x_2{}^k, \ldots, x_n{}^k)}{||(1, x_2{}^k, \ldots, x_n{}^k)||}$$

is contained in $L_p \subset \mathbb{C}^n$, then the same property holds for the sequence

$$\frac{(1, x_2{}^k, \ldots, x_n{}^k)}{||(1, x_2{}^k, \ldots, x_n{}^k)||},$$

hence $x_i{}^k \to 0$ for all $i = 2, 3, \ldots, n$ and therefore $p_k \to p$.

proof of Proposition 5.1. Let $p \in \pi^{-1}(0)$ and $\{p_k\}$ any sequence of points in $\pi^{-1}(U)\backslash\pi^{-1}(0)$ such that $p_k \to p$.

Since $h \in C^1$, we have

$$\mathrm{h}(\pi(p_k)) = d\,\mathrm{h}_0(\pi(p_k)) + r(\pi(p_k)), \text{ where } \frac{r(\pi(p_k))}{||\pi(p_k)||} \to 0 \text{ as } k \to 0.$$

Then

$$\frac{\mathrm{h}(\pi(p_k))}{||\pi(p_k)||} = d\,\mathrm{h}_0\left(\frac{(\pi(p_k))}{||\pi(p_k)||}\right) + \frac{r(\pi(p_k))}{||\pi(p_k)||}. \tag{11}$$

By proposition 5.5, $\pi(p_k)$ is tangent to L_p at 0, hence any point of accumulation of the sequence $\{(\pi(p_k))/\|\pi(p_k)||\}$ is contained in L_p. Thus, it is easy to see from equation

(11) that any point of accumulation of the sequence $\{\mathrm{h}(\pi(p_k))/||\pi(p_k)||\}$ is contained in $d\,\mathrm{h}_0(L_p)$ and the same holds for the sequence

$$\frac{\mathrm{h}(\pi(p_k))}{||\,\mathrm{h}(\pi(p_k))||} = \frac{\mathrm{h}(\pi(p_k))}{||\pi(p_k)||}\frac{||\pi(p_k)||}{||\,\mathrm{h}(\pi(p_k))||}.$$

From proposition 5.2, we have that $d\,\mathrm{h}_0(L_p)$ is a complex line. Then $\{\mathrm{h}(\pi(p_k)\})$ is tangent to $d\,\mathrm{h}_0(L_p)$ at 0. It follows by proposition 5.5 that $\pi^{-1}\circ \mathrm{h}\circ\pi(p_k) = h(p_k) \to q$, where $q \in \pi^{-1}(0)$ is such that $L_q = d\,\mathrm{h}_0(L_p)$. We remark that $q = H(p)$, where H is the diffeomorphism of $\pi^{-1}(0) \simeq \mathbb{P}^{n-1}$ onto itself induced by $d\,\mathrm{h}_0$ (5.2). We define $h(p) = H(p)$ for all p in $\pi^{-1}(0)$. Finally, we prove that $h : \pi^{-1}(0) \to \pi^{-1}(0)$ is a homeomorphism. Let $p \in \pi^{-1}(0)$ and let p_k in $\widehat{\mathbb{C}^n}$ such that $p_k \to p$ as $k \to \infty$. Let x be a limit point of the sequence $\{h(p_k)\}$, that is, $x = \lim h(q_k)$, where $\{q_k\}$ is a subsequence of $\{p_k\}$. Clearly, there exists a subsequence $\{r_k\}$ of $\{q_k\}$ such that: either $\{r_k\}$ is contained in $\pi^{-1}(0)$, or $\{r_k\}$ is contained in $\pi^{-1}(U)\backslash\pi^{-1}(0)$. If $\{r_k\}$ is contained in $\pi^{-1}(0)$ we have that $h(r_k) = H(r_k)$ for all k and

$$x = \lim_{k\to\infty} h(q_k) = \lim_{k\to\infty} h(r_k) = \lim_{k\to\infty} H(r_k) = H(p) = h(p),$$

since H is continuous. On the other hand, if $\{r_k\}$ is contained in $\pi^{-1}(U)\backslash\pi^{-1}(0)$, we have from above that

$$x = \lim_{k\to\infty} h(r_k) = H(p) = h(p).$$

Therefore h is continuous. Analogously we prove that h^{-1} is also continuous.

Part II

The differentiable case

6 Introduction

Let $\mathcal{F}$ be a holomorphic foliation by curves of a neighborhood U of $0 \in \mathbb{C}^2$ with a unique singularity at $0 \in \mathbb{C}^2$. We assume that $\mathcal{F}$ is generated by the holomorphic vector field

$$V = \sum_{i=1}^{n} a_i \frac{\partial}{\partial z_i},\ a_i \in \mathcal{O}_U,\ g.c.d.(a_1, a_2, \ldots, a_n) = 1.$$

The algebraic multiplicity of $\mathcal{F}$ (at $0 \in \mathbb{C}^2$) is the minimum vanishing order at $0 \in \mathbb{C}^2$ of the functions a_i. Let $\widetilde{\mathcal{F}}$ be a holomorphic foliation by curves of a neighborhood $\widetilde{U}$ of $0 \in \mathbb{C}^2$ and Let $\mathrm{h} : U \to \widetilde{U}$ be a topological equivalence between $\mathcal{F}$ and $\widetilde{\mathcal{F}}$, that is, a homeomorphism taking leaves of $\mathcal{F}$ to leaves of $\widetilde{\mathcal{F}}$. A natural question, posed by J.F.Mattei is: are the algebraic multiplicities of $\mathcal{F}$ and $\widetilde{\mathcal{F}}$ the same?. In [2], the authors give a positive answer if $\mathcal{F}$ is a *generalized curve*, that is, if the desingularization of $\mathcal{F}$ does not contain complex saddle-nodes. If $\mathcal{F}$ is *dicritical*, that is, after a blow up the exceptional divisor is not invariant by the strict transform of $\mathcal{F}$, the conjecture is also true: in this case, it is not difficult to show that the algebraic multiplicity of $\mathcal{F}$ is equal to the index of $\mathcal{F}$ (as defined in [2]) along a generic separatrix. Then the topological invariance of the algebraic multiplicity of a dicritical singularity is a consequence of the topological invariance of the index along a curve, which is proved in [2]. Thus, from now on we assume that $\mathcal{F}$ is nondicritical. In this work we impose conditions on the topological equivalence $\mathrm{h} : U \to \widetilde{U}$ and prove the following.

Theorem 6.1. *Let* $\mathrm{h} : U \to \widetilde{U}$ *be a topological equivalence between* $\mathcal{F}$ *and* $\widetilde{\mathcal{F}}$ *and assume that* h *preserves the orientation of* $\mathbb{C}^2$. *Suppose that* h *is differentiable at* $0 \in \mathbb{C}^2$ *and such that* $d\,\mathrm{h}(0) : \mathbb{R}^4 \to \mathbb{R}^4$ *is a real isomorphism. Then the algebraic multiplicities of* $\mathcal{F}$ *and* $\widetilde{\mathcal{F}}$ *are the same.*

Let $\pi : \widehat{\mathbb{C}^2} \to \mathbb{C}^2$ be the blow up at $0 \in \mathbb{C}^2$. Given a complex line P passing through $0 \in \mathbb{C}^2$, we say that P is *regular for* $\mathcal{F}$, if the strict transform of P by π intersects the divisor E at a regular point of the strict transform of $\mathcal{F}$. The following theorem is a key step in the proof of Theorem 6.1.

Theorem 6.2. *Let* $\mathrm{h} : U \to \widetilde{U}$ *be a topological equivalence between* $\mathcal{F}$ *and* $\widetilde{\mathcal{F}}$ *and assume that* h *preserves the orientation of* $\mathbb{C}^2$. *Let* P *and* $\widetilde{P}$ *be two complex lines passing through* $0 \in \mathbb{C}^2$ *which are regular for* $\mathcal{F}$ *and* $\widetilde{\mathcal{F}}$ *respectively. Suppose that* $P \cap U$ *is homeomorphic to a disc and* $\mathrm{h}(P \cap U) = \mathrm{h}(\widetilde{P} \cap \widetilde{U})$. *Then the algebraic multiplicities of* $\mathcal{F}$ *and* $\widetilde{\mathcal{F}}$ *are equal.*

The paper is organized as follows. In section 7 we prove a weaker version of Theorem 6.2. In section 8 we stay and prove a topological lemma, fundamental for the following sections. We prove Theorem 6.2 in section 9. Finally, in section 10 we prove Theorem 6.1.

7 A first theorem.

Let $\mathrm{h} : U \to \widetilde{U}$ be a topological equivalence between $\mathcal{F}$ and $\widetilde{\mathcal{F}}$. Let $\mathcal{F}_0$ and $\widetilde{\mathcal{F}}_0$ be the strict transforms of $\mathcal{F}$ and $\widetilde{\mathcal{F}}$ respectively. Let W and $\widetilde{W}$ be denote the sets $\pi^{-1}(U)$ and

$\pi^{-1}(\widetilde{U})$ respectively. Let

$$h : W\backslash E \to \widetilde{W}\backslash E$$

be the homeomorphism defined by $h = \pi^{-1}\,\mathrm{h}\,\pi$. We have a natural fibration ρ on $\widehat{\mathbb{C}^2}$ which fibers are the strict transforms of the complex lines passing through $0 \in \mathbb{C}^2$. Consider $p, \widetilde{p} \in E$ and let L_p and $L_{\widetilde{p}}$ be the fibers of ρ passing through p and $\widetilde{p}$ respectively. This section is devoted to prove the following.

Theorem 7.1. *Suppose that p and $\widetilde{p}$ are regular points of $\mathcal{F}_0$ and $\widetilde{\mathcal{F}}_0$ respectively. Let Ω be a neighborhood of p in $\widehat{\mathbb{C}^2}$. Suppose that h extends to $(W\backslash E)\cup\Omega$ as a homeomorphism onto its image, such that $h(L_p \cap W) = L_{\widetilde{p}} \cap \widetilde{W}$. Then the algebraic multiplicities of $\mathcal{F}$ and $\widetilde{\mathcal{F}}$ are the same.*

Let ν be the algebraic multiplicity of $\mathcal{F}$ at 0 and let $p_1, ..., p_k$ be the singularities of $\mathcal{F}_0$ on E. We have the following relation due to Van Den Essen:

$$\sum_{i=1}^{k} \mu(\mathcal{F}_0, p_i) = \mu(\mathcal{F}, 0) - \nu^2 + \nu + 1,$$

where $\mu(\mathcal{F}, p)$ is the Milnor number of $\mathcal{F}$ at p. Let $s = \sum_{i=1}^{k} \mu(\mathcal{F}_0, p_i)$. In the same way, let $\widetilde{s}$ be the sum of the Milnor numbers of the singularities on E of $\widetilde{\mathcal{F}}_0$. Then, since the Milnor number is a topological invariant, it is sufficient to prove that $s = \widetilde{s}$.

Let $\mathcal{D} \subset E \cap \Omega$ be a closed disc containing p, which does not contain singularities of $\mathcal{F}_0$ and such that $h(\mathcal{D})$ does not contains singularities of $\widetilde{\mathcal{F}}_0$. Let D and $\widetilde{D}$ be the closed discs in E equal to the closure of $E\backslash\mathcal{D}$ and $E\backslash h(\mathcal{D})$ respectively. Then h maps $W\backslash D$ homeomorphically onto $\widetilde{W}\backslash\widetilde{D}$, and the interiors of D and $\widetilde{D}$ contain all the singularities of $\mathcal{F}_0$ and $\widetilde{\mathcal{F}}_0$ respectively. Observe that h is a topological equivalence between $\mathcal{F}_0|_{W\backslash D}$ and $\widetilde{\mathcal{F}}_0|_{\widetilde{W}\backslash\widetilde{D}}$. Since $h(L_p \cap W) = L_{\widetilde{p}} \cap \widetilde{W}$, we have the homeomorphism

$$h : (W\backslash D)\backslash L_p \to (\widetilde{W}\backslash\widetilde{D})\backslash L_{\widetilde{p}}.$$

We know that $W\backslash L_p$ and $\widetilde{W}\backslash L_{\widetilde{p}}$ are isomorphic to $\mathbb{C}^2$, where the divisor can be represented by the vertical line $\{z_1 = 0\}$ and the sets $W\backslash L_p$ and $\widetilde{W}\backslash L_{\widetilde{p}}$ give neighborhoods V and $\widetilde{V}$ of $\{z_1 = 0\}$. Thus, we may think that the foliations $\mathcal{F}_0$ and $\widetilde{\mathcal{F}}_0$ are defined on the sets V and $\widetilde{V}$ in $\mathbb{C}^2$, and that

$$h : V\backslash D \subset \mathbb{C}^2 \to \widetilde{V}\backslash\widetilde{D} \subset \mathbb{C}^2$$

is a topological equivalence between $\mathcal{F}_0$ and $\widetilde{\mathcal{F}}_0$. Observe that $\mathcal{F}_0$ is globally defined by a holomorphic vector field on V and the same holds for $\widetilde{\mathcal{F}}_0$ on $\widetilde{V}$. The disc D is contained in $\{z_1 = 0\}$ and we may assume that $D = \{(0, z_2) : |z_2| \leq r\}$, where $r > 0$.

We proceed now to compute s. Let Z be a holomorphic vector field, which generates the foliation $\mathcal{F}_0$ on V. Let B be a neighborhood of D homeomorphic to a ball, such that ∂B is homeomorphic to S^3 and $\overline{B} \subset V$. Since all the singularities of $\mathcal{F}_0$ are contained in $D \subset B$, it can be proved that the sum of the Milnor numbers of the singularities of $\mathcal{F}_0$ is equal to the degree of the map

$$\frac{Z}{||Z||} : \partial B \to \mathbb{S}^3,$$

$$\frac{Z}{||Z||}(z) = \frac{Z(z)}{||Z(z)||}.$$

Let $\mathcal{B}$ be a neighborhood of $\overline{B}$ homeomorphic to a ball and such that $\overline{\mathcal{B}} \subset V$. Since V is a neighborhood of $\{z_1 = 0\}$, for $\varepsilon > 0$ small enough, the set $\{|z_1| < 2\varepsilon, |z_2| < 4r\}$, which contains D, is contained in V. Then, we may chose B and $\mathcal{B}$ such that

$$\mathcal{B} \subset \{|z_1| < \varepsilon, |z_2| < 2r\}.$$

The last hypothesis will be used only in the proof of Lemma 7.5.

Consider the sets $\widetilde{B} = h(B \backslash D) \cup \widetilde{D}$, $\widetilde{\mathcal{B}} = h(\mathcal{B} \backslash D) \cup \widetilde{D}$ and $\widetilde{V} = h(V \backslash D) \cup \widetilde{D}$. It is easy to see that $\widetilde{B}$, $\widetilde{\mathcal{B}}$ and $\widetilde{V}$ are neighborhoods of $\widetilde{D}$ in $\mathbb{C}^2$.



Let

$$\varphi : \overline{\mathbb{D}}_\varepsilon \times \overline{\mathcal{B}} \to V \subset \mathbb{C}^2$$

and

$$\widetilde{\varphi} : \overline{\mathbb{D}}_\varepsilon \times \widetilde{\mathcal{B}} \to \widetilde{V} \subset \mathbb{C}^2$$

be the local complex flows of Z and $\widetilde{Z}$ respectively, where $\mathbb{D}_\varepsilon = \{T \in \mathbb{C} : ||T|| < \varepsilon\}$ with ε small enough. Now, we follow the ideas used in [2] to prove the topological invariance of the Milnor number.

Lemma 7.2. *There exists continuous functions $\tau : \mathcal{B} \backslash D \to (0, \varepsilon)$ and $\widetilde{\tau} : h(\mathcal{B} \backslash D) \to \mathbb{D}_\varepsilon \backslash \{0\}$ such that for all $z \in \mathcal{B} \backslash D$ we have:*

(i) $\varphi(\tau(z), z) \in \mathcal{B} \backslash D$.

(ii) $\varphi(t\tau(z), z) \neq z$, *for any* $t \in (0, 1]$.

(iii) $h(\varphi(\tau(z), z)) = \widetilde{\varphi}(\widetilde{\tau}(h(z)), h(z))$.

We say that a function $f : U \to \mathbb{R}$ is *lower(upper) semi-continuous* if given $\epsilon > 0$ and $x_0 \in U$, there is a neighborhood Ω of x_0 in U such that $f(x) \geq f(x_0) - \epsilon$ ($f(x) \leq f(x_0) + \epsilon$) for all $x \in \Omega$. We need the following lemma.

Lemma 7.3. *Let U be an open set in $\mathbb{R}^n$ and let $f : U \to \mathbb{R}$ and $g : U \to \mathbb{R}$ be an upper and a lower semicontinuous function respectively. Suppose that $f < g$. Then there exists a continuous function $h : U \to \mathbb{R}$ such that $f < h < g$. In particular, if g is a strictly positive lower semicontinuous function, then there exists a continuous function h such that $0 < h < g$.*

Proof of Lemma 7.2. Clearly, given $z \in \mathcal{B}\backslash D$ there exists $\delta > 0$ such that $\varphi(*, z)$ is injective on $\mathbb{D}_\delta$. Then define $\delta(z) > 0$ as the supreme of $\delta' \leq \varepsilon$ such that $\varphi(*, z)$ is injective on $\mathbb{D}_{\delta'}$.

Assertion 1. *The function $\delta : \mathcal{B}\backslash D \to (0, \varepsilon]$ is lower semicontinuous.*

Proof. Fix $z_0 \in \mathcal{B}\backslash D$ and let $\epsilon > 0$. We will prove that for z close enough to z_0 we have $\delta(z) \geq \delta(z_0) - \epsilon$. Suppose by contradiction that for $z_k \to z_0$ we have that $\varphi(*, z_k)$ is not injective on $\mathbb{D}_{\delta(z_0)-\epsilon}$. Then there are points t_k, t'_k in $\mathbb{D}_{\delta(z_0)-\epsilon}$, with $t_k \neq t'_k$ and such that $\varphi(t_k, z_k) = \varphi(t'_k, z_k)$ for all k. By taking a subsequence we may assume that $t_k \to a$ and $t'_k \to a'$ with $a, a' \in \overline{\mathbb{D}}_{\delta(z_0)-\epsilon} \subset \mathbb{D}_{\delta(z_0)}$. By continuity we have

$$\varphi(a, z_0) = \lim_{k\to\infty} \varphi(t_k, z_k) = \lim_{k\to\infty} \varphi(t'_k, z_k) = \varphi(a', z_0)$$

and, since $\varphi(*, z_0)$ is injective on $\mathbb{D}_{\delta(z_0)}$, we deduce that $a = a'$. Let $z' = \varphi(a, z_0)$ and take a neighborhood Ω of z' and $\delta_0 > 0$ such that $\varphi(*, z)$ is injective on $\mathbb{D}_{\delta_0}$ for all $z \in \Omega$. For k big enough we have that $\varphi(a, z_k) \in \Omega$ and $(t_k - a), (t'_k - a') \in \mathbb{D}_{\delta_0}$. Then, since

$$\varphi(t_k - a, \varphi(a, z_k)) = \varphi(t_k, z_k) = \varphi(t'_k, z_k) = \varphi(t'_k - a', \varphi(a', z_k)),$$

we have that $t_k - a = t'_k - a'$, hence $t_k = t'_k$, which is a contradiction.

Assertion 2. *Consider $\bar{\delta} : \mathcal{B}\backslash D \to (0, \varepsilon]$, where $\bar{\delta}(z)$ is the supreme of $\delta' < \varepsilon$ such that $\varphi(\mathbb{D}_{\delta'}, z) \subset \mathcal{B}\backslash D$. Then $\bar{\delta}$ is a lower semicontinuous function.*

Proof. Fix z_0 and let $\epsilon > 0$. The set $\varphi(\overline{\mathbb{D}}_{\bar{\delta}(z_0)-\epsilon}, z_0)$ is compact and is contained in $\mathcal{B}\backslash D$. If z is close enough to z_0 we have that $\varphi(\overline{\mathbb{D}}_{\bar{\delta}(z_0)-\epsilon}, z)$ is also contained in $\mathcal{B}\backslash D$. Then $\bar{\delta}(z) \geq \bar{\delta}(z_0) - \epsilon$ and it follows that $\bar{\delta}$ is lower semicontinuous.

Consider $\widetilde{\delta} : h(\mathcal{B}\backslash D) \to (0, \varepsilon]$, where $\widetilde{\delta}(w)$ is the supreme of $\delta' < \varepsilon$ such that $\widetilde{\varphi}(*, w)$ is injective on $\mathbb{D}_{\delta'}$. As in Assertion 1, we can prove that $\widetilde{\delta}$ is a lower semicontinuous function.

Assertion 3. *Define $\hat{\delta} : \mathcal{B}\backslash D \to (0, \varepsilon]$, where $\hat{\delta}(z)$ is the supreme of $\delta' < \varepsilon$ such that $h(\varphi(\mathbb{D}_{\delta'}, z))$ is contained in $\widetilde{\varphi}(\mathbb{D}_{\widetilde{\delta}(h(z))}, h(z))$. Then $\hat{\delta}$ is a lower semicontinuous function.*

Proof. Fix z_0 and let $\epsilon > 0$. Since $h(\varphi(\mathbb{D}_{\hat{\delta}(z_0)}, z_0))$ is contained in $\widetilde{\varphi}(\mathbb{D}_{\widetilde{\delta}(h(z_0))}, h(z_0))$, there is $\epsilon' > 0$ such that $h(\varphi(\overline{\mathbb{D}}_{\hat{\delta}(z_0)-\epsilon}, z_0))$ is contained in $\widetilde{\varphi}(\mathbb{D}_{\widetilde{\delta}(h(z_0))-\epsilon'}, h(z_0))$. Let Σ be a disc passing through $h(z_0)$ and transverse to the foliation. Since $\widetilde{\delta}$ is lower semicontinuous, we may take Σ small enough such that $\widetilde{\varphi}(*, z)$ is injective on $\overline{\mathbb{D}}_{\widetilde{\delta}(h(z_0))-\epsilon'}$ for all $z \in \Sigma$. Moreover, we may take Σ small enough such that $\widetilde{\varphi}$ is injective on

$\mathbb{D}_{\widetilde{\delta}(h(z_0))-\epsilon'} \times \Sigma$. Let M denote the open set $\widetilde{\varphi}(\mathbb{D}_{\widetilde{\delta}(h(z_0))-\epsilon'} \times \Sigma)$ and let $M' = \widetilde{\varphi}(\mathbb{D}_{\epsilon'/2} \times \Sigma)$. We may take a neighborhood Ω of z_0 such that $h(\Omega) \subset M'$ and $\widetilde{\delta}(h(z)) \geq \widetilde{\delta}(h(z_0)) - \epsilon'/2$ for all $z \in \Omega$, because $\widetilde{\delta}$ is lower semicontinuous. Since $h(\varphi(\overline{\mathbb{D}}_{\hat{\delta}(z_0)-\epsilon}, z_0))$ is compact and is contained in M, we may assume Ω small enough such that $h(\varphi(\overline{\mathbb{D}}_{\hat{\delta}(z_0)-\epsilon}, z))$ is contained in M for all $z \in \Omega$. Fix $z \in \Omega$. Since $h(z) \in M'$, there is $w' \in \Sigma$ and t', with $|t'| < \epsilon'/2$, such that $h(z) = \widetilde{\varphi}(t', w')$. Since $h(\varphi(\overline{\mathbb{D}}_{\hat{\delta}(z_0)-\epsilon}, z))$ is contained in M, we deduce that it is contained in $\widetilde{\varphi}(\mathbb{D}_{\widetilde{\delta}(h(z_0))-\epsilon'}, w')$. Then, given w in $h(\varphi(\overline{\mathbb{D}}_{\hat{\delta}(z_0)-\epsilon}, z))$, we have that $w = \widetilde{\varphi}(t'', w')$ with $|t''| < \widetilde{\delta}(h(z_0)) - \epsilon'$. Thus

$$w = \widetilde{\varphi}(t'', w') = \widetilde{\varphi}(t'' - t', \widetilde{\varphi}(t', w')) = \widetilde{\varphi}(t'' - t', h(z)),$$

where $|t'' - t'| \leq |t''| + |t'| < \widetilde{\delta}(h(z_0)) - \epsilon' + \epsilon'/2 = \widetilde{\delta}(h(z_0)) - \epsilon'/2 \leq \widetilde{\delta}(h(z))$. Then $h(\varphi(\overline{\mathbb{D}}_{\hat{\delta}(z_0)-\epsilon}, z))$ is contained in $\widetilde{\varphi}(\mathbb{D}_{\widetilde{\delta}(h(z))}, h(z))$ and it follows that $\hat{\delta}$ is lower semicontinuous.

It is easy to see that the function $g = \min\{\delta, \bar{\delta}, \hat{\delta}\}$ is also lower semicontinuous. Then, by Lemma 7.3, there exists a positive continuous function τ on $\mathcal{B}\backslash D$ such that $\tau < \delta, \bar{\delta}, \hat{\delta}$. By the definition of $\bar{\delta}$, (i) is satisfied. Since $\varphi(*, z)$ is injective on $\mathbb{D}_{\bar{\delta}}$ and $\tau(z) \in \mathbb{D}_{\bar{\delta}}$, we have that (ii) holds. Now, we shall define $\widetilde{\tau}$. Let $w = h(z) \in h(\mathcal{B}\backslash D)$. Since $\tau < \hat{\delta}$, we have that $h(\varphi(\tau(z), z))$ is contained in $\widetilde{\varphi}(\mathbb{D}_{\widetilde{\delta}(h(z))}, h(z))$ and by injectivity there exists a unique $\widetilde{\tau}(h(z))$ in $\mathbb{D}_{\widetilde{\delta}(h(z))}$ such that $h(\varphi(\tau(z), z)) = \widetilde{\varphi}(\widetilde{\tau}(h(z)), h(z))$. Now, it is easy to see that $\widetilde{\tau}$ is continuous and therefore (iii) holds. $\square$

Proof of Lemma 7.3. Consider $x \in U$ and $a_x \in \mathbb{R}$, such that $f(x) < a_x < g(x)$. It follows from the definition of lower and upper semicontinuous function that there exists a neighborhood V_x of x in U such that $f(y) < a_x < g(y)$ for all $y \in V_x$. We may take a subset $I \subset U$, such that $U \subset \bigcup_{i \in I} V_i$ and $\{V_i\}_{i \in I}$ is locally finite. Thus, we have $f(x) < a_i < g(x)$ for all $x \in V_i$. Let $\{\psi_i\}_{i \in I}$ be a partition of the unity subordinate to $\{V_i\}_{i \in I}$. Then, we define $h : U \to \mathbb{R}$ by

$$h(x) = \sum_{i \in I} \psi_i(x) a_i.$$

Clearly, h is continuous. If $x \in V_i$, then $f(x) < a_i < g(x)$, hence $\psi_i(x) f(x) < \psi_i(x) a_i < \psi_i(x) g(x)$ and it follows that $f < h < g$. $\square$

From Lemma 7.2, we have the maps

$$f : \mathcal{B}\backslash D \to \mathcal{B}\backslash D,$$

$$f(z) = \varphi(\tau(z), z)$$

and

$$\widetilde{f} : \widetilde{\mathcal{B}}\backslash \widetilde{D} \to \widetilde{\mathcal{B}}\backslash \widetilde{D},$$

$$\widetilde{f}(w) = \widetilde{\varphi}(\widetilde{\tau}(w), w)$$

with

$$h \circ f = \widetilde{f} \circ h$$

and such that f and $\widetilde{f}$ are without fixed points.

There exists $\psi, \widetilde{\psi} : \mathbb{C}^2 \to \mathbb{C}^2$ with the following properties:

(i) $\psi(D) = 0$ and $\widetilde{\psi}(\widetilde{D}) = 0$.

(ii) $\psi : \mathbb{C}^2 \backslash D \to \mathbb{C}^2 \backslash \{0\}$ and $\widetilde{\psi} : \mathbb{C}^2 \backslash \widetilde{D} \to \mathbb{C}^2 \backslash \{0\}$ are homeomorphisms.

(iii) ψ and $\widetilde{\psi}$ are equal to the identity out of B and $\widetilde{B}$ respectively.

We define

$$\begin{array}{rcl} f' & = & \psi f \psi^{-1} : \mathcal{B}\backslash\{0\} \to \mathcal{B}\backslash\{0\} \subset \mathbb{C}^2, \\ \widetilde{f}' & = & \widetilde{\psi} \widetilde{f} \widetilde{\psi}^{-1} : \widetilde{\mathcal{B}}\backslash\{0\} \to \widetilde{\mathcal{B}}\backslash\{0\} \subset \mathbb{C}^2, \\ h' & = & \widetilde{\psi} h \psi^{-1} : V \to \widetilde{V}. \end{array}$$

Then we have the following:

(i) f' and $\widetilde{f}'$ do not have fixed points.

(ii) On ∂B, we have $f' = f$ and $\widetilde{f}' = \widetilde{f}$.

(iii) h' is a homeomorphism with $h'(0) = 0$ and such that $h' \circ f' = \widetilde{f}' \circ h'$.

Thus, there are well defined maps:

$$\begin{array}{rcl} (f' - \mathrm{id}) & : & \mathcal{B}\backslash\{0\} \to \mathbb{C}^2\backslash\{0\}, \\ (\widetilde{f}' - \mathrm{id}) & : & \widetilde{\mathcal{B}}\backslash\{0\} \to \mathbb{C}^2\backslash\{0\}. \end{array}$$

Observe that $H_3(\mathcal{B}\backslash\{0\}) \subset H_3(\mathbb{C}^2\backslash\{0\})$ and this inclusion is an isomorphism between the groups. Then $(f' - \mathrm{id})$ induces a map

$$(f' - \mathrm{id})_* : H_3(\mathbb{C}^2\backslash\{0\}) \to H_3(\mathbb{C}^2\backslash\{0\})$$

at the homology level.

Lemma 7.4. $(f' - \mathrm{id})_*$ *is the multiplication by* s.

Proof. We have that $\partial B \subset \mathcal{B}$ is a generator of $H_3(\mathbb{C}^2\backslash\{0\})$. It is known that, homologically:

$$(f' - \mathrm{id})(\mathbb{S}^3) = (f' - \mathrm{id})(\partial B) = n\mathbb{S}^3,$$

where n is the degree of the map:

$$g : \partial B \to \mathbb{S}^3,$$

$$g(z) = \frac{(f' - \mathrm{id})}{||(f' - \mathrm{id})||}(z).$$

Thus, it is sufficient to prove that $\deg(g) = s$. Observe that $g = \frac{(f-\mathrm{id})}{||(f-\mathrm{id})||}$, since $f' = f$ on ∂B. By (ii) of Lemma 7.2 the map

$$G : [0,1] \times \partial B \to \mathbb{S}^3,$$

$$G(t,z) = \frac{\varphi(t\tau(z),z) - z}{||\varphi(t\tau(z),z) - z||}, \quad t \neq 0,$$

$$G(0,z) = \frac{\tau(z)}{||\tau(z)||} \cdot \frac{Z(z)}{||Z(z)||}$$

is well defined. Evidently, $G(1,z) = g(z)$. On the other hand:

$$\begin{aligned}
\lim_{t\to 0} G(t,z) &= \frac{\tau(z)}{||\tau(z)||} \lim_{t\to 0} \left|\left| \frac{\varphi(t\tau(z),z) - z}{t\tau(z)} \right|\right|^{-1} \cdot \lim_{t\to 0} \frac{\varphi(t\tau(z),z) - z}{t\tau(z)} \\
&= \frac{\tau(z)}{||\tau(z)||} \lim_{s\to 0} \left|\left| \frac{\varphi(s,z) - z}{s} \right|\right|^{-1} \cdot \lim_{s\to 0} \frac{\varphi(s,z) - z}{s} \\
&= \frac{\tau(z)}{||\tau(z)||} \cdot \frac{Z(z)}{||Z(z)||}.
\end{aligned}$$

It follows that G is continuous and therefore is a homotopy between $g(z)$ and $G(0,z) = \frac{\tau(z)}{||\tau(z)||} \cdot \frac{Z(z)}{||Z(z)||}$. Now, since $\pi_3(\mathbb{S}^1) = \{0\}$, the map $\tau/|\tau| : \partial B \to \mathbb{S}^1$ is homotopic to the constant $1 \in \mathbb{S}^1$ and g is homotopic to $Z/||Z||$. Therefore $\deg(g) = \deg(Z/||Z||) = s$.

In the same way, we have that

$$(\widetilde{f}' - \mathrm{id})_* : H_3(\mathbb{C}^2\backslash\{0\}) \to H_3(\mathbb{C}^2\backslash\{0\})$$

is the multiplication by $\widetilde{s}$.

Let

$$h'_* : H_3(\mathbb{C}^2\backslash\{0\}) \to H_3(\mathbb{C}^2\backslash\{0\})$$

be the isomorphism induced by h'. Clearly, the following lemma implies Proposition 6.2.

Lemma 7.5. *The following diagram commutes:*

$$\begin{array}{ccc}
H_3(\mathbb{C}^2\backslash\{0\}) & \xrightarrow{(f'-\mathrm{id})_*} & H_3(\mathbb{C}^2\backslash\{0\}) \\
\downarrow{\scriptstyle h'_*} & & \downarrow{\scriptstyle h'_*} \\
H_3(\mathbb{C}^2\backslash\{0\}) & \xrightarrow{(\widetilde{f}'-\mathrm{id})_*} & H_3(\mathbb{C}^2\backslash\{0\})
\end{array}$$

Proof. Recall that $\mathcal{B}$ was chosen such that

$$\mathcal{B} \subset \{|z_1| < \varepsilon, |z_2| < 2r\} \subset \{|z_1| < 2\varepsilon, |z_2| < 4r\} \subset V.$$

Since $h' \circ f' = \widetilde{f}' \circ h'$ we have $(\widetilde{f}' - \mathrm{id}) \circ h' = \widetilde{f}' \circ h' - h' = h' \circ f' - h'$. It is sufficient to prove that $h' \circ f' - h'$ and $h' \circ (f' - \mathrm{id}) : \mathcal{B}\backslash\{0\} \to \mathbb{C}^2\backslash\{0\}$ are homotopic. For any

$z \in \mathcal{B}\backslash\{0\}$ and $t \in [0,1]$ we have that $f'(z), (1-t)z \in \mathbb{D}_\epsilon \times \mathbb{D}_{2r}$. Then $(f'(z) + (1-t)z)$ is contained in $\mathbb{D}_{2\epsilon} \times \mathbb{D}_{4r} \subset V$. Therefore, the map:

$$F : [0,1] \times (\mathcal{B}\backslash\{0\}) \to \mathbb{C}^2\backslash\{0\},$$

$$F(t,z) = h'(f'(z) - (1-t)z) - h'(tz)$$

is well defined. F is continuous and $F(t,z) \neq 0$ for all $(t,z) \in [0,1] \times (\mathcal{B}\backslash\{0\})$ because $F(t,z) = 0$ implies $h'(f'(z) - (1-t)z) = h'(tz)$ and since h' is a homeomorphism $f'(z) - (1-t)z = tz$, hence $f'(z) = z$, which contradicts $f'(z) \neq z$. Thus F is a homotopy between $h' \circ f' - h'$ and $h' \circ (f' - \mathrm{id})$. $\square$

8 A topological fact.

Let $\pi : \widehat{\mathbb{C}^2} \to \mathbb{C}^2$ be the blow up at $0 \in \mathbb{C}^2$ and let $E = \pi^{-1}(0)$. Let $\rho : \widehat{\mathbb{C}^2} \to E$ be the natural projection. If $x \in E$, the set $\rho^{-1}(x)$ is the strict transform of a complex line passing through $0 \in \mathbb{C}^2$ and $x \in \rho^{-1}(x)$. Let M be a complex manifold. We say that $\mathcal{D}$ is a complex disc in M, if $\mathcal{D} \subset M$ and there is a map $f : \overline{\mathbb{D}} \to M$, which is a homeomorphism onto $\mathcal{D}$ and is holomorphic on $\mathbb{D}$. Let V be any subset of M containing $\partial\mathcal{D}$. The map $f|_{S^1} : S^1 \to \partial\mathcal{D} \subset M$ defines a 1-cycle in V and represents an element in $H_1(V)$ which does not depend on f. We denote this 1-cycle by $\partial\mathcal{D}$ independly of the set V. For simplicity, we write $\gamma = \gamma'$ in $H_1(M)$ for means that the 1-cycles γ and γ' represents the same element in the group $H_1(M)$. The following Lemma is a reason for assuming that the topological equivalence h preserves the orientation of $\mathbb{C}^2$.

Lemma 8.1. *Let* $\mathrm{h} : U \to U'$ *be a homeomorphism, where* U *and* U' *are neighborhoods of* $0 \in \mathbb{C}^2$ *homeomorphic to balls. Let* P *and* P' *be two complex lines passing through* $0 \in \mathbb{C}^2$*. Suppose that* $P \cap U$ *is homeomorphic to a disc and* $\mathrm{h}(P \cap U) = P' \cap U'$*. Let* L *and* L' *be the strict transforms of* P *and* P' *respectively. Let* p *and* p' *be the points of intersection of* L *and* L' *with* E *respectively. Denote by* W *and* W' *the sets* $\pi^{-1}(U)$ *and* $\pi^{-1}(U')$ *in* $\widehat{\mathbb{C}^2}$ *and let* $h : W\backslash E \to W'\backslash E$ *be the homeomorphism defined by* $h = \pi^{-1}\,\mathrm{h}\,\pi$*. Let* $V \subset W$ *be a neighborhood of* p *and let*

$$\varphi : \mathbb{D} \times \mathbb{D} \to V$$

be a biholomorphism such that $\varphi(\{0\} \times \mathbb{D}) = L \cap V$ *and* $\varphi(\mathbb{D} \times \{0\}) = E \cap V$*. Let* r *with* $0 < r < 1$ *and consider the disc* $\mathcal{B}_w = \varphi(w, |z| \leq r)$*, where* $w \in \mathbb{D}$*. Let* Ω *be a neighborhood of* p' *in* E*, homeomorphic to a disc. Let* $V' \subset \widehat{\mathbb{C}^2}$ *be the set* $\rho^{-1}(\Omega)$*. Let* $\mathcal{A}' \subset V'\backslash E$ *and* $\mathcal{B}' \subset V'\backslash L'$ *be complex discs transverse to* L' *and* E *respectively. Then, for* $|w|$ *small enough we have the following:*

(i) If h *preserves the orientation of* $\mathbb{C}^2$*, then*

$$h(\partial\mathcal{B}_w) = \xi\partial\mathcal{B}' \quad \textit{in} \quad H_1(V'\backslash(L' \cup E)),$$

where $\xi = +1$ *or* -1*.*

(ii) If h *inverts the orientation of* $\mathbb{C}^2$*, then*

$$h(\partial\mathcal{B}_w) = -2\xi\partial\mathcal{A}' + \xi\partial\mathcal{B}' \quad in \quad H_1(V'\backslash(L' \cup E)),$$

where $\xi = +1$ *or* -1*.*

Remark. With some hypothesis on the foliation $\mathcal{F}$, we have in fact that the topological equivalence h necessarily preserves the orientation of $\mathbb{C}^2$. Precisely, we have the following.

Proposition 8.2. *Let* $\mathcal{F}$ *be a holomorphic foliation by curves on* U *which has* $0 \in \mathbb{C}^2$ *as its unique singularity. Suppose that* $\mathcal{F}$ *has three smooth and transverse separatrices. Suppose that* $\widetilde{\mathcal{F}}$ *is another holomorphic foliation of a neighborhood* $\widetilde{U}$ *of* $0 \in \mathbb{C}^2$ *and let*

$$\mathrm{h} : U \to \widetilde{U}$$

be a topological equivalence between $\mathcal{F}$ *and* $\widetilde{\mathcal{F}}$*. Then* h *preserves the orientation of* $\mathbb{C}^2$*.*

Let $U \subset \mathbb{C}^2$ be an open set homeomorphic to a ball. Let P be a complex line in $\mathbb{C}^2$ and suppose that $U \cap P$ is homeomorphic to a disc. It follows by Alexander's duality theorem that $H_1(U\backslash P) \simeq \mathbb{Z}$. Let $\mathcal{D} \subset \mathbb{C}^2$ be a complex disc transverse to P. The 1-cycle $\partial\mathcal{D}$ represents an element in $H_1(U\backslash P) \simeq \mathbb{Z}$, which does not depends on the disc $\mathcal{D}$. We know that $\partial\mathcal{D}$ is a generator of the group and we say that it is the *positive generator of* $H_1(U\backslash P)$. Given a homeomorphism $f : M \to M'$, where M and M' are oriented manifolds, we define $\deg(f)$ to be 1 or -1 depending on whether f preserves or reverses orientation.

Lemma 8.3. *Let* $\mathrm{h} : U \to U'$ *be a homeomorphism, where* U *and* U' *are neighborhoods of* $0 \in \mathbb{C}^2$ *homeomorphic to balls. Let* P *and* P' *be two complex lines passing through* $0 \in \mathbb{C}^2$*. Suppose that* $P \cap U$ *is homeomorphic to a disc and* $\mathrm{h}(P \cap U) = P' \cap U'$*. Let* a *and* a' *be 1-cycles in* $U\backslash P$ *and* $U'\backslash P'$ *representing the positive generators of* $H_1(U\backslash P)$ *and* $H_1(U'\backslash P')$ *respectively. Then*

$$\mathrm{h}(a) = \deg(\mathrm{h})\deg(\mathrm{h}\,|_P)a' \quad in \quad H_1(U'\backslash P').$$

Proof of Lemma 8.1. If $\mathcal{B}'' \subset V'\backslash L'$ is any complex disc transverse to E, we have that $\partial\mathcal{B}''$ is homologous $\partial\mathcal{B}'$ in $H_1(V'\backslash(L' \cup E))$. Thus, we may change the disc $\mathcal{B}'$ if necessary and assume that it is contained in W'. Let b' be the 1-cycle defined by $b' = \pi(\partial\mathcal{B}')$. Then, since $\pi(\mathcal{B}') \subset U'$ is a complex disc transverse to P' and $\pi(\partial\mathcal{B}') = \partial\pi(\mathcal{B}')$, we have that b' is a positive generator of $H_1(U'\backslash P')$. Analogously, if $b = \pi(\partial\mathcal{B}_w)$, we deduce that b is a positive generator of $H_1(U\backslash P)$. It follows from Lemma 8.3 that:

$$\mathrm{h}(b) = \psi\xi b' \quad \text{in} \quad H_1(U'\backslash P'),$$

where $\psi = \deg(\mathrm{h})$ and $\xi = \deg(\mathrm{h}\,|_P)$. Then, since $\pi^{-1} : U'\backslash P' \to W'\backslash(L' \cup E)$ is well defined, we have that

$$\pi^{-1}(\mathrm{h}(b)) = \psi\xi\pi^{-1}(b') \quad \text{in} \quad H_1(W'\backslash(L' \cup E))$$

and thus

$$h(\partial\mathcal{B}_w) = \psi\xi\partial\mathcal{B}' \quad \text{in} \quad H_1(W'\backslash(L' \cup E)). \tag{12}$$

Observe that $\pi(\mathcal{A}')$ is a complex disc transverse to P'. Then the cycle $\partial\pi(\mathcal{A}') = \pi(\partial\mathcal{A}')$ represents the positive generator of $H_1(U'\backslash P')$. Thus, we deduce that $\pi(\partial\mathcal{A}') = \pi(\partial\mathcal{B}')$ in $H_1(U'\backslash P')$ and therefore

$$\partial\mathcal{A}' = \partial\mathcal{B}' \quad \text{in} \quad H_1(W'\backslash(L' \cup E)). \tag{13}$$

Let $\mathcal{C}$ be the disc $\varphi(0, |z| \leq r)$ in L. Let $\mathcal{C}'$ be a disc in L' containing p'. Since h maps $\mathcal{C}$ homeomorphically into L' with $h(p) = p'$, the cycle $h(\partial\mathcal{C})$ is a generator of the group $H_1(L'\backslash\{p'\})$ and we have $h(\partial\mathcal{C}) = \deg(h|_L)\partial\mathcal{C}'$. Thus, since $h|_L$ preserves orientation if an only if h$|_P$ does, we have that $h(\partial\mathcal{C}) = \xi\partial\mathcal{C}'$ in $H_1(L'\backslash\{p'\})$. Since $L'\backslash\{p'\}$ is contained in $V'\backslash E$, we conclude that

$$h(\partial\mathcal{C}) = \xi\partial\mathcal{C}' \quad \text{in} \quad H_1(V'\backslash E). \tag{14}$$

Observe that $\partial\mathcal{C}' = \partial\mathcal{B}'$ in $H_1(V'\backslash E)$. Moreover, $\partial\mathcal{C} = \varphi(0, |z| = r)$ is homologous to $\partial\mathcal{B}_w = \varphi(w, |z| = r)$ in the set $T = \varphi(|z| \leq |w|, |z| = r)$. It is easy to see that for $|w|$ small enough, the set $h(T)$ is contained in $V'\backslash E$. Then $h(\partial\mathcal{C})$ and $h(\partial\mathcal{B}_w)$ are homologous in $V'\backslash E$. It follows from (14) and the observations above that for $|w|$ small enough:

$$h(\partial\mathcal{B}_w) = \xi\partial\mathcal{B}' \quad \text{in} \quad H_1(V'\backslash E). \tag{15}$$

We know that there exists integers n and m such that

$$h(\partial\mathcal{B}_w) = n\partial\mathcal{A}' + m\partial\mathcal{B}' \quad \text{in} \quad H_1(V'\backslash(L' \cup E)).$$

Then, since $V'\backslash(L' \cup E) \subset V'\backslash E$:

$$h(\partial\mathcal{B}_w) = n\partial\mathcal{A}' + m\partial\mathcal{B}' \quad \text{in} \quad H_1(V'\backslash E),$$

hence

$$h(\partial\mathcal{B}_w) = m\partial\mathcal{B}' \quad \text{in} \quad H_1(V'\backslash E),$$

because $\partial\mathcal{A}' = 0$ in $H_1(V'\backslash E)$. From this and (15) we conclude that $m = \xi$. Then

$$h(\partial\mathcal{B}_w) = n\partial\mathcal{A}' + \xi\partial\mathcal{B}' \quad \text{in} \quad H_1(V'\backslash(L' \cup E))$$

and, since $V'\backslash(E \cup L')$ is contained in $W'\backslash(E \cup L')$, we have that

$$h(\partial\mathcal{B}_w) = n\partial\mathcal{A}' + \xi\partial\mathcal{B}' \quad \text{in} \quad H_1(W'\backslash(L' \cup E)). \tag{16}$$

From (13) we have $\partial\mathcal{A}' = \partial\mathcal{B}'$ in $H_1(W'\backslash(L' \cup E))$. Replacing in (16) we obtain:

$$h(\partial\mathcal{B}_w) = n\partial\mathcal{B}' + \xi\partial\mathcal{B}' \quad \text{in} \quad H_1(W'\backslash(L' \cup E)).$$

Thus, from (12) we have:

$$\psi\xi\partial\mathcal{B}' = n\partial\mathcal{B}' + \xi\partial\mathcal{B}' \quad \text{in} \quad H_1(W'\backslash(L' \cup E))$$

and therefore $n = (\psi - 1)\xi$. This proves the Lemma. □